

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number  
WO 02/46439 A2(51) International Patent Classification: C12N 15/82,  
15C9, 5/10, 9/12, A01H 5/00, G01N 33/53

(21) International Application Number: PCT/EP01/14154

(22) International Filing Date: 4 December 2001 (04.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00870288.8 4 December 2000 (04.12.2000) EP  
60/256,204 15 December 2000 (15.12.2000) US(71) Applicant (for all designated States except US): UNI-  
VERSITEIT UTRECHT [NL/NL]; Heidelberglaan 8,  
NL-3584 CS Utrecht (NL).

(72) Inventors: and

(75) Inventors/Applicants (for US only): SCHERES, Ben  
[NL/NL]; Vlieland 106, NL-3524 AC Utrecht (NL).  
HEIDSTRA, Renze [NL/NL]; Hellendoornseberg 13,  
NL-3524 LJ Utrecht (NL).(74) Agents: DE CLERCQ, Ann et al.; De Clercq, Brants &  
Partners, l.; Gevaertdreef 10a, B-9830 Sint-Martens-Laleim  
(BE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

## Published:

- without international search report and to be republished upon receipt of that report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A NOVEL ROOT SPECIFIC PROMOTER DRIVING THE EXPRESSION OF A NOVEL LRR RECEPTOR-LIKE KINASE

(57) Abstract: The present invention relates to the field of plant molecular biology, more particularly to the root-specific gene expression in plants. The invention provides nucleic acids for a novel transcriptional regulatory root-specific promoter and nucleic acid and protein sequences coding for a new LRR receptor-kinase protein, further specified as a root clavata 1 homolog (RCH1). Further provided are compositions comprising nucleic acids, polypeptides, antibodies and vectors. The invention further provides for methods for modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology comprising the modification of expression in particular cells, tissues or organs of a plant of the novel LRR receptor-like kinase or comprising the expressing of a gene of interest under the control of the novel transcriptional regulatory root-specific promoter. Further are provided compounds interacting with the new polypeptides for use as herbicides or growth regulators.

## A NOVEL ROOT SPECIFIC PROMOTER DRIVING THE EXPRESSION OF A NOVEL LRR RECEPTOR-LIKE KINASE

### FIELD OF THE INVENTION

5 The present invention relates to the field of plant molecular biology, more particularly to the root-specific gene expression in plants. The isolation of a root specific operon, comprising a transcriptional regulatory promoter that contributes to root-specific gene expression and its operably linked gene encoding a novel Leucine Rich Repeat (LRR) receptor-like kinase, is disclosed. Said transcriptional regulatory promoter may be used  
10 for driving root-specific expression of at least one gene of interest in a transgenic plant and said encoded LRR receptor-like kinase gene may be used to alter the features and/or to confer a selective advantage to transgenic plants.

### BACKGROUND TO THE INVENTION

#### 15 Plant promoters

Initiation of transcription is generally understood to be the predominant controlling factor in determining expression of a gene. The transcriptional control elements, which may interact with DNA binding proteins, are generally embedded in the sequence 5'-flanking or upstream of the transcribed gene. These DNA sequence elements promote the  
20 formation of transcriptional regulatory complexes that either activate or repress the expression of the gene 3' downstream of the promoter.

The sequences of the promoter vary in length and base pair composition from gene to gene. Within the sequence of the promoter, many regulatory sequence elements can be embedded that control the promoter activity. These specific elements, called DNA-  
25 boxes, contribute to the promoter a defined feature or a defined activation pattern. Many of these boxes are recognition sites to which regulatory transcription factors can bind and are part of a tight control mechanism of the promoter. Numerous plant promoter DNA boxes have been described, such as tissue-specific boxes (Chaubet et al. 1996), pyrimidine boxes (Gubler & Jacobsen 1992), which influence the level of expression or G-boxes (Dolferus et al. 1994) which reduce promoter activity by cold or dehydration.

30

Constitutive promoters are promoters that drive the expression of genes in a continuous manner, whereas ubiquitous promoters drive the expression of the gene throughout the entire plant.

5 Expression of heterologous DNA sequences in a host is dependent upon the presence of an operably linked promoter that is also present in that host, whereby the choice of promoter will determine when, where or how strong the heterologous DNA sequence will be expressed in the host organism. Frequently those skilled in the art desire to limit the expression of the transfected DNA sequence to a certain time-period (e.g. to a particular phase of plant development), to a defined tissue of the host (e.g. leaves), or to a certain  
10 expression level. Controlling the expression of the heterologous genes in transgenic plants is considered to provide several advantages to the plant over ubiquitous and constitutive expression. There is a broad need in many industrial applications for transcriptional control elements capable of driving tissue specific gene-expression in plants. It is therefore important that such control elements are continuously provided and  
15 that new control elements with their specific transcriptional features are isolated and characterized.

### Promoter regulation

Promoters from both monocots and dicots that have a root root-preferred expression  
20 pattern have already been described (Fowler, Bernhardt, & Tierney 1999; Hwang & Goodman 1995; Lagarde et al. 1996; Okada et al. 2000; Okumura et al. 1994; Pysh et al. 1999; Schneider et al. 1998; Vidmar et al. 2000; Wanapu & Shinmyo 1996; Winicov 2000; WO9007001 (Baszczynski et al. 2000; Borg et al. 1997; Masuda, Sakuta, & Satoh 1999; Webb et al. 2000) (de Pater & Schilperoort 1992; Ely, Evans, & Schuch  
25 2000; Goddemeier, Wulff, & Feix 1998; Kikuchi et al. 1999; Suzuki, Fowler, & Tierney 1993; Unbe et al. 1998; Woo & Hawes 1997; Xu et al. 1995) (Yamamoto et al. 1991) (Gritsch et al. 1997; Keller & Lamb 1989; Lauter 1996; Olson, Oetiker, & Yang 1995; van der Zaaij et al. 1991). Many of them are root-preferred rather than root-specific and are expressed at low levels in tissues other than the root. These promoters are not suitable  
30 for industrial applications where it is of the utmost importance that the heterologous product is expressed in the root only. As an example, ectopic expression of cell cycle control elements to obtain growth arrest in some parts of the root (e.g. lateral roots),

should be strictly regulated to that tissue in order not to interfere with the overall growth of the plant. In contrast, other promoters are strictly and exclusively confined to root-specific expression and can therefore be valuable for some applications demanding tight control of the expression. In this group of root-specific promoter subgroups can be distinguished based on the root-tissue they are expressed in. The HRGPnt3 gene of tobacco for instance is only expressed in the pericycle and the endodermis of the root, more specifically in the discrete subset of cells, involved in the initiation of the lateral roots (Keller et al. 1989). This makes the promoter of HRGPnt3 highly specific. Another example is the XSP30 sequence, which is only activated in the xylem of the root.

For persons skilled in the art it is of importance to have at their disposal a wide variety of root-specific promoters with different root tissue-specificity from which they can choose to confer a specific feature to the transgenic plant.

Many of these root-preferred promoters are not only limited to tissue specific expression, but are also subjected to a tight control mechanism deciding when the promoter should be active. This decision depends on the function and the effects of the gene product that is switched on/off by the related promoter. Some gene products are switched on/off at a certain developmental stage of the plant, at a certain moment of the cell cycle, in stress situations originating from weather factors, upon injury caused by pathogens or upon a change in the environmental resources. A typical example is the ARSK1 gene of *Arabidopsis thaliana*, that is activated when the roots are exposed to air during growth (dehydration) or by treatment of roots with abscisic acid or salt (Hwang et al. 1995).

However, for many industrial and agronomic applications, where the aim is to confer a special feature at any time of the developmental, environmental or physical state of the plant, a constitutive promoter will be needed. For example when the gene product of the transgene is to be isolated and purified from the plant for commercial purposes.

It is important for the persons skilled in the art to have at their disposal a wide variety of promoters with different time-control features from which they can chose to confer a specific feature to the transgenic plant.

## Plant root development

Root development is an essential determinant of plant growth and crop yield since the root is the main channel to extract nutrients from the environment. Any changes in the



root that contribute to more functional organ tissue will result in higher crop yield. Also any regulatory element that is involved in the root development and whose control is subjected to developmental events can contribute better yield properties to a transgenic plant.

- 5 Cells of the shoot and root meristems are mitotically active and they extend the undifferentiated tissue pattern of the embryo as a pool of cells from which cells can differentiate into several tissues of the adult plant. This process continues throughout the life span of the plant. It is clear that the root meristem plays a pivotal role in the formation of strong roots and therefor also in affecting the growth and crop yield.
- 10 In root meristem, the emerging picture is that the integrity of the root meristem is kept by balancing cell proliferation and cell differentiation, and that differentiation-inhibiting signals originating from the quiescent center are involved (van den Berg et al. 1997). Inhibition of differentiation only occurs in cells that contact the quiescent center, which reveals that short-range or contact-dependent signals are involved in this process.
- 15 However which compounds play a role in this process and whether positional cues guiding root cell fate are targeted via plasmodesmata or via the cell wall remains unknown (van den Berg et al. 1997).

- In the shoot meristem more progress has been made in characterizing molecular components involved in meristem maintenance. *Clavata* (*clv*) mutants accumulate
- 20 massive pools of undifferentiated cells in the central zone of shoot and floral meristem. The *CLAVATA* genes show a shoot and floral meristem specific expression pattern. Root meristems of *clv*-mutants are unaffected (Clark, Running, & Meyerowitz 1993). The *CLAVATA* loci (*CLV1*, *CLV2* and *CLV3*) appear to promote the transition toward differentiation of cells at the shoot and floral meristems, and/or to restrict the proliferation
- 25 of cells at the center of these meristems (Clark et al. 1993). *CLV1* acts with *CLV3* and *WUSCHEL* (*WUS*) and *SHOOT MERISTEMLESS* (*STM*) to maintain the integrity of the central zone which acts as a reservoir of stem cells. A model for stem-cell maintenance in the root meristem and in the shoot meristem is depicted in Figure 1.

### 30 **CLV1 homologs and CLV1-redundancy**

The *CLAVATA1* gene resides on the *Arabidopsis thaliana* chromosome 1. *Clv1* mRNA is found in the L3 tunica cell layer of the central zone and encodes a protein with leucine-rich repeats (LRR, of a type found in a number of proteins, notably the products of

several plant-pathogen resistance genes), a putative serin/threonin kinase domain and a predicted extracellular receptor domain. Expression of the *clv1* gene is initiated in the heart stage of the embryo, when cotyledonary primordia are apparent, and is independent of the *STM* (*SHOOT MERISTEMLESS*) activity (Long & Barton 1998).

5 *CLAVATA1* is a leucine-rich receptor-like kinase. The *Arabidopsis thaliana* genome project has so far identified over 50 LRR transmembrane receptor serine/threonine kinases. It can be estimated that the total *Arabidopsis thaliana* genome contains in the order of 100 family members. Other members of the *CLAVATA* family are *ERECTA* (*ER*) (Torii et al. 1996), *BRASSINOLIDE INSENSITIVE* (*BRI*) (Li & Chory 1997) and *HAESA* (*HAE*) (Jinn, Stone, & Walker 2000), genes which have a function in plant development. Analogues of the *Arabidopsis thaliana* *CLAVATA* genes were also found in soybean (*Glycine max*). They were isolated from the wild type and the fasciating mutant of soybean and were designated GmCLVA1 and GmCLV1B (Yamamoto, Karakaya, & Knap 2000).

15 It is the aim of the present invention to isolate novel promoters and genes which are useful for altering features of transgenic plants and thus confer selective advantage to transgenic plants.

All the aims of the present invention have been met by the formulation of the following preferred embodiments of the current invention.

## SUMMARY OF THE INVENTION

In the present invention a DNA operon was isolated from *Arabidopsis thaliana*, which comprises a novel root-specific promoter, driving a gene encoding a Leucine Rich Repeat (LRR) receptor-like kinase. Unexpectedly, the inventors discovered that the new operon was specifically expressed in roots and more surprisingly in the meristem of the main and lateral roots, in the main and lateral root vascular tissue and in the lateral root primordia.

The surprisingly specific expression pattern, as well as the very strong expression level, makes the newly found promoter a very attractive and selective tool to drive root-specific expression of any gene of interest. The ability to target meristem cells of the root makes this promoter clearly distinct from the root-specific promoters presently known in the art. Another surprising aspect of the invention is the identification of the new LRR receptor-

like kinase gene, which is the first gene identified in roots showing homology with a *clavata*-gene. This gene is specifically expressed in roots in a specific developmental stage of the root cells (namely in the meristematic phase) and can therefore contribute agronomic interesting features to a transgenic plant when transfected herein. Therefore the root-specific promoter is further denominated as ROOT *CLAVATA* HOMOLOG 1 promoter (RCH1prom) and the LRR receptor-like kinase as RCH1.

Accordingly, the invention embodies an isolated DNA sequence with nucleotide sequence as given in SEQ ID NO 1 (Figure 5) comprising a novel root specific regulatory promoter sequence with nucleotide sequence as given in SEQ ID NO 2 (Figure 6) or SEQ ID NO 18 (Figure 13), and an isolated DNA sequence with nucleotide sequence as given in SEQ ID NO 3 (Figure 8), encoding a novel ROOT *CLAVATA* HOMOLOG 1 gene, with amino acid sequence given in SEQ ID NO 4 or SEQ ID NO 19 (Figure 9).

Furthermore the invention embodies a method for modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology comprising the modification of expression in particular cells of the root of a particular transgene operably linked to the RCH1 promoter sequence of the present invention. Provided in the current invention are methods to effect expression of any gene of interest in particular cells of the plant root. The present invention also relates to a method to effect expression of a ROOT *CLAVATA* HOMOLOG 1 protein or a homologue or a derivative thereof in a plant cell, tissue or organ. The present invention clearly extends to any plant produced by the inventive methods described herein.

## DETAILED DESCRIPTION OF THE INVENTION

One of the technical problems underlying the present invention is to provide regulatory elements that are active in the root only and that show no expression in other tissues. Additional to the highly selective and tissue-specific regulation of a promoter, other features such as very strong activity, continuous activity, and termination of activity at a certain developmental stage of the cells, would contribute an unique combination of regulatory elements to specifically regulate the expression of a gene of interest in the roots of a transgenic plant, thereby providing a means for modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology. Also the isolation of root-specific DNA sequences encoding proteins that could play a role in root

development is envisaged for use in modification of cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology.

In a first embodiment, the present invention relates to an isolated nucleic acid comprising a transcriptional regulatory root promoter sequence and/or a nucleic acid sequence  
5 encoding a novel LRR receptor-like kinase (protein) or an immunologically active and/or functional fragment of such a regulatory element or protein (kinase) selected from the group consisting of:

- (a) nucleic acid comprising at least part of the DNA sequence as given in any of  
10 SEQ ID NOs 1, 2, 3 or 18, or the complement thereof,
- (b) nucleic acid comprising the RNA sequences corresponding to at least part of any of SEQ ID NOs 1, 2, 3 or 18, or the complement thereof,
- (c) nucleic acid specifically hybridizing, preferably under stringent conditions, with the nucleotide sequence as defined in (a) or (b),
- (d) nucleic acid encoding a protein comprising the amino acid sequence as given in  
15 SEQ ID NO 4, SEQ ID NO 5 or SEQ ID NO 19,
- (e) nucleic acid which is degenerated as a result of the genetic code to a nucleotide sequence encoding a protein as given in SEQ ID NO 4, SEQ ID NO 19 or to a nucleotide sequence as defined in (a) to (d),
- (f) nucleic acid which is diverging due to the differences in codon usage between  
20 the organisms to a nucleotide sequence encoding a protein as given in SEQ ID NO 4, SEQ ID NO 19 or to a nucleic acid sequence as defined in (a) to (d),
- (g) nucleic acid encoding a protein as given in SEQ ID NO 4, SEQ ID NO 19 or as defined in (a) to (d), which is diverging due to the differences between alleles encoding a protein
- (h) nucleic acid encoding a fragment of a protein encoded by a DNA sequence as  
25 given in SEQ ID NO 1 or SEQ ID NO 3 or encoding a fragment of a protein encoded by a nucleic acid as defined in any one of (a) to (g), wherein said fragment comprises the sequence as represented in SEQ ID NO 5, and,
- (k) nucleic acid encoding a protein as defined in SEQ ID NO 4, SEQ ID NO 19 or  
30 nucleic acid as defined in any one of (a) to (h), said nucleic acid interrupted by intervening DNA sequences.

The sequence as depicted in SEQ ID NO 1 is part of a sequence deposited under the Genbank Accession number AB017061, which represents the entire *Arabidopsis*

*thaliana* chromosome 5. The nucleic acid of the invention is not the nucleic acid with the sequence as deposited under the Genbank Accession number AB017061 (Figure 14).

Furthermore, in a preferred embodiment, the present invention relates to an isolated nucleic acid comprising a novel root-specific promoter sequence, termed RCH1prom, isolated from *Arabidopsis thaliana* and consisting of the nucleic acid as represented in SEQ ID NO 2 or SEQ ID NO 18, or a functional part of said sequence, which is able to regulate gene expression in a root specific manner as SEQ ID NO 2 or SEQ ID NO 18 itself. More specifically, said isolated DNA sequence provides a novel type of a root-specific promoter, which differs from other known root-specific promoters in the fact that it is unexpectedly active in only distinct cells of the root, namely the meristem of the main and the lateral root, as well as in the vascular tissue of the main and lateral root and in the lateral root primordia.

The isolation of the root specific promoter RCH1 of the present invention is described in Example 1.

Another aspect of the invention is the isolation and sequencing of a root-specific Leucin Rich Repeat (LRR) receptor-like kinase gene from root tissue of *Arabidopsis thaliana* and is also described in Example 1. The nucleotide sequence is listed in the current specification as SEQ ID NO 3 and illustrated in Figure 8. This sequence is present as part of the sequence deposited under the GenBank Accession number AB017061, which represents the entire *Arabidopsis thaliana* chromosome 5.

The novelty of the RCH1 promoter sequence relative to known root-specific promoters is substantiated by its unique expression pattern in plant roots. The analysis of the RCH1 expression patterns was done using a RCH1prom-GUS hybrid operon that was transformed to *Arabidopsis thaliana* as outlined in Example 2. Analogous experiments with the RCH1prom-GUS hybrid operon were performed to establish the expression pattern of RCH1 in embryo's (Example 8 and Figure 15).

The RCH1 promoter is active in the meristem of the main and the lateral roots, as well as in the vascular tissue of the main and the lateral root and in the lateral root primordia. The promoter of the present invention is the first known promoter that regulates a CLAVATA HOMOLOG gene in roots, and is clearly distinct from other described CLAVATA-operons which are all active in the shoot and floral meristem. The present data thus clearly distinguish RCH1prom from other plant root-specific promoters, which

are not expressed in the meristem of the main and lateral root, in the vascular tissue of the main and lateral root and in the lateral root primordia.

From the results obtained by the RCH1::GFP expression studies in root it was even more clear that the RCH1 promoter is strongly active in the endodermis, cortex, epidermis, lateral root cap, in short in the "division zone". The RCH1 promoter is also active but in a lower manner, in the quiescent center and in the vascular tissue.

A further characteristic of the RCH1 promoter is its very high expression level. Based on the incubation time of the RCH1prom-GUS hybrid operon transformed plant with the GUS substrate (1 hour) and the very high output signal in the root of that plant (Example 2, Figure 7D), it is clear for persons skilled in the art that the RCH1 promoter is very strong.

A further characteristic of the present invention is that the activity of the RCH1 promoter in the meristem cells stops when these cells start to differentiate and to elongate to become cells of specified root tissue other than root vascular tissue. Surprisingly the promoter is activated again in cells of the pericycle that form lateral root primordia. This expression is extended in the meristem of the lateral root, but again, the cells that differentiate into lateral root tissue other than vascular tissue do not retain RCH1 promoter activity.

Functional parts of the promoter of the invention will have at least one of these characteristics.

According to a further embodiment, the present invention relates to an isolated nucleic acid comprising a transcriptional regulatory root promoter comprising at least part of a sequence as given in SEQ ID NO 1, 2 or 18.

According to another embodiment, the present invention relates to an isolated nucleic acid as represented in SEQ ID NO 2 or 18 wherein said transcriptional regulatory root promoter is operably linked to said nucleic acid sequence encoding a novel LRR receptor-like kinase or to an immunologically active and/or functional fragment thereof.

According to the invention, distinctive fragments of the newly identified promoter, which contribute specific features to said promoter, can be fused to another nucleic acid sequence (e.g. a promoter) in order to alter the sequential and/or regulatory features of the latter.

The invention thus relates to an isolated nucleic acid having transcriptional regulatory root specific, root-meristem specific or root-vascular-tissue specific promoter activity

comprising at least part of the DNA sequence as given in any of SEQ ID 1, 2 or 18 and a second transcriptional regulatory sequence.

In one embodiment, said second transcriptional regulatory sequence is for instance an inducible box or sequence, which is placed within the RCH1 promoter, wherein the function of the RCH1 promoter is not hampered but instead can be regulated, for instance by induction. In the latter embodiment, the RCH1 promoter is not used to confer root-specificity to other promoters, but a second sequence is used to confer an additional regulatory feature to the RCH1 promoter.

In another embodiment said second transcriptional regulatory sequence is a promoter sequence not normally exhibiting root-specificity, thereby resulting in a hybrid promoter with root-specificity.

Accordingly, another embodiment of the present invention is to confer root-specificity, root-meristem-specificity, and root-vascular-tissue-specificity and/or root abundant gene expression to other promoter sequences. This can be achieved by fusing elements of the here-disclosed RCH1 promoter to the promoter sequence of interest preferably the widely used ubiquitin promoter. Such modifications can be achieved by routine experimentation by those skilled in the art.

It should be further understood that the invention preferably relates to any of the isolated nucleic acids as described above which is DNA, cDNA, genomic DNA, synthetic DNA or RNA wherein T is replaced by U.

The invention also relates to a nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with any of the nucleic acids of the invention. The invention also relates to a nucleic acid molecule of at least 15 nucleotides in length specifically amplifying a nucleic acid of the invention.

Further, the invention relates to a vector comprising a nucleic acid of the invention. Said vector may be an expression vector wherein the (said) nucleic acid sequence encoding a novel LRR receptor-like kinase or an immunologically active and/or functional fragment thereof, is operably linked to one or more control sequences allowing the expression in prokaryotic and/or eukaryotic host cells.

The invention also relates to a vector comprising at least part of a nucleic acid according to the invention and wherein a transcriptional regulatory root promoter of the invention or functional part(s) thereof, is operably linked to one or more genes of interest.

The invention also relates to a host cell containing a nucleic acid molecule of the invention or a vector of the invention wherein said nucleic acid or vector has been introduced by transformation, transfection or infection. Preferred host cells that are part of the invention are bacterial, insect, fungal, plant or animal cells.

- 5 Therefore, in a preferred embodiment, the invention relates to an isolated nucleic acid as represented in SEQ ID NO 3 encoding a LRR receptor-like kinase characterized by an amino acid sequence as given in SEQ ID NO 4 or 19. More specifically, said isolated nucleic acid encodes a plant ROOT CLAVATA HOMOLOG 1 (RCH1) of a novel type which is unexpectedly expressed in roots. Said novel plant RCH1 shows significant  
10 homology with CLAVATA 1, a protein involved in the maintenance of floral and shoot meristem cells.

An initial comparative amino acid sequence homology search using the BLASTP 2.0.8 software (Altschul et al. 1997) revealed that most significant alignments of RCH1 protein are produced with receptor-like protein kinases, more specifically with leucine rich  
15 receptor-like kinases such as CLAVATA1 (CLV1). Furthermore, the most homologous protein with a known function of this list, is the CLAVATA 1 receptor-like kinase from *Arabidopsis thaliana*. Therefore the novel gene was denominated a ROOT CLAVATA HOMOLOG 1 gene (RCH1).

An amino acid sequence alignment of the RCH1 protein and CLAVATA 1 from *A.*  
20 *thaliana* is shown in Figure 10. Calculation of the percentage of identical amino acids residues between RCH1 and CLV1 teaches that RCH1 is 39% identical to CLV1. Additional to 39% of identical amino acids, there is 49% of the amino acids that belong to the same class of amino acids and therefore have the same physical and chemical properties.

- 25 The classification of RCH1 as a novel type of CLAVATA is documented further in figures 11 and 12.

The conclusions we can draw from figure 11 is that based on the sequence, RCH1 is closer related to CLV1 compared to other known CLV1 homologues, such as ERECTA, BRASSINOLIDE INSENSITIVE and HAESA.

- 30 The fact that the novel LRR receptor like protein described in the present invention is rather a root clavata 1 homologue than another kind of LRR receptor-like kinase (e.g. resistance genes) is illustrated in Figure 12 in which the evolutionary relationship between clavata homologues and resistance genes is shown. The resistance genes



(RPS5, RPS2, RPP8, RPM1, RPS4, N, RPP5) are described in "Meyers BC, Dickerman AW, Michelmore RW, Sivaramakrishnan S, Sobral BW, Young ND. Plant disease resistance genes encode members of an ancient and diverse protein family within the nucleotide-binding super family. Plant J. 1999 Nov;20(3):317-32".

- 5 The present invention relates to a ROOT *CLAVATA* HOMOLOG 1 gene that differs from other known *CLAVATA*-like genes, in the fact that it is active in the root meristem, whereas the other known *CLAVATA* genes and *CLAVATA* homologs (like *ERECTA*, *BRASSINOLIDE* *INSENSITIVE* and *HAESA*) are only active in the shoot and floral meristem.
- 10 According to another embodiment, the invention also relates to nucleic acids encoding a functional plant LRR receptor-like kinase comprising one or more protein regions, distinguishing said plant LRR receptor-like kinase from those LRR receptor-like kinases known in the art.

The inventors found that the predicted amino acid sequence (BAB10317.1, receptor protein kinase-like protein) corresponding to the database entry with accession number AB017061 (*A. thaliana* chromosome 5), was missing a part of the amino acid sequence (namely, 25 amino acids as represented in SEQ ID NO. 5 of the present patent application). This extra stretch of 25 amino acids was identified by the cloning and the sequencing of the corresponding cDNA of the present invention. Therefore, the

15 predicted prior art sequence is believed not to represent a functional enzyme.

20 The invention thus relates to an isolated nucleic acid selected from the group consisting of:

- (a) a nucleic acid consisting of at least part, preferably at least a functional part, of the DNA sequence as given in SEQ ID NO 1 or 3, or the complement thereof,
- 25 (b) a nucleic acid encoding a protein as given in SEQ ID NO 4 or 19 or encoding a fragment of said protein, wherein said fragment comprises the sequence as represented in SEQ ID NO 5, and
- (c) a nucleic acid encoding a protein with an amino acid sequence which is at least 60%, preferably at least 65 %, 70 %, 75%, 80%, 85%, more preferably at least 90% or 95% identical to the protein as given in SEQ ID NO 4 or 9, wherein said
- 30 amino acid sequence comprises the sequence as represented in SEQ ID NO 5,

characterised in that said nucleic acid encodes a novel LRR receptor-like kinase protein or an immunologically active and/or functional fragment of such a protein, and further provided that said nucleic acid is not one of the nucleic acids as deposited under the GenBank Accession numbers AB017061 or AQ966419.

5 The present invention also relates to an isolated LRR receptor-like kinase (protein) comprising one of the polypeptides selected from the group consisting of:

(a) a polypeptide as given in SEQ ID NO 4 or SEQ ID NO 19,

(b) a polypeptide comprising the amino acid sequence as given in SEQ ID NO 5, and,

10 (c) a polypeptide comprising the sequence represented in SEQ ID NO 5 and encoded by a nucleic acid as given in SEQ ID NO 1 or 3,

or a homologue or a derivative of said protein, or a fragment or an immunologically active and/or functional fragment thereof, provided that said homologue is not the amino acid sequence as described under the GenBank Accession number BAB10317.1.

15 With the expression "homologue" when referring to homologues of the LRR receptor-like kinase (protein), a polypeptide is meant which is at least 65% identical to the amino acid sequence of RCH1 as represented in SEQ ID NO 4 or 19.

More preferred, the invention relates to a protein consisting of an amino acid sequence as given in SEQ ID NO 4 or SEQ ID NO 19. More specifically, the present invention  
20 relates to a root CLAVATA homologue or a functional homologue thereof. A preferred fragment of said protein is represented in SEQ ID NO 5.

A further embodiment of the invention comprises homologues, derivatives, immunologically active and/or functional fragments of the LRR receptor-like kinase (protein) according to the invention, and proteins comprising said homologues,  
25 derivatives, immunologically active and/or functional fragments of said LRR receptor like kinase (protein).

As such, the present invention also relates to an isolated polypeptide comprising the sequence represented in SEQ ID NO 5 encodable by a nucleic acid molecule of the invention as defined above, or a homologue or a derivative thereof, or a fragment or an  
30 immunologically active and/or functional fragment thereof.

In a more preferred embodiment, the invention relates to a polypeptide, encodable by a nucleic acid molecule of the invention and which has an amino acid sequence as given in SEQ ID NO 4 or SEQ ID NO 19 or comprising an amino acid sequence as given in

SEQ ID NO 5, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.

Any of said proteins can be produced in a biological system, e.g. a cell culture. Alternatively any of said proteins is chemically manufactured e.g. by solid phase peptide  
5 synthesis. Said proteins or fragments thereof can be part of a fusion protein as is the case in e.g. a two-hybrid assay which enables e.g. the identification of proteins interacting with the LRR receptor-like kinase according to the present invention.

Therefore, according to another embodiment, the invention also relates to a method for  
10 producing a polypeptide or a protein of the invention comprising culturing a host cell further specified above under conditions allowing the expression of the polypeptide and recovering the produced polypeptide from the culture.

The proteins or fragments thereof obtained by a method of the invention are furthermore useful e.g. to modulate the interaction between a LRR receptor-like kinase according to the invention and a ligand of the receptor and/or other identified interacting protein  
15 partners. Chemically synthesized peptides are particularly useful e.g. as a source of antigens for the production of antisera and/or antibodies.

The current invention thus furthermore encompasses antisera and/or antibodies specifically recognizing the LRR receptor-like kinase according to the invention or immunologically active parts or epitopes thereof. Said antisera and/or antibodies are  
20 useful in many areas related to the invention including: (i) identification in any organism, preferably plants, of other LRR receptor-like kinase and their genes according to the invention; (ii) quantification of synthesis in organisms and/or recombinant organisms of the LRR receptor-like kinase according to the invention; (iii) purification of the LRR receptor-like kinase according to the invention; (iv) immunoprecipitation of the LRR  
25 receptor-like kinase according to the invention e.g. as a way to identify other protein partners complexing with said LRR receptor-like kinase; (v) immunolocalization of the LRR receptor-like kinase according to the invention which is expressed in an organism or a recombinant organism.

The invention also relates to a method for the production of transgenic plants, plant cells  
30 or plant tissues comprising the introduction of a nucleic acid molecule of the invention in an expressible format or a vector as described earlier in said plant, plant cell or plant tissue

The invention also relates to a method as described above further comprising regenerating a plant from said plant cell.

In a further embodiment, the invention also relates to a transgenic plant cell comprising a nucleic acid sequence of the invention which is operably linked to regulatory elements  
5 allowing transcription and/or expression of said nucleic acid in plant cells or obtainable by one of the methods described above. Preferably, in said transgenic plant cell one of the nucleic acids of the invention is stably integrated into the genome of said plant cell. The invention further relates to a transgenic plant or plant tissue comprising said plant cells and to a harvestable part of said plant or tissue. Said harvestable part of said plant  
10 or tissue is preferably selected from the group consisting of seeds, leaves, fruits, stem cultures, rhizomes, tubers and bulbs.

The invention further extends to the progeny derived from any of the plants or plant parts described above.

The present invention is applicable to any plant, in particular a monocotyledonous plants  
15 and dicotyledonous plants including a fodder or forage legume, ornamental plant, food crop, tree, or shrub selected from the list comprising *Acacia spp.*, *Acer spp.*, *Actinidia spp.*, *Aesculus spp.*, *Agathis australis*, *Albizia amara*, *Alsophilla tricolor*, *Andropogon spp.*, *Arachis spp.*, *Areca catechu*, *Astelia fragrans*, *Astragalus cicer*, *Baikiaea plurijuga*, *Betula spp.*, *Brassica spp.*, *Bruguiera gymnorrhiza*, *Burkea africana*, *Butea frondosa*, *Cadaba farinosa*, *Calliandra spp.*, *Camellia sinensis*, *Canna indica*, *Capsicum spp.*, *Cassia spp.*,  
20 *Centroema pubescens*, *Chaenomeles spp.*, *Cinnamomum cassia*, *Coffea arabica*, *Colophospermum mopane*, *Coronillia varia*, *Cotoneaster serotina*, *Crataegus spp.*, *Cucumis spp.*, *Cupressus spp.*, *Cyathea dealbata*, *Cydonia oblonga*, *Cryptomeria japonica*, *Cymbopogon spp.*, *Cynthea dealbata*, *Cydonia oblonga*, *Dalbergia monetaria*,  
25 *Davallia divaricata*, *Desmodium spp.*, *Dicksonia squarosa*, *Diheteropogon amplexans*, *Dioclea spp.*, *Dolichos spp.*, *Dorycnium rectum*, *Echinochloa pyramidalis*, *Ehretia spp.*, *Eleusine coracana*, *Eragrostis spp.*, *Erythrina spp.*, *Eucalyptus spp.*, *Euclea schimperi*, *Eulalia villosa*, *Fagopyrum spp.*, *Feijoa sellowiana*, *Fragaria spp.*, *Flemingia spp.*, *Freyinetia banksii*, *Geranium thunbergii*, *Ginkgo biloba*, *Glycine javanica*, *Glicidia spp.*,  
30 *Gossypium hirsutum*, *Grevillea spp.*, *Guibourtia coleosperma*, *Hedysarum spp.*, *Hemarthra altissima*, *Heteropogon contortus*, *Hordeum vulgare*, *Hyparrhenia rufa*, *Hypericum erectum*, *Hyperthelia dissoluta*, *Indigo incarnata*, *Iris spp.*, *Leptarrhena pyrolifolia*, *Lepidiza spp.*, *Lettuca spp.*, *Leucaena leucocephala*, *Loudetia simplex*,

*Lotonous bainesii*, *Lotus* spp., *Macrotyloma axillare*, *Malus* spp., *Manihot esculenta*,  
*Medicago sativa*, *Metasequoia glyptostroboides*, *Musa sapientum*, *Nicotianum* spp.,  
*Onobrychis* spp., *Ornithopus* spp., *Oryza* spp., *Peltophorum africanum*, *Pennisetum*  
5 *sp.*, *Persea gratissima*, *Petunia* spp., *Phaseolus* spp., *Phoenix canariensis*, *Phormium*  
*cookianum*, *Photinia* spp., *Picea glauca*, *Pinus* spp., *Pisum sativum*, *Podocarpus totara*,  
*Pogonarthria fleckii*, *Pogonarthria squarrosa*, *Populus* spp., *Prosopis cineraria*,  
*Pseudotsuga menziesii*, *Pterolobium stellatum*, *Pyrus communis*, *Quercus* spp.,  
*Rhaphiolepis umbellata*, *Rhopalostylis sapida*, *Rhus natalensis*, *Ribes grossularia*,  
*Ribes* spp., *Robinia pseudoacacia*, *Rosa* spp., *Rubus* spp., *Salix* spp., *Schyzachyrium*  
10 *sanguineum*, *Sciadopitys verticillata*, *Sequoia sempervirens*, *Sequoiadendron*  
*giganteum*, *Sorghum bicolor*, *Spinacia* spp., *Sporobolus fimbriatus*, *Stiburus*  
*alopecuroides*, *Stylosanthos humilis*, *Tadehagi* spp., *Taxodium distichum*, *Themeda*  
*triandra*, *Trifolium* spp., *Triticum* spp., *Tsuga heterophylla*, *Vaccinium* spp., *Vicia*  
*spp.*, *Vitis vinifera*, *Watsonia pyramidata*, *Zantedeschia aethiopica*, *Zea mays*, amaranth,  
15 artichoke, asparagus, broccoli, brussel sprout, cabbage, canola, carrot, cauliflower,  
celery, collard greens, flax, kale, lentil, oilseed rape, okra, onion, potato, rice, soybean,  
straw, sugarbeet, sugar cane, sunflower, tomato, squash, and tea, amongst others, or  
the seeds of any plant specifically named above or a tissue, cell or organ culture of any  
of the above species.

20 Yet according to another embodiment the present invention relates to a method for  
conferring root-specificity and/or root-meristem-specificity and/or root-vascular-tissue-  
specificity and/or root-endodermis-specificity and/or root-cortex-specificity and/or root-  
epidermis-specificity and/or lateral-root cap-specificity and/or abundant expression in  
roots to other promoter sequences comprising the fusion of at least part of the DNA  
25 sequence as given in SEQ ID NO 1, 2 or 18 to a (second) transcriptional regulatory  
promoter sequence normally not exhibiting root-specificity.

The invention further relates to method for root-specific expression of a gene(s) of  
interest comprising operably linking of said gene(s) of interest to a transcriptional  
regulatory root-specific promoter comprising at least part of the nucleic acids as given in  
30 SEQ ID NO 1, 2 or 18 as defined above, and possibly in combination/or a (second)  
transcriptional regulatory promoter sequence not normally exhibiting root-specificity, as  
defined earlier.

The invention further relates to a method for modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology comprising the modification of expression in the meristem of the main and the lateral roots, or in the vascular tissue of the root, or in the lateral root primordia of a gene(s) of interest operably linked to a transcriptional regulatory root promoter as defined earlier.

The ability to drive tissue-specific expression of any gene of interest in plants is considered to be of great agronomic importance. This art can be used to exclude expression in eatable parts of the plant, which averts the uptake of ectopic components. Alternatively the expression of the transgene can be restricted to the relevant tissues of the plant so the total organism does not spend too much energy in superfluous production of the foreign component. This technique can also reduce potential yield loss by limiting the expression of some pernicious, yet useful agronomic genes, to specific tissues only. Said genes of interest, which can be expressed in a root-specific manner under the control of the present invention can also be involved in phytoremediation, enhanced root-growth (like cell cycle genes or cytokinin oxidase) or disease tolerance. These methods are more elaborated in detail below.

The invention described here provides a new transcriptional regulatory element which can be operably linked to a gene effecting and/or modifying root metabolites which can be involved in phytoremediation. Examples are genes involved in the production of citric and maleic acids that can be secreted by the plant and subsequently form complexes with the aluminium in the soil. Root specific expression of the aluminium-resistance gene that give rise to said metabolites can therefore allow plants to grow on acid soils, like those of south-eastern United States, Central and south America, North Africa and parts of India and China, where the aluminium is set free as an ion that poisons plant roots.

Alternatively, the promoter of the present invention also provides a tool to be used in a strategy for environmental remediation. Accordingly, a metal-resistance gene such as the TaPCS1 or the CAD1 gene, contributing cadmium tolerance, can be expressed specifically in roots to enable enhanced growth of transgenic plants on soils contaminated with cadmium. The metals can consequently be extracted from the soil and stored in the plant material, which could then be harvested and incinerated.

Therefore according to preferred embodiments, the present invention relates to a method for phytoremediation or environmental remediation comprising the expression of

a gene(s) of interest under the control of a transcriptional regulatory root promoter as defined earlier.

In conclusion the present invention provides a method for broadening the range of soils or environments in which a plant may thrive or that can be used for agriculture.

- 5 Plant root development is an essential determinant of plant growth and crop yield, so that yield could be enhanced by induced changes in expression of root-specific regulatory factors. Cyclin D for example, which promotes the cell division, can be expressed under the root-specific control of the current invention and hence contribute to increased growth rate and crop yield. Also genes that are involved in biosynthesis and  
10 perception of phytohormones like auxins, cytokinines, ethylene, abscisic acid and gibberellins can be used to enhance root growth and development. However, the root-specific expression of these genes might be pivotal for the overall welfare of the transgenic plant.

- It is known that the constitutive and ubiquitous expression of the cytokinin oxidase in  
15 *Arabidopsis thaliana* leads to the promotion of an enlarged root system and enhanced root biomass, but also to diminished shoot growth. To overcome this problem the cytokinin oxidase can be operably linked to the promoter of the present invention as described in example 6, which will enhance root growth without negatively affecting shoot growth. This enhanced root growth in the plant can be used for bioremediation or  
20 phytoremediation.

- Also the said gene of interest, that can be operably linked to the promoter of the present invention, can be a root transcription factor, such as Alfin. Alfin binds to promoter fragments of predominantly root-specific and salt inducible genes and its activity has major implications for crop plant yield (Winicov 2000) (Winicov & Bastola 1999)). In  
25 conclusion the present invention provides a method for enhancing plant growth and crop yield.

- It is also clear for the skilled artisan that the promoter of the present invention provides a useful tool to be used for the introduction of resistance to soil-borne pathogen attack to plants, a widespread target for genetic improvement of crop plants. Certain disease-  
30 inducing microorganisms attack the belowground plant tissue and any genetic modification to contribute resistance to such organisms will require expression of the resistance-gene in the roots. To overcome this problem, the coding region that is operably linked to the promoter of the present invention preferably encodes a protein

which is toxic to root-attacking organisms and more preferably the protein is an insecticidal endotoxin of *Bacillus thuringiensis*. Similarly to the above-mentioned pathogens, there are pests attacking plants. Certain insect species attack green leaf tissue, whereas other, for example coleoptera, attack the roots. Corn root worm  
5 (*Diabrotica undecimpunctata howardi* Barber) for example is a particularly difficult pest to control or to eradicate. It attacks the plant below the soil line, where insecticides are difficult or impossible to apply effectively. Corn root worm can be eradicated by root-specific expression of the Cry III gene, which produces a component, specifically toxic for coleoptera. Also root-specific expression of the limonene cyclase gene in  
10 combination with GPPP synthetase would be larvicidal. Another pest, nematodes, that penetrate the root and form root-knots or cysts, which cause severe damage to crops throughout the world, could be treated with the root-specific expression of nematode resistance genes such as Hs1<sup>P<sub>root</sub>-1</sup>, Ni-1 and Gpa2.

In conclusion the invention disclosed here provides a method for conferring enhanced  
15 resistance to pathogens to a transgenic plant which pathogens attack the belowground plant tissue comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined earlier.

In yet another embodiment of the invention, the invention provides a method to enhanced freezing tolerance comprising the expression of a gene(s) of interest under  
20 the control of a transcriptional regulatory root-specific promoter as defined earlier. Preferred is the overexpressing of the *Arabidopsis thaliana* DNA binding factors CBF1 and DREB1A that induce cold-regulated (COR) genes (Jaglo-Ottosen et al. 1998). However, constitutive expression of these genes in all organs of the plant led to growth retardation under normal growth conditions. Root-specific expression of these freezing-  
25 tolerance genes under control of the promoter sequence of the present invention can make the plant tolerant to cold temperatures of the soil without affecting the growth of the plant. This technique can have important agronomic potential since lots of soils are inaccessible for crop production because of presence of permanently frozen layers in the soil.

Alternatively, other genes of interest can tolerize the transgenic plant for other weather  
30 conditions like drought (e.g. oat arginin decarboxylase gene) and high salt conditions (e.g. MliD gene). In conclusion, the present invention provides a method to contribute to a transgenic plant stress-tolerance.



Further advantages to root-preferred gene expression include the production of useful proteins in an industrial setting. Light-sensitive proteins may be synthesized in root tissue such that said proteins are not exposed to light. Therefore the present invention also relates to a method for the production of light-sensitive proteins comprising the expression of a gene encoding said light-sensitive protein under the control of a transcriptional regulatory root-specific promoter of the invention.

Also considered as an embodiment of the invention is the use of the RCH1 promoter for applications based on the inhibition of expression of native DNA sequences within the plant's root to achieve a desired phenotype resulting from the silencing of the native gene. In this case, such inhibition might be accomplished with transformation of the plant to comprise a transcriptional regulatory root-specific promoter of the present invention operably linked to e.g. an antisense nucleotide sequence to the gene of interest, a gene silencing construct or a ribozyme.

Those skilled in the art will be aware that plants secure the formation of organs throughout their life span by developing and maintaining a collection of stem cells termed the meristem. The promoter of the present invention offers the opportunity to deliver a certain protein specifically to these cells. For example overexpression of genes that influence the proliferation and/or the differentiation of the root meristem could contribute to a higher turnover of the meristem cells and therefor enhance the root architecture.

The promoter of the present invention drives the *Arabidopsis thaliana* gene ROOT CLAVATA HOMOLOG 1 (RCH1). Since the *clavata*-genes are key role players in shoot-meristem maintenance, persons skilled in the art may recognize the possible involvement of the RCH1 operon in root meristem maintenance. Therefor one could assume that the control of this promoter is influenced by signals that promote meristem maintenance. This might be an advantage when aiming to stimulate the root meristem-formation, which eventually could result in more roots, higher uptake of resources and in higher crop yield.

Therefore, according to preferred embodiments, the invention relates to methods for stimulating meristem formation and/or maintaining root meristem comprising the expression of a gene that influences the proliferation and/or differentiation of the root meristeme under the control of a transcriptional regulatory root (-specific) promoter of the invention.

The present invention therefore relates to a method for stimulating root meristem formation or for root-meristeme maintenance comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase, for instance the LRR receptor-like kinase of the invention, wherein optionally said nucleic acid is operably linked to a plant-operable promoter sequence, for instance said plant-operable promoter being the transcriptional regulatory root promoter as defined earlier.

As mentioned supra, the combined gene expression data and homology to CLAVATA 1 indicate that the novel plant LRR receptor-like kinase of the invention, RCH1 (SEQ ID No. 3, 4, 19), is very likely to exert a function that is involved in the development and/or maintenance of the root meristem. As the root is the main plant organ for nutrient uptake from the environment, enhanced root formation could eventually lead to overall growth enhancement and better yield. Therefore preferred genes to be expressed in the methods for stimulating meristem formation and/or for root meristeme maintenance are genes encoding the polypeptide(s) of the invention.

The present invention also relates to a method for enhancing root formation and/or root growth comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase, for instance the LRR receptor-like kinase of the invention, wherein optionally said nucleic acid is operably linked to a plant-operable promoter sequence, for instance said plant-operable promoter being the transcriptional regulatory root promoter as defined earlier.

The present invention therefore relates to a method for enhancing overall growth and yield comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase, for instance the LRR receptor-like kinase of the invention, wherein optionally said nucleic acid is operably linked to a plant-operable promoter sequence, for instance said plant-operable promoter being the transcriptional regulatory root promoter as defined earlier.

The present invention also relates to a method for enhancing overall growth and yield comprising the expression of a protein of the invention under the control of a transcriptional regulatory root (or root-specific) promoter of the invention.

Also in yet other embodiments of the invention, methods are provided for modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or

physiology comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase, for instance the LRR receptor-like kinase of the invention and defined above, wherein optionally said nucleic acid is operably linked to a plant-operable promoter sequence, for instance said plant-operable promoter being the transcriptional regulatory root promoter as defined earlier.

The present invention further relates to a method to confer pathogen resistance to a transgenic plant comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase, for instance the LRR receptor-like kinase of the invention and defined above, wherein optionally said nucleic acid is operably linked to a plant-operable promoter sequence, for instance said plant-operable promoter being the transcriptional regulatory root promoter as defined earlier.

Many pathogen resistance genes are LRR receptor-like proteins, which initiate a signal transduction pathway in the root cells, leading to the activation of defense mechanisms against the pathogen. The novel LRR receptor-like kinase might contribute to pathogen resistance to the transgenic plant when transformed herein.

Yet another embodiment of the invention relates to methods to perform a functional analysis of the DNA sequence according to the present invention, encoding a novel LRR receptor-like protein. For persons skilled in the art it is possible to make single or double knock-out mutants of *Arabidopsis thaliana*, wherein the DNA of the present invention is inactivated. Also introduction of the antisense sequence of the DNA of the invention, as well as the overexpression of its sense sequence in a transgenic plant can elucidate a specific feature in this transgenic plant. Furthermore, the loss of function of this gene of the invention by introduction of dominant negative mutations can result in interesting characteristics of the mutant plant.

Another aspect of the present invention is the revelation of a new target for herbicides or growth regulators. When said gene of the invention is an essential gene for plant cell viability, it can be a target for a herbicide or growth regulator.

Therefore the invention relates to a method for identifying and obtaining proteins interacting with a (LRR receptor-like kinase) polypeptide of the invention comprising a screening assay wherein said polypeptide is used or expressed.

In a preferred embodiment the invention relates to a method for identifying and obtaining proteins interacting with an LRR receptor-like kinase (protein) comprising a two-hybrid

screening system wherein a nucleic acid encoding a polypeptide of the invention as a bait and a cDNA library as prey are expressed.

The invention also relates to a method for modulating the interaction between an LRR receptor-like kinase (protein) of the invention and interacting proteins obtainable by a method as described above.

The invention further relates to a method for identifying and obtaining compounds interacting with an LRR receptor-like kinase (protein) comprising the steps of:

- a) providing a two-hybrid screening system wherein a polypeptide or protein of the invention and a protein interacting with said (LRR receptor-like kinase) polypeptide or protein or an interacting protein obtainable by a method of claim as claimed above are expressed,
- b) interacting said compound with the complex formed by the expressed proteins as defined in a),
- c) detecting a second complex, wherein the presence of said second complex identifies a compound which specifically binds to one of said polypeptides or said second complex, and
- d) identifying the compound.

The invention also relates to a method for identifying compounds or mixtures of compounds which specifically bind to a polypeptide of the invention, comprising:

- a) combining a polypeptide of the invention with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,
- b) detecting complex formation, wherein the presence of a complex identifies a molecule which specifically binds said polypeptide.

The invention further relates to the use of a molecule identified by means of one of the methods described above as a plant growth regulator or herbicide.

The invention also relates to a method for the production of a plant growth regulator or herbicide composition comprising the steps of one of the methods described above and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture.

The invention also relates to the use of any of the nucleic acid molecules, vectors, polypeptides or antibodies of described herein for modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology.

The invention also relates to a diagnostic composition comprising at least one of the nucleic acid molecules, vectors, polypeptides or antibodies of the invention.

It is further important to notice that CLAVATA 1 has many analogues in other plant species, so it is likely that also the promoter of RCH1, the RCH1 ORF or the complete RCH1 operon can be functional in those other plant species, and possibly exert the same specific activation pattern. This can be investigated by root-specific reporter gene expression in rice mediated by the *Arabidopsis thaliana* RCH1 promoter as described in example 5.

#### Definitions and elaborations to the embodiments

The terms "protein(s)", "peptide(s)" or "oligopeptide(s)", when used herein refer to amino acids in a polymeric form of any length. Said terms also include known amino acid modifications such as disulphide bond formation, cysteinylolation, oxidation, glutathionylation, methylation, acetylation, farnesylation, biotinylation, stearylolation, formylation, lipoic acid addition, phosphorylation, sulphation, ubiquitination, myristoylation, palmitoylation, geranylgeranylation, cyclization (e.g. pyroglutamic acid formation), oxidation, deamidation, dehydration, glycosylation (e.g. pentoses, hexosamines, N-acetylhexosamines, deoxyhexoses, hexoses, sialic acid etc.), acylation and radiolabels (e.g.  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^3\text{H}$ ) as well as non-naturally occurring amino acid residues, L-amino acid residues and D-amino acid residues.

"Homologues" or "Homologs" of a protein of the invention are those peptides, oligopeptides, polypeptides, proteins and enzymes which contain amino acid substitutions, deletions and/or additions relative to the said protein with respect to which they are a homolog, without altering one or more of its functional properties, in particular without reducing the activity of the resulting. For example, a homolog of said protein will consist of a bioactive amino acid sequence variant of said protein. To produce such homologs, amino acids present in the said protein can be replaced by other amino acids having similar properties, for example hydrophobicity, hydrophilicity, hydrophobic moment, antigenicity, propensity to form or break  $\alpha$ -helical structures or  $\beta$ -sheet structures, and so on. An overview of physical and chemical properties of amino acids is given in Table 1.

Substitutional variants of a protein of the invention are those in which at least one residue in said protein amino acid sequence has been removed and a different residue

inserted in its place. Amino acid substitutions are typically of single residues, but may be clustered depending upon functional constraints placed upon the polypeptide; insertions will usually be of the order of about 1-10 amino acid residues, and deletions will range from about 1-20 residues. Preferably, amino acid substitutions will comprise conservative amino acid substitutions, such as those described *supra*.

**Table 1: Properties of naturally occurring amino acids.**

Charge properties / hydrophobicity	Side group	Amino Acid
nonpolar hydrophobic	Aliphatic Aliphatic, S-containing Aromatic Imino	Ala, Ile, Leu, Val Met Phe, Trp Pro
polar uncharged	Aliphatic Amide Aromatic Hydroxyl Sulphydryl	Gly Asn, Gln Tyr Ser, Thr Cys
positively charged	Basic	Arg, His, Lys
negatively charged	Acidic	Asp, Gly

Insertional amino acid sequence variants of a protein of the invention are those in which one or more amino acid residues are introduced into a predetermined site in said protein. Insertions can comprise amino-terminal and/or carboxy-terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Generally, insertions within the amino acid sequence will be smaller than amino or carboxyl terminal fusions, of the order of about 1 to 10 residues. Examples of amino- or carboxy-terminal fusion proteins or peptides include the binding domain or activation domain of a transcriptional activator as used in the yeast two-hybrid system, phage coat proteins, (histidine)<sub>6</sub>-tag, glutathione S-transferase, protein A, maltose-binding protein, dihydrofolate reductase, Tag•100

epitope (EETARFQPGYRS), c-myc epitope (EQKLISEEDL), FLAG®-epitope (DYKDDDK), lacZ, CMP (calmodulin-binding peptide), HA epitope (YPYDVPDYA), protein C epitope (EDQVDPRLIDGK) and VSV epitope (YTDIEMNRLGK).

Deletional variants of a protein of the invention are characterised by the removal of one or more amino acids from the amino acid sequence of said protein.

Homologues of the peptides or polypeptides of the invention contain a number of amino acid substitutions, deletions and/or additions relative to said peptide or polypeptide such that said homologues are still substantially different from the peptides or polypeptides known in the art. The difference between proteins can be calculated in terms of % identity according to methods well known in the art. Preferably, the homologues of the peptides or polypeptides of the present invention comprise the amino acid sequence represented in SEQ ID NO 5, optionally comprising minor amino acid substitutions such as those described in table 1. Other homologues of the polypeptides of the invention have an amino acid sequence which is at least 65 % identical to the sequence represented in SEQ ID NO 4 or 19.

Amino acid variants of a protein of the invention may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis and the like, or by recombinant DNA manipulations. The manipulation of DNA sequences to produce variant proteins which manifest as substitutional, insertional or deletional variants are well known in the art. For example, techniques for making substitution mutations at predetermined sites in DNA having known sequence are well known to those skilled in the art, such as by M13 mutagenesis, T7-Gen in vitro mutagenesis kit (USB, Cleveland, OH), QuickChange Site Directed mutagenesis kit (Stratagene, San Diego, CA), PCR-mediated site-directed mutagenesis or other site-directed mutagenesis protocols.

"Derivatives" of a protein of the invention are those peptides, oligopeptides, polypeptides, proteins and enzymes which comprise at least about five contiguous amino acid residues of said polypeptide but which retain the biological activity of said protein and which may further comprise additional naturally-occurring, altered glycosylated, acylated or non-naturally occurring amino acid residues compared to the amino acid sequence of a naturally-occurring form of said polypeptide. Alternatively or in addition, a derivative may comprise one or more non-amino acid substituents compared to the amino acid sequence of a naturally-occurring form of said polypeptide, for

example a reporter molecule or other ligand, covalently or non-covalently bound to the amino acid sequence such as, for example, a reporter molecule which is bound thereto to facilitate its detection. Preferably, the derivatives of the proteins of the present invention comprise the amino acid sequence as represented in SEQ ID NO 5.

- 5 With "immunologically active" is meant that a molecule or specific fragments thereof such as epitopes or haptens are recognized by, i.e. bind to antibodies. Preferably, the immunologically active or fragments of the proteins of the present invention comprise the amino acid sequence as represented in SEQ ID NO 5.

- 10 With "functional fragment" or "functional homologue" is meant a protein which comprises at least about 5, 10, 20, 50, 75, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1050, 1100, 1110, 1120, 1130, 1131, 1132, 1133 or 1134 contiguous amino acid residues of said polypeptide but which retains the biological activity of said protein. Preferably, the functional fragments or functional homologues of the polypeptide or proteins of the invention comprise the amino acid sequence as represented in SEQ ID NO 5 but  
15 wherein minor amino acid substitutions are allowed (see table 1).

In the context of the current invention are embodied homologs, derivatives and/or immunologically active fragments of any of the inventive LRR receptor-like kinase protein or polypeptide or homolog, derivative or fragment thereof as defined supra.

- "Antibodies" include monoclonal, polyclonal, synthetic or heavy chain camel antibodies  
20 as well as fragments of antibodies such as Fab, Fv or scFv fragments. Monoclonal antibodies can be prepared by the techniques as described in e.g. Liddle and Cryer (1991) which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized animals. Furthermore, antibodies or fragments thereof to a molecule or fragments thereof can be obtained by using methods as described in e.g. Harlow and  
25 Lane (1988). In the case of antibodies directed against small peptides such as fragments of a protein of the invention, said peptides are generally coupled to a carrier protein before immunization of animals. Such protein carriers include keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), ovalbumin and Tetanus toxoid. The carrier protein enhances the immune response of the animal and provides epitopes for  
30 T-cell receptor binding sites. The term "antibodies" furthermore includes derivatives thereof such as labelled antibodies. Antibody labels include alkaline phosphatase, PKH2, PKH26, PKH67, fluorescein (FITC), Hoechst 33258, R-phycoerythrin (PE), rhodamine (TRITC), Quantum Red, Texas Red, Cy3, biotin, agarose, peroxidase, gold



spheres and radiolabels (e.g.  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^3\text{H}$ ). Tools in molecular biology relying on antibodies against a protein include protein gel blot analysis, screening of expression libraries allowing gene identification, protein quantitative methods including ELISA and RIA, immunoaffinity purification of proteins, immunoprecipitation of proteins (e.g. Magyar et al. 1997) and immunolocalization. Other uses of antibodies and especially of peptide antibodies include the study of proteolytic processing (Löffler et al. 1994, Woulfe et al. 1994), determination of protein active sites (Lerner 1982), the study of precursor and post-translational processing (Baron and Baltimore 1982, Lerner et al. 1981, Sernler et al. 1982), identification of protein domains involved in protein-protein interactions (Murakami et al. 1992) and the study of exon usage in gene expression (Tamura et al. 1991).

Embodied in the current invention are antibodies recognizing a LRR receptor-like kinase polypeptide or protein, or homolog, derivative or fragment thereof as defined supra or specific epitopes of said polypeptide or protein.

The terms "gene(s)", "polynucleotide(s)", "nucleic acid sequence(s)", "nucleotide sequence(s)", "DNA sequence(s)" or "nucleic acid molecule(s)", when used herein refer to nucleotides, either ribonucleotides or deoxyribonucleotides or a combination of both, in a polymeric form of any length. Said terms furthermore include double-stranded and single-stranded DNA and RNA. Said terms also include known nucleotide modifications such as methylation, cyclization and 'caps' and substitution of one or more of the naturally occurring nucleotides with an analog such as inosine. Modifications of nucleotides include the addition of acridine, amine, biotin, cascade blue, cholesterol, Cy3<sup>®</sup>, Cy5<sup>®</sup>, Cy5.5<sup>®</sup> Dabcyl, digoxigenin, dinitrophenyl, Edans, 6-FAM, fluorescein, 3'-glyceryl, HEX, IRD-700, IRD-800, JOE, phosphate psoralen, rhodamine, ROX, thiol (SH), spacers, TAMRA, TET, AMCA-S<sup>®</sup>, SE, BODIPY<sup>®</sup>, Marina Blue<sup>®</sup>, Pacific Blue<sup>®</sup>, Oregon Green<sup>®</sup>, Rhodamine Green<sup>®</sup>, Rhodamine Red<sup>®</sup>, Rhodol Green<sup>®</sup> and Texas Red<sup>®</sup>. Polynucleotide backbone modifications include methylphosphonate, 2'-OMe-methylphosphonate RNA, phosphorothiorate, RNA, 2'-OMeRNA. Base modifications include 2-amino-dA, 2-aminopurine, 3'-(ddA), 3'dA(cordycepin), 7-deaza-dA, 8-Br-dA, 8-oxo-dA, N<sup>6</sup>-Me-dA, abasic site (dSpacer), biotin dT, 2'-OMe-5Me-C, 2'-OMe-propynyl-C, 3'-(5-Me-dC), 3'-(ddC), 5-Br-dC, 5-I-dC, 5-Me-dC, 5-F-dC, carboxy-dT, convertible dA, convertible dC, convertible dG, convertible dT, convertible dU, 7-deaza-dG, 8-Br-dG, 8-oxo-dG, O<sup>6</sup>-Me-dG, S6-DNP-dG, 4-methyl-indole, 5-nitroindole, 2'-OMe-inosine, 2'-dl, O<sup>6</sup>-

phenyl-dI, 4-methyl-indole, 2'-deoxynebularine, 5-nitroindole, 2-aminopurine, dP(purine analogue), dK(pyrimidine analogue), 3-nitropyrrole, 2-thio-dT, 4-thio-dT, biotin-dT, carboxy-dT, O<sup>4</sup>-Me-dT, O<sup>4</sup>-triazol dT, 2'-OMe-propynyl-U, 5-Br-dU, 2'-dU, 5-F-dU, 5-I-dU, O<sup>4</sup>-triazol dU and radiolabels (e.g. <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S, <sup>14</sup>C, <sup>32</sup>P, <sup>33</sup>P, <sup>3</sup>H). Said terms also encompass peptide nucleic acids (PNAs), a DNA analogue in which the backbone is a pseudopeptide consisting of N-(2-aminoethyl)-glycine units rather than a sugar. PNAs mimic the behaviour of DNA and bind complementary nucleic acid strands. The neutral backbone of PNA results in stronger binding and greater specificity than normally achieved. In addition, the unique chemical, physical and biological properties of PNA have been exploited to produce powerful biomolecular tools, antisense and antigene agents, molecular probes and biosensors. With "recombinant DNA molecule" or "chimeric gene" is meant a hybrid DNA produced by joining pieces of DNA from different sources. With "heterologous nucleotide sequence" is intended a sequence that is not naturally occurring with the promoter sequence. While this nucleotide sequence is heterologous to the promoter sequence, it may be homologous, or native, or heterologous, or foreign, to the plant host. "Sense strand" refers to the strand of a double-stranded DNA molecule that is homologous to a mRNA transcript thereof. The "anti-sense strand" contains an inverted sequence which is complementary to that of the "sense strand".

A "coding sequence" or "open reading frame" or "ORF" is defined as a nucleotide sequence that can be transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences, i.e. when said coding sequence or ORF is present in an expressible format. Said coding sequence or ORF is bounded by a 5' translation start codon and a 3' translation stop codon. A coding sequence or ORF can include, but is not limited to RNA, mRNA, cDNA, recombinant nucleotide sequences, synthetically manufactured nucleotide sequences or genomic DNA. Said coding sequence or ORF can be interrupted by intervening nucleic acid sequences.

Table 2. Degeneracy of the genetic code.

Amino Acid	Three-letter code	One-letter code	Possible codons					
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Asparagine	Asn	N	AAC	AAU				
Aspartic Acid	Asp	D	GAC	GAU				
Cysteine	Cys	C	UGC	UGU				
Glutamic Acid	Glu	E	GAA	GAG				
Glutamine	Gln	Q	CAA	CAG				
Glycine	Gly	G	GGA	GGC	GGG	GGU		
Histidine	His	H	CAC	CAU				
Isoleucine	Ile	I	AUA	AUC	AUU			
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU
Lysine	Lys	K	AAA	AAG				
Methionine	Met	M	AUG					
Phenylalanine	Phe	F	UUC	UUU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				
Valine	Val	V	GUA	GUC	GUG	GUU		
Possible "STOP" codons								
UAA	UAG	UGA						

- Genes and coding sequences essentially encoding the same protein but isolated from different sources can consist of substantially divergent nucleic acid sequences. Reciprocally, substantially divergent nucleic acid sequences can be designed to effect expression of essentially the same protein. Said nucleic acid sequences are the result of e.g. the existence of different alleles of a given gene, of the degeneracy of the genetic code or of differences in codon usage. Thus, as indicated in Table 2, amino acids such

as methionine and tryptophan are encoded by a single codon whereas other amino acids such as arginine, leucine and serine can each be translated from up to six different codons. Differences in preferred codon usage are illustrated below for *Agrobacterium tumefaciens* (a bacterium), *A. thaliana*, *M. sativa* (two dicotyledonous plants) and *Oryza sativa* (a monocotyledonous plant). These examples were extracted from (<http://www.kazusa.or.jp/codon>). To give one example, the codon GGC (for glycine) is the most frequently used codon in *A. tumefaciens* (36.2 %), is the second most frequently used codon in *O. sativa* but is used at much lower frequencies in *A. thaliana* and *M. sativa* (9 % and 8.4 %, respectively). Of the four possible codons encoding glycine (see Table 6), said GGC codon is most preferably used in *A. tumefaciens* and *O. sativa*. However, in *A. thaliana* this is the GGA (and GGU) codon whereas in *M. sativa* this is the GGU (and GGA) codon.

"Hybridization" is the process wherein substantially homologous complementary nucleotide sequences anneal to each other. The hybridization process can occur entirely in solution, i.e. both complementary nucleic acids are in solution. Tools in molecular biology relying on such a process include the polymerase chain reaction (PCR; and all methods based thereon), subtractive hybridization, random primer extension, nuclease S1 mapping, primer extension, reverse transcription, cDNA synthesis, differential display of RNAs, and DNA sequence determination. The hybridization process can also occur with one of the complementary nucleic acids immobilized to a matrix such as magnetic beads, Sepharose beads or any other resin. Tools in molecular biology relying on such a process include the isolation of poly (A+) mRNA. The hybridization process can furthermore occur with one of the complementary nucleic acids immobilized to a solid support such as a nitrocellulose or nylon membrane or immobilized by e.g. photolithography to e.g. a silicious glass support (the latter known as nucleic acid arrays or microarrays or as nucleic acid chips). Tools in molecular biology relying on such a process include RNA and DNA gel blot analysis, colony hybridization, plaque hybridization, in situ hybridization and microarray hybridization. In order to allow hybridization to occur, the nucleic acid molecules are generally thermally or chemically denatured to melt a double strand into two single strands and/or to remove hairpins or other secondary structures from single stranded nucleic acids. The stringency of hybridization is influenced by conditions such as temperature, salt concentration and hybridization buffer composition. High stringency conditions for hybridization include high

temperature and/or low salt concentration (salts include NaCl and Na<sub>3</sub>-citrate) and/or the inclusion of formamide in the hybridization buffer and/or lowering the concentration of compounds such as SDS (detergent) in the hybridization buffer and/or exclusion of compounds such as dextran sulfate or polyethylene glycol (promoting molecular crowding) from the hybridization buffer. Conventional hybridization conditions are described in e.g. Sambrook et al. (1989) but the skilled craftsman will appreciate that numerous different hybridization conditions can be designed in function of the known or the expected homology and/or length of the nucleic acid sequence. Sufficiently low stringency hybridization conditions are particularly preferred to isolate nucleic acids heterologous to the DNA sequences of the invention defined supra. Elements contributing to said heterology include allelism, degeneration of the genetic code and differences in preferred codon usage as discussed supra.

Clearly, the current invention embodies the use of the inventive DNA sequences encoding a LRR receptor-like kinase, homolog, derivative and/or immunologically fragment thereof as defined higher in any method of hybridization. The current invention furthermore also relates to DNA sequences specifically hybridizing to said inventive DNA sequences.

DNA sequences as defined in the current invention can be interrupted by intervening sequences. With "intervening sequences" is meant any nucleic acid sequence which disrupts a coding sequence comprising said inventive DNA sequence or which disrupts the expressible format of a DNA sequence comprising said inventive DNA sequence. Removal of the intervening sequence restores said coding sequence or said expressible format. Examples of intervening sequences include introns, mobilizable DNA sequences such as transposons and DNA tags such as e.g. a T-DNA. With "mobilizable DNA sequence" is meant any DNA sequence that can be mobilized as the result of a recombination event.

To effect expression of a protein in a cell, tissue or organ, preferably of plant origin, either the protein may be introduced directly to said cell, such as by microinjection or ballistic means or alternatively, an isolated nucleic acid molecule encoding said protein may be introduced into said cell, tissue or organ in an expressible format.

Preferably, the DNA sequence of the invention comprises a coding sequence or open reading frame (ORF) encoding a LRR receptor-like kinase or a homolog or derivative thereof or an immunologically active fragment thereof as defined supra. The preferred

protein of the invention comprises the amino acid sequence of said LRR receptor-like kinase.

With "vector" or "vector sequence" is meant a DNA sequence which can be introduced in an organism by transformation and can be stably maintained in said organism. Vector maintenance is possible in e.g. cultures of *Escherichia coli*, *A. tumefaciens*, *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*. Other vectors such as phagemids and cosmid vectors can be maintained and multiplied in bacteria and/or viruses. Vector sequences generally comprise a set of unique sites recognized by restriction enzymes, the multiple cloning site (MCS), wherein one or more non-vector sequence(s) can be inserted.

With "non-vector sequence" is accordingly meant a DNA sequence which is integrated in one or more of the sites of the MCS comprised within a vector.

"Expression vectors" form a subset of vectors which, by virtue of comprising the appropriate regulatory sequences enabling the creation of an expressible format for the inserted non-vector sequence(s), thus allowing expression of the protein encoded by said non-vector sequence(s). Expression vectors are known in the art enabling protein expression in organisms including bacteria (e.g. *E. coli*), fungi (e.g. *S. cerevisiae*, *S. pombe*, *Pichia pastoris*), insect cells (e.g. baculoviral expression vectors), animal cells (e.g. COS or CHO cells) and plant cells (e.g. potato virus X-based expression vectors, see e.g. Vance et al. 1998 - WO9844097).

The current invention clearly includes any vector or expression vector comprising a non-vector DNA sequence comprising the promoter sequence according to the present invention or a non-vector sequence encoding a LRR receptor-like kinase, homolog, derivative and/or immunologically active fragment thereof as defined supra.

As an alternative to expression vector-mediated protein production in biological systems, chemical protein synthesis can be applied. Synthetic peptides can be manufactured in solution phase or in solid phase. Solid phase peptide synthesis (Merrifield 1963) is, however, the most common way and involves the sequential addition of amino acids to create a linear peptide chain. Solid phase peptide synthesis includes cycles consisting of three steps: (i) immobilization of the carboxy-terminal amino acid of the growing peptide chain to a solid support or resin; (ii) chain assembly, a process consisting of activation, coupling and deprotection of the amino acid to be added to the growing peptide chain; and (iii) cleavage involving removal of the completed peptide chain from the resin and

removal of the protecting groups from the amino acid side chains. Common approaches in solid phase peptide synthesis include Fmoc/tBu (9-fluorenylmethyloxycarbonyl/t-butyl) and Boc (t-butyloxycarbonyl) as the amino-terminal protecting groups of amino acids. Amino acid side chain protecting groups include methyl (Me), formyl (CHO), ethyl (Et), acetyl (Ac), t-butyl (t-Bu), anisyl, benzyl (Bzl), trifluoroacetyl (Tfa), N-hydroxysuccinimide (ONSu, OSu), benzoyl (Bz), 4-methylbenzyl (Meb), thioanizyl, thiocresyl, benzyloxymethyl (Bom), 4-nitrophenyl (ONp), benzyloxycarbonyl (Z), 2-nitrobenzoyl (NBz), 2-nitrophenylsulphenyl (Nps), 4-toluenesulphonyl (Tosyl, Tos), pentafluorophenyl (Pfp), diphenylmethyl (Dpm), 2-chlorobenzyloxycarbonyl (Cl-Z), 2,4,5-trichlorophenyl, 2-bromobenzyloxycarbonyl (Br-Z), triphenylmethyl (Trityl, Trt), and 2,5,7,8-pentamethylchroman-6-sulphonyl (Pmc). During chain assembly, Fmoc or Boc are removed resulting in an activated amino-terminus of the amino acid residue bound to the growing chain. The carboxy-terminus of the incoming amino acid is activated by conversion into a highly reactive ester, e.g. by HBTU. With current technologies (e.g. PerSeptive Biosystems 9050 synthesizer, Applied Biosystems Model 431A Peptide Synthesizer), linear peptides of up to 50 residues can be manufactured. A number of guidelines is available to produce peptides that are suitable for use in biological systems including (i) limiting the use of difficult amino acids such as cys, met, trp (easily oxidized and/or degraded during peptide synthesis) or arg; (ii) minimize hydrophobic amino acids (can impair peptide solubility); and (iii) prevent an amino-terminal glutamic acid (can cyclize to pyroglutamate).

By "expressible format" is meant that the isolated nucleic acid molecule is in a form suitable for being transcribed into mRNA and/or translated to produce a protein, either constitutively or following induction by an intracellular or extracellular signal, such as an environmental stimulus or stress (mitogens, anoxia, hypoxia, temperature, salt, light, dehydration, etc) or a chemical compound such as IPTG (isopropyl- $\beta$ -D-thiogalactopyranoside) or such as an antibiotic (tetracycline, ampicillin, rifampicin, kanamycin), hormone (e.g. gibberellin, auxin, cytokinin, glucocorticoid, brassinosteroid, ethylene, abscisic acid etc), hormone analogue (iodoacetic acid (IAA), 2,4-D, etc), metal (zinc, copper, iron, etc), or dexamethasone, amongst others. As will be known to those skilled in the art, expression of a functional protein may also require one or more post-translational modifications, such as glycosylation, phosphorylation, dephosphorylation,

or one or more protein-protein interactions, amongst others. All such processes are included within the scope of the term "expressible format".

Preferably, expression of a protein in a specific cell, tissue, or organ, preferably of plant origin, is effected by introducing and expressing an isolated nucleic acid molecule encoding said protein, such as a cDNA molecule, genomic gene, synthetic oligonucleotide molecule, mRNA molecule or open reading frame, to said cell, tissue or organ, wherein said nucleic acid molecule is placed operably in connection with suitable regulatory sequences including a promoter, preferably a plant-expressible promoter, and a terminator sequence.

"Regulatory sequence" refers to control DNA sequences which are necessary to affect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoters, ribosomal binding sites, and terminators. In eukaryotes generally control sequences include promoters, terminators and enhancers or silencers.

The term "control sequence" is intended to include, at a minimum, all components the presence of which are necessary for expression, and may also include additional advantageous components and which determines when, how much and where a specific gene is expressed.

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences derived from a classical eukaryotic genomic gene, including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence and additional regulatory elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner.

The term "promoter" also includes the transcriptional regulatory sequences of a classical prokaryotic gene, in which case it may include a -35 box sequence and/or a -10 box transcriptional regulatory sequences.

The term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers, activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

The term "functional part" when referred to nucleic acids in relation to the above defined transcriptional regulatory sequences, refers to a part or parts of the nucleic acid having



the activity to specifically drive or promote transcription from sequences which are located downstream of said nucleic acid sequence.

Promoters may contain additional copies of one or more specific regulatory elements, to further enhance expression and/or to alter the spatial expression and/or temporal expression of a nucleic acid molecule to which it is operably connected. Such regulatory elements may be placed adjacent to a heterologous promoter sequence to drive expression of a nucleic acid molecule in response to e.g. copper, glucocorticoids, dexamethasone, tetracycline, gibberellin, cAMP, abscisic acid, auxin, wounding, ethylene, jasmonate or salicylic acid or to confer expression of a nucleic acid molecule to specific cells, tissues or organs such as meristems, leaves, roots, embryo, flowers, seeds or fruits.

In the context of the present invention, the promoter preferably is a plant-expressible promoter sequence. Promoters, however, that also function or solely function in non-plant cells such as bacteria, yeast cells, insect cells and animal cells are not excluded from the invention. By "plant-expressible" is meant that the promoter sequence, including any additional regulatory elements added thereto or contained therein, is at least capable of inducing, conferring, activating or enhancing expression in a plant cell, tissue or organ, preferably a monocotyledonous or dicotyledonous plant cell, tissue, or organ.

The terms "plant-operable" and "operable in a plant" when used herein, in respect of a promoter sequence, shall be taken to be equivalent to a plant-expressible promoter sequence.

In the present context, a "regulatable promoter sequence" is a promoter that is capable of conferring expression on a structural gene in a particular cell, tissue, or organ or group of cells, tissues or organs of a plant, optionally under specific conditions, however does generally not confer expression throughout the plant under all conditions. Accordingly, a regulatable promoter sequence may be a promoter sequence that confers expression on a gene to which it is operably connected in a particular location within the plant or alternatively, throughout the plant under a specific set of conditions, such as following induction of gene expression by a chemical compound or other elicitor.

Preferably, the regulatable promoter used in the performance of the present invention confers expression in a specific location within the plant, either constitutively or following induction, however not in the whole plant under any circumstances. Included within the scope of such promoters are cell-specific promoter sequences, tissue-specific promoter

sequences, organ-specific promoter sequences, cell cycle specific gene promoter sequences, inducible promoter sequences and constitutive promoter sequences that have been modified to confer expression in a particular part of the plant at any one time, such as by integration of said constitutive promoter within a transposable genetic element (*Ac*, *Ds*, *Spm*, *En*, or other transposon). The skilled craftsman will understand that a "constitutive promoter" is a promoter that is transcriptionally active in an organism, preferably a plant, during most, but not necessarily all phases of its growth and development. Similarly, the skilled craftsman will understand that a "ubiquitous promoter" is a promoter that is transcriptionally active throughout most, but not necessarily all parts of an organism, preferably a plant.

Generally by "weak promoter" is intended a promoter that drives expression of a coding sequence at a low level. By "low level" is intended at levels of about 1/10,000 transcripts to about 1/100,000 transcripts, to about 1/500,000 transcripts. Conversely, a "strong promoter" drives expression of a coding sequence at high level, or at about 1/10 transcripts to about 1/100 transcripts to about 1/1,000 transcripts.

The term "cell-specific" shall be taken to indicate that expression is predominantly in a particular cell or cell-type, preferably of plant origin, albeit not necessarily exclusively in said cell or cell-type.

Similarly, the term "tissue-specific" shall be taken to indicate that expression is predominantly in a particular tissue or tissue-type, preferably of plant origin, albeit not necessarily exclusively in said tissue or tissue-type.

Similarly, the term "organ-specific" shall be taken to indicate that expression is predominantly in a particular organ, preferably of plant origin, albeit not necessarily exclusively in said organ. "Root specific" means that the promoter is expressed in the root only and not in other tissues of the plant.

By "root-preferred" it is intended that expression of the heterologous nucleotide sequence is most abundant root, but could also have low expression levels elsewhere in the plant. While some level of expression of the heterologous nucleotide sequence occurs in other plant tissue types, expression occurs most abundantly in the root including primary, lateral and adventitious roots.

By "root" is intended any part of the root structure, including, but not limited to, the root cap, apical meristem, protoderm, ground meristem, procambium, endodermis, cortex, vascular cortex, epidermis, and the like.

Similarly, the term "cell cycle specific" shall be taken to indicate that expression is predominantly cyclic and occurring in one or more, not necessarily consecutive phases of the cell cycle albeit not necessarily exclusively in cycling cells, preferably of plant origin.

- 5 Contrarily, the term ubiquitous shall be taken to indicate that expression is throughout the entire organism.

Those skilled in the art will be aware that an "inducible promoter" is a promoter the transcriptional activity of which is increased or induced in response to a developmental, chemical, environmental, or physical stimulus. Similarly, the skilled craftsman will  
10 understand that a "constitutive promoter" is a promoter that is transcriptionally active throughout most, but not necessarily all parts of an organism, preferably a plant, during most, but not necessarily all phases of its growth and development.

Those skilled in the art will readily be capable of selecting appropriate promoter sequences for use in regulating appropriate expression of the LRR receptor-like kinase  
15 as described supra from publicly-available or readily-available sources, without undue experimentation.

Placing a nucleic acid molecule under the regulatory control of a promoter sequence, or in operable connection with a promoter sequence, means positioning said nucleic acid molecule such that expression is controlled by the promoter sequence. A promoter is  
20 usually, but not necessarily, positioned upstream, or at the 5'-end, and within 2 kb of the start site of transcription, of the nucleic acid molecule which it regulates. In the construction of heterologous promoter/structural gene combinations it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the same as the distance between that promoter and the gene it  
25 controls in its natural setting (i.e., the gene from which the promoter is derived). As is known in the art, some variation in this distance can be accommodated without loss of promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting (i.e., the gene from which it is derived).  
30 Again, as is known in the art, some variation in this distance can also occur.

"Expression" means the production of a protein or nucleotide sequence in the cell itself or in a cell-free system. It includes transcription into an RNA product, post-transcriptional

modification and/or translation to a protein product or polypeptide from a DNA encoding that product, as well as possible post-translational modifications.

"Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence  
5 "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is preferably used.

Examples of promoters suitable for use in gene constructs of the present invention  
10 include those listed in Table 3, amongst others. The promoters listed in Table 5 are provided for the purposes of exemplification only and the present invention is not to be limited by the list provided therein. Those skilled in the art will readily be in a position to provide additional promoters that are useful in performing the present invention.

In the case of constitutive promoters or promoters that induce expression throughout the  
15 entire plant, it is preferred that such sequences are modified by the addition of nucleotide sequences derived from one or more of the tissue-specific promoters listed in Table 8, or alternatively, nucleotide sequences derived from one or more of the above-mentioned tissue-specific inducible promoters, to confer tissue-specificity thereon. For example, the CaMV 35S promoter may be modified by the addition of maize *Adh1*  
20 promoter sequence, to confer anaerobically-regulated root-specific expression thereon, as described previously (Ellis *et al.*, 1987). Another example describes conferring root specific or root abundant gene expression by fusing the CaMV35S promoter to elements of the maize glycine-rich protein GRP3 gene (Feix and Wulff 2000 - WO0015662). Such modifications can be achieved by routine experimentation by those skilled in the art.

The term "terminator" refers to a DNA sequence at the end of a transcriptional unit which  
25 signals termination of transcription. Terminators are 3'-non-translated DNA sequences containing a polyadenylation signal, which facilitates the addition of polyadenylate sequences to the 3'-end of a primary transcript. Terminators active in cells derived from viruses, yeasts, moulds, bacteria, insects, birds, mammals and plants are known and  
30 described in the literature. They may be isolated from bacteria, fungi, viruses, animals and/or plants.

Examples of terminators particularly suitable for use in the gene constructs of the present invention include the *Agrobacterium tumefaciens* nopaline synthase (NOS) gene

terminator, the *Agrobacterium tumefaciens* octopine synthase (OCS) gene terminator sequence, the Cauliflower mosaic virus (CaMV) 35S gene terminator sequence, the *Oryza sativa* ADP-glucose pyrophosphorylase terminator sequence (t3'Bt2), the *Zea mays* zein gene terminator sequence, the *rbcs-1A* gene terminator, and the *rbcs-3A* gene terminator sequences, amongst others.

**Table 3. Exemplary plant-expressible promoters for use  
in the performance of the present invention**

<b>I: CELL-SPECIFIC, TISSUE-SPECIFIC, AND ORGAN-SPECIFIC PROMOTERS</b>		
<b>GENE SOURCE</b>	<b>EXPRESSION PATTERN</b>	<b>REFERENCE</b>
<i><math>\alpha</math>-amylase (Amy32b)</i>	Aleurone	Lanahan et al, Plant Cell 4:203-211, 1992; Skriver et al, Proc Natl Acad Sci USA 88:7266-7270, 1991
cathepsin $\beta$ -like gene	Aleurone	Cejudo et al, Plant Mol Biol 20:849-856, 1992
<i>Agrobacterium rhizogenes rolB</i>	Cambium	Nilsson et al, Physiol Plant 100:456-462, 1997
AtPRP4	Flowers	<a href="http://salus.mediam.edu/mmg/tierney/html">http://salus.mediam.edu/mmg/tierney/html</a>
chalcone synthase (chsA)	Flowers	Van der Meer et al, Plant Mol Biol 15:95- 109, 1990
LAT52	Anther	Twell et al, Mol Gen Genet 217:240-245, 1989
<i>apetala-3</i>	Flowers	
chitinase	fruit (berries, grapes, etc)	Thomas et al. CSIRO Plant Industry, Urrbrae, South Australia, Australia; <a href="http://winetitles.com.au/gwrdc/csh95-1.html">http://winetitles.com.au/gwrdc/csh95-1.html</a>
rbcs-3A	green tissue (eg leaf)	Lam et al, Plant Cell 2:857-866, 1990; Tucker et al., Plant Physiol 113:1303-1308, 1992
leaf-specific genes	Leaf	Baszczynski et al, Nucl Acid Res 16:4732, 1988
AtPRP4	Leaf	<a href="http://salus.mediam.edu/mmg/tierney/html">http://salus.mediam.edu/mmg/tierney/html</a>
chlorella virus adenine methyltransferase gene promoter	Leaf	Mitra and Higgins, Plant Mol Biol 26:85-93, 1994
aldP gene promoter from rice	Leaf	Kagaya et al, Mol Gen Genet 248:668-674, 1995

rbcs promoter from rice or tomato	Leaf	Kyozuka et al, Plant Physiol 102:991-1000, 1993
<i>Pinus cab-6</i>	Leaf	Yamamoto et al, Plant Cell Physiol 35:773-778, 1994
rubisco promoter	Leaf	
cab (chlorophyll a/b binding protein)	Leaf	
pea Blec4 gene	Vegetative and floral epidermal tissues	Mandaci and Dobres, Plant Mol Biol 34:961-965
SAM22	senescent leaf	Crowell et al, Plant Mol Biol 18:459-466, 1992
<i>ltp</i> gene (lipid transfer gene)		Fleming et al, Plant J 2:855-862, 1992
<i>R. japonicum nif</i> gene	nodule	United States Patent No 4 803165
<i>B. japonicum nifH</i> gene	nodule	United States Patent No 5008194
GmENOD40	nodule	Yang et al, Plant J 3:573-585, 1993
PEP carboxylase (PEPC)	nodule	Pathirana et al, Plant Mol Biol 20:437-450, 1992
Leghaemoglobin (Lb)	nodule	Gordon et al, J Exp Bot 44:1453-1465, 1993
<i>Tungro bacilliform</i> virus gene	phloem	Bhattacharyya-Pakrasi et al, Plant J 4:71-79, 1992
pollen-specific genes	pollen; microspore	Albani et al, Plant Mol Biol 15:605, 1990; Albani et al, Plant Mol Biol 18:501, 1991
Zm13	pollen	Guerrero et al, Mol Gen Genet 224:161-168, 1993
apg gene	microspore	Twel et al, Sex Plant Reprod 6:217-224, 1993
maize pollen-specific gene	pollen	Hamilton et al, Plant Mol Biol 18:211-218, 1992
Sunflower pollen-expressed gene	pollen	Baltz et al, Plant J 2:713-721, 1992
<i>B. napus</i> pollen-	pollen; anther; tapetum	Arnoldo et al, J Cell Biochem, Abstract No.

specific gene		Y101, 204, 1992
root-expressible genes	roots	Tingey et al, EMBO J 6:1, 1987
Tobacco auxin-inducible gene	root tip	Van der Zaal et al, Plant Mol Biol 16:983, 1991
$\beta$ -tubulin	root	Oppenheimer et al, Gene 63:87, 1988
Tobacco root-specific genes	root	Conkling et al, Plant Physiol 93:1203, 1990
<i>B. napus</i> G1-3b gene	root	United States Patent No 5401836
SbPRP1	roots	Suzuki et al, Plant Mol Biol 21:109-119, 1993
AtPRP1; AtPRP3	roots; root hairs	<a href="http://salus.med.unc.edu/mmg/tierney/html">http://salus.med.unc.edu/mmg/tierney/html</a>
RD2 gene	root cortex	<a href="http://www2.cnsu.edu/ncsu/research">http://www2.cnsu.edu/ncsu/research</a>
TobRB7 gene	root vasculature	<a href="http://www2.cnsu.edu/ncsu/research">http://www2.cnsu.edu/ncsu/research</a>
AtPRP4	leaves; flowers; lateral root primordia	<a href="http://salus.med.unc.edu/mmg/tierney/html">http://salus.med.unc.edu/mmg/tierney/html</a>
seed-specific genes	seed	Simon et al, Plant Mol Biol 5:191, 1985; Scofield et al, J Biol Chem 262:12202, 1987; Baszczynski et al, Plant Mol Biol 14:633, 1990
Brazil Nut albumin	seed	Pearson et al, Plant Mol Biol 18:235-245, 1992
Legumin	seed	Ellis et al, Plant Mol Biol 10:203-214, 1988
Glutelin (rice)	seed	Takaiwa et al, Mol Gen Genet 208:15-22, 1986; Takaiwa et al, FEBS Lett 221:43-47, 1987
Zeln	seed	Matzke et al, Plant Mol Biol 14:323-32 1990
NapA	seed	Stalberg et al, Planta 199:515-519, 1996
wheat LMW and HMW glutenin-1	endosperm	Mol Gen Genet 216:81-90, 1989; Nucl Acids Res 17:461-462, 1989
wheat SPA	seed	Albani et al, Plant Cell 9:171-184, 1997
cZ19B1, maize 19 kDa zeln	seed	WO0011177
mitps, maize	seed	WO0011177



myoinositol-1-Pi synthase		
wheat $\alpha$ , $\beta$ , $\gamma$ -gliadins	endosperm	EMBO J 3:1409-1415, 1984
barley <i>ltr1</i> promoter	endosperm	
barley B1, C, D, hordein	endosperm	Theor Appl Gen 98:1253-1262, 1999; Plant J 4:343-355, 1993; Mol Gen Genet 250:750-60, 1996
barley DOF	endosperm	Mena et al, Plant J 116:53-62, 1998
<i>blz2</i>	endosperm	EP99106056.7
synthetic promoter	endosperm	Vicente-Carbajosa et al, Plant J 13:629-640, 1998
rice prolamin NRP33	endosperm	Wu et al, Plant Cell Physiol 39: 885-889, 1998
rice $\alpha$ -globulin Glb-1	endosperm	Wu et al, Plant Cell Physiol 39:885-889, 1998
maize END genes	endosperm	WO0012733
barley END1	endosperm	WO9808961
barley NUC1	nucellus	WO9808961
rice OSH1	embryo	Sato et al, Proc Natl Acad Sci USA 93:8117-8122, 1996
rice $\alpha$ -globulin REB/OHP-1	endosperm	Nakase et al, Plant Mol Biol 33:513-522, 1997
rice ADP-glucose PP	endosperm	Trans Res 6:157-168, 1997
maize ESR gene family	endosperm	Plant J 12:235-246, 1997
sorgum $\gamma$ -kafirin	endosperm	Plant Mol Biol 32:1029-1035, 1996
KNOX	embryo	Postma-Haarsma et al, Plant Mol Biol 39:257-271, 1999
rice oleosin	embryo and aleuron	Wu et al, J Biochem 123:386, 1998
sunflower oleosin	seed (embryo and dry seed)	Cummins et al, Plant Mol Biol 19:873-876, 1992
<i>LEAFY</i>	shoot meristem	Weigel et al, Cell 69:843-859, 1992
<i>Arabidopsis thaliana knat1</i>	shoot meristem	Accession number AJ131822
<i>Malus domestica</i>	shoot meristem	Accession number Z71981

<i>kn1</i>		
<i>CLAVATA1</i>	shoot meristem	Accession number AF049870
stigma-specific genes	stigma	Nasrallah et al, Proc Natl Acad Sci USA 85:5551, 1988; Trick et al, Plant Mol Biol 15:203, 1990
class I patatin gene	tuber	Liu et al, Plant Mol Biol 153:386-395, 1991
PCNA rice	meristem	Kosugi et al, Nucl Acids Res 19:1571-1576, 1991; Kosugi and Ohashi, Plant Cell 9:1607-1619, 1997
Pea TubA1 tubulin	Dividing cells	Stotz and Long, Plant Mol Biol 41:601-614, 1999
<i>Arabidopsis cdc2a</i>	cycling cells	Chung and Parish, FEBS Lett 362:215-219, 1995
<i>Arabidopsis</i> Rop1A	Anthers; mature pollen + pollen tubes	Li et al, Plant Physiol 118:407-417, 1998
<i>Arabidopsis</i> AtDMC1	Meiosis-associated	Klimyuk and Jones, Plant J 11:1-14, 1997
Pea PS-IAA4/5 and PS-IAA6	Auxin-inducible	Wong et al, Plant J 9:587-599, 1996
Pea farnesyltransferase	Meristematic tissues; phloem near growing tissues; light- and sugar-repressed	Zhou et al, Plant J 12:921-930, 1997
Tobacco ( <i>N. sylvestris</i> ) cyclin B1;1	Dividing cells / meristematic tissue	Trehin et al, Plant Mol.Biol. 35:667-672, 1997
Catharanthus roseus Mitotic cyclins CYS (A-type) and CYM (B-type)	Dividing cells / meristematic tissue	Ito et al, Plant J 11:983-992, 1997
<i>Arabidopsis</i> cyc1At (=cyc B1;1) and cyc3aAt (A-type)	Dividing cells / meristematic tissue	Shaul et al, Proc Natl Acad Sci USA 93:4868-4872, 1996
<i>Arabidopsis</i> tef1 promoter box	Dividing cells / meristematic tissue	Regad et al, Mol Gen Genet 248:703-711, 1995
<i>Catharanthus roseus</i>	Dividing cells /	Ito et al, Plant Mol Biol 24:863-878, 1994

cyc07	meristematic tissue	
<b>II: EXEMPLARY CONSTITUTIVE PROMOTERS</b>		
GENE SOURCE	EXPRESSION PATTERN	REFERENCE
Actin	constitutive	McElroy et al, Plant Cell 2:163-171, 1990
CAMV 35S	constitutive	Odell et al, Nature 313:810-812, 1985
CaMV 19S	constitutive	Nilsson et al, Physiol Plant 100:456-462, 1997
GOS2	constitutive	de Pater et al, Plant J 2:837-844, 1992
Ubiquitin	constitutive	Christensen et al, Plant Mol Biol 18:675-689, 1992
rice cyclophilin	constitutive	Buchholz et al, Plant Mol Biol 25:837-843, 1994
maize histone H3	constitutive	Lepetit et al, Mol Gen Genet 231:276-285, 1992
alfalfa histone H3	constitutive	Wu et al, Nucleic Acids Res 17:3057-3063, 1989; Wu et al, Plant Mol Biol 11:641-649, 1988
actin 2	constitutive	An et al, Plant J 10:107-121, 1996
<b>III: EXEMPLARY STRESS-INDUCIBLE PROMOTERS</b>		
NAME	STRESS	REFERENCE
P5CS (delta(1)-pyrroline-5-carboxylate synthase)	salt, water	Zhang et al, Plant Sci 129:81-89, 1997
cor15a	cold	Hajela et al, Plant Physiol 93:1246-1252, 1990
cor15b	cold	Willehm et al, Plant Mol Biol 23:1073-1077, 1993
cor15a (-305 to +78 nt)	cold, drought	Baker et al, Plant Mol Biol 24: 01-713, 1994
rd29	salt, drought, cold	Kasuga et al, Nature Biotechnol 18:287-291, 1999
heat shock proteins,	heat	Barros et al, Plant Mol Biol 19 665-75,

including artificial promoters containing the heat shock element (HSE)		1992. Marrs et al, Dev Genet 14:27-41, 1993. Schoffl et al, Mol Gen Genet 217:246-53, 1989.
SmHSP (small heat shock proteins)	heat	Waters et al, J Exp Bot 47:325-338, 1996
wcs120	cold	Ouellete et al, FEBS Lett 423:324-328, 1998
ci7	cold	Kirch et al, Plant Mol Biol 33:897-909, 1997
Adh	cold, drought, hypoxia	Dolferus et al, Plant Physiol 105:1075-87, 1994
pws18	salt and drought	Joshee et al, Plant Cell Physiol 39:64-72, 1998
ci21A	cold	Schneider et al, Plant Physiol 113:335-45, 1997
Trg-31	drought	Chaudhary et al, Plant Mol Biol 30:1247-57, 1996
Osmotin	osmotic	Raghothama et al, Plant Mol Biol 23:1117-28, 1993
LapA	wounding, environmental	WO99/03977 University of California/INRA

#### IV: EXEMPLARY PATHOGEN-INDUCIBLE PROMOTERS

NAME	PATHOGEN	REFERENCE
RB7	Root-knot nematodes (Meloidogyne spp.)	US5760386 - North Carolina State University; Opperman et al, Science 263:221-23, 1994
PR-1, 2, 3, 4, 5, 8, 11	fungal, viral, bacterial	Ward et al, Plant Cell 3:1085-1094, 1991; Reiss et al 1996; Lebel et al, Plant J 16:223-233, 1998; Melchers et al, Plant J 5:469-480, 1994; Lawton et al, Plant Mol Biol, 19:735-743, 1992
HMG2	nematodes	WO9503690 - Virginia Tech Intellectual Properties Inc .
Abi3	Cyst nematodes	unpublished

	(Heterodera spp.)	
ARM1	nematodes	Barthels et al, Plant Cell 9:2119-2134, 1997 WO 98/31822 – Plant Genetic Systems
Att0728	nematodes	Barthels et al, Plant Cell 9: 2119-2134, 1997 PCT/EP98/07761
Att1712	nematodes	Barthels et al, Plant Cell 9, 2119-2134, 1997 PCT/EP98/07761
Gst1	Different types of pathogens	Strittmatter et al, Mol Plant-Microbe Interact 9:68-73, 1996
LEMMI	nematodes	WO 92/21757 – Plant Genetic Systems
CLE	geminivirus	PCT/EP99/03445 - CINESTAV
PDF1.2	Fungal including <i>Alternaria brassicicola</i> and <i>Botrytis cinerea</i>	Manners et al, Plant Mol Biol, 38:1071-1080, 1998
Thi2.1	Fungal – <i>Fusarium oxysporum f. sp. matthiolae</i>	Vignutelli et al, Plant J 14:285-295, 1998
DB#226	nematodes	Bird and Wilson, Mol Plant-Microbe Interact 7:419-442, 1994 WO 95.322888
DB#280	nematodes	Bird and Wilson, Mol Plant-Microbe Interact 7:419-442, 1994 WO 95.322888
Cat2	nematodes	Niebel et al, Mol Plant-Microbe Interact 8:371-378, 1995
□Tub	nematodes	Aristizabal et al (1996), 8 <sup>th</sup> International Congress on Plant-Microbe Interaction, Knoxville US B-29
SHSP	nematodes	Fenoll et al (1997) In: Cellular and molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grondler and S.A. Ohi (Eds.),
Tsw12	nematodes	Fenoll et al (1997) In: Cellular and

		molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grundler and S.A. Ohi (Eds.)
Hs1(pro1)	nematodes	WO 98/122335 - Jung
NsLTP	viral, fungal, bacterial	Molina and Garcia-Olmedo FEBS Lett, 316:119-122, 1993
RIP	viral, fungal	Tumer et al, Proc Natl Acad Sci USA 94:3866-3871, 1997

Those skilled in the art will be aware of additional promoter sequences and terminator sequences which may be suitable for use in performing the invention. Such sequences may readily be used without any undue experimentation.

5 In the context of the current invention, "ectopic expression" or "ectopic overexpression" of a gene or a protein are conferring to expression patterns and/or expression levels of said gene or protein normally not occurring under natural conditions. Ectopic expression can be achieved in a number of ways including operably linking of a coding sequence encoding said protein to an isolated homologous or heterologous promoter in order to  
10 create a chimeric gene and/or operably linking said coding sequence to its own isolated promoter (i.e. the unisolated promoter naturally driving expression of said protein) in order to create a recombinant gene duplication or gene multiplication effect. With "ectopic co-expression" is meant the ectopic expression or ectopic overexpression of two or more genes or proteins. The same or, more preferably, different promoters are used  
15 to confer expression of said genes or proteins.

Preferably, the promoter sequence used in the context of the present invention is operably linked to a coding sequence or open reading frame (ORF) encoding a LRR receptor-like kinase or a homolog, derivative and/or an immunologically active fragment thereof as defined supra.

20 "Dominant negative version or variant" refers to a mutant protein, which interferes with the activity of the corresponding wild-type protein.

"Downregulation of expression" as used herein means lowering levels of gene expression and/or levels of active gene product and/or levels of gene product activity. Decreases in expression may be accomplished by e.g. the addition of coding sequences  
25 or parts thereof in a sense orientation (if resulting in co-suppression) or in an antisense

orientation relative to a promoter sequence and furthermore by e.g. insertion mutagenesis (e.g. T-DNA insertion or transposon insertion) or by gene silencing strategies as described by e.g. Angell and Bailcombe (1998 - WO9836083), Lowe et al. (1989 - WO9853083), Lederer et al. (1999 - WO9915682) or Wang et al. (1999 - WO9953050). Genetic constructs aimed at silencing gene expression may have the nucleotide sequence of said gene (or one or more parts thereof) contained therein in a sense and/or antisense orientation relative to the promoter sequence. Another method to downregulate gene expression comprises the use of ribozymes, e.g. as described in Atkins et al. 1994 (WO9400012), Lenée et al. 1995 (WO9503404), Lutziger et al. 2000 (WO0000619), Prinsen et al. 1997 (WO9713865) and Scott et al. 1997 (WO9738116).

Modulating, including lowering, the level of active gene products or of gene product activity can be achieved by administering or exposing cells, tissues, organs or organisms to said gene product, a homolog, analogue, derivative and/or immunologically active fragment thereof. Immunomodulation is another example of a technique capable of downregulation levels of active gene product and/or of gene product activity and comprises administration of or exposing to or expressing antibodies to said gene product to or in cells, tissues, organs or organisms wherein levels of said gene product and/or gene product activity are to be modulated. Such antibodies comprise "plantibodies", single chain antibodies, IgG antibodies and heavy chain camel antibodies as well as fragments thereof.

Modulating, including lowering, the level of active gene products or of gene product activity can furthermore be achieved by administering or exposing cells, tissues, organs or organisms to an inhibitor or activator of said gene product or the activity thereof. Such inhibitors or activators include proteins (comprising e.g. proteinases and kinases) and chemical compounds identified according to the current invention as described supra.

In the context of the invention the term "agonist" refers to a substance that can be either a protagonist or an antagonist, i.e. can have either positive or negative effects, can be an enhancer or an inhibitor or a modulator as well.

In the context of the current invention is envisaged the downregulation of the expression of a LRR receptor-like kinase gene as defined higher. The invention further comprises downregulation of levels of a LRR receptor-like kinase or of a LRR receptor-like kinase activity whereby said LRR receptor-like kinase has been defined supra.

By "cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology" is meant that one or more developmental and/or morphological and/or biochemical and/or physiological characteristics of a plant is altered by the performance of one or more steps pertaining to the invention described herein.

5 "Cell fate" refers to the cell-type or cellular characteristics of a particular cell that are produced during plant development or a cellular process therefor, in particular during the cell cycle or as a consequence of a cell cycle process.

"Plant development" or the term "plant developmental characteristic" or similar term shall, when used herein, be taken to mean any cellular process of a plant that is involved  
10 in determining the developmental fate of a plant cell, in particular the specific tissue or organ type into which a progenitor cell will develop. Cellular processes relevant to plant development will be known to those skilled in the art. Such processes include, for example, morphogenesis, photomorphogenesis, shoot development, root development, vegetative development, reproductive development, stem elongation, flowering, and  
15 regulatory mechanisms involved in determining cell fate, in particular a process or regulatory process involving the cell cycle.

"Plant morphology" or the term "plant morphological characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to the external appearance of a plant, including any one or more structural features or combination of  
20 structural features thereof. Such structural features include the shape, size, number, position, colour, texture, arrangement, and patterning of any cell, tissue or organ or groups of cells, tissues or organs of a plant, including the root, stem, leaf, shoot, petiole, trichome, flower, petal, stigma, style, stamen, pollen, ovule, seed, embryo, endosperm, seed coat, aleurone, fibre, fruit, cambium, wood, heartwood, parenchyma, aerenchyma,  
25 sieve element, phloem or vascular tissue, amongst others.

"Plant biochemistry" or the term "plant biochemical characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to the metabolic and catalytic processes of a plant, including primary and secondary metabolism and the products thereof, including any small molecules, macromolecules or chemical  
30 compounds, such as but not limited to starches, sugars, proteins, peptides, enzymes, hormones, growth factors, nucleic acid molecules, celluloses, hemicelluloses, calloses, lectins, fibres, pigments such as anthocyanins, vitamins, minerals, micronutrients, or macronutrients, that are produced by plants.



"Plant physiology" or the term "plant physiological characteristic" or similar term will, when used herein, be understood to refer to the functional processes of a plant, including developmental processes such as growth, expansion and differentiation, sexual development, sexual reproduction, seed set, seed development, grain filling, asexual reproduction, cell division, dormancy, germination, light adaptation, photosynthesis, leaf expansion, fiber production, secondary growth or wood production, amongst others; responses of a plant to externally-applied factors such as metals, chemicals, hormones, growth factors, environment and environmental stress factors (eg. anoxia, hypoxia, high temperature, low temperature, dehydration, light, daylength, flooding, salt, heavy metals, amongst others), including adaptive responses of plants to said externally-applied factors.

Means for introducing recombinant DNA into plant tissue or cells include, but are not limited to, transformation using  $\text{CaCl}_2$  and variations thereof, in particular the method described by Hanahan (1983), direct DNA uptake into protoplasts (Krens *et al.* 1982; Paszkowski *et al.* 1984), PEG-mediated uptake to protoplasts (Armstrong *et al.* 1990) microparticle bombardment, electroporation (Fromm *et al.*, 1985), microinjection of DNA (Crossway *et al.*, 1986), microparticle bombardment of tissue explants or cells (Christou *et al.* 1988), vacuum-infiltration of tissue with nucleic acid, or in the case of plants, T-DNA-mediated transfer from *Agrobacterium* to the plant tissue as described essentially by An *et al.* (1985), Dodds *et al.* (1985), Herrera-Estrella *et al.* (1983a, 1983b). Methods for transformation of monocotyledonous plants are well known in the art and include *Agrobacterium*-mediated transformation (Cheng *et al.* 1997 - WO9748814; Hansen 1998 - WO9854961; Hiei *et al.* 1994 - WO9400977; Hiei *et al.* 1998 - WO9817813; Rikiishi *et al.* 1999 - WO9904618; Saito *et al.* 1995 - WO9506722), microprojectile bombardment (Adams *et al.* 1999 - US5969213; Bowen *et al.* 1998 - US5736369; Chang *et al.* 1994 - WO9413822; Lundquist *et al.* 1999 - US5874265/US5990390; Vasil and Vasil 1995 - US5405765; Walker *et al.* 1999 - US5955362), DNA uptake (Eyal *et al.* 1993 - WO9318168), microinjection of *Agrobacterium* cells (von Holt 1994 - DE4309203) and sonication (Finer *et al.* 1997 - US5693512).

For microparticle bombardment of cells, a microparticle is propelled into a cell to produce a transformed cell. Any suitable ballistic cell transformation methodology and apparatus can be used in performing the present invention. Exemplary apparatus and procedures are disclosed by Stomp *et al.* (U.S. Patent No. 5122466) and Sanford and Wolf (U.S.

Patent No. 4945050). When using ballistic transformation procedures, the gene construct may incorporate a plasmid capable of replicating in the cell to be transformed.

Examples of microparticles suitable for use in such systems include 1 to 5  $\mu\text{m}$  gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

A whole plant may be regenerated from the transformed or transfected cell, in accordance with procedures well known in the art. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a gene construct of the present invention and a whole plant regenerated therefrom.

The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristem, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem).

The term "organogenesis", as used herein, means a process by which shoots and roots are developed sequentially from meristematic centers.

The term "embryogenesis", as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes.

Preferably, the plant is produced according to the inventive method is transfected or transformed with a genetic sequence, or amenable to the introduction of a protein, by any art-recognized means, such as microprojectile bombardment, microinjection, *Agrobacterium*-mediated transformation (including the 'flower dip' transformation method; Bechtold and Pelletier 1998, Trieu et al. 2000), protoplast fusion, or electroporation, amongst others. Most preferably said plant is produced by *Agrobacterium*-mediated transformation.

The "seedling" is the juvenile plant that arises from the mature embryo after seed germination.

With "differentiation of a cell" it is understood that the cell develops unique features to be engaged for a specific function. Mostly differentiation is irreversible.

*Agrobacterium*-mediated transformation or agrolistic transformation of plants, yeast, moulds or filamentous fungi is based on the transfer of part of the transformation vector

sequences, called the T-DNA, to the nucleus and on integration of said T-DNA in the genome of said eukaryote.

With "*Agrobacterium*" is meant a member of the *Agrobacteriaceae*, more preferably *Agrobacterium* or *Rhizobacterium* and most preferably *Agrobacterium tumefaciens*.

- 5 With "T-DNA", or transferred DNA, is meant that part of the transformation vector flanked by T-DNA borders which is, after activation of the *Agrobacterium* *vir* genes, nicked at the T-DNA borders and is transferred as a single stranded DNA to the nucleus of an eukaryotic cell.

- When used herein, with "T-DNA borders", "T-DNA border region", or "border region" are  
10 meant either right T-DNA border (RB) or left T-DNA border (LB). Such a border comprises a core sequence flanked by a border inner region as part of the T-DNA flanking the border and/or a border outer region as part of the vector backbone flanking the border. The core sequences comprise 22 bp in case of octopine-type vectors and 25 bp in case of nopaline-type vectors. The core sequences in the right border region and  
15 left border region form imperfect repeats. Border core sequences are indispensable for recognition and processing by the *Agrobacterium* nicking complex consisting of at least VirD1 and VirD2. Core sequences flanking a T-DNA are sufficient to promote transfer of said T-DNA. However, efficiency of transformation using transformation vectors carrying said T-DNA solely flanked by said core sequences is low. Border inner and outer regions  
20 are known to modulate efficiency of T-DNA transfer (Wang et al. 1987). One element enhancing T-DNA transfer has been characterized and resides in the right border outer region and is called *overdrive* (Peralta et al. 1986, van Haaren et al. 1987).

- With "T-DNA transformation vector" or "T-DNA vector" is meant any vector encompassing a T-DNA sequence flanked by a right and left T-DNA border consisting of  
25 at least the right and left border core sequences, respectively, and used for transformation of any eukaryotic cell.

With "T-DNA vector backbone sequence" or "T-DNA vector backbone sequences" is meant all DNA of a T-DNA containing vector that lies outside of the T-DNA borders and, more specifically, outside the nicking sites of the border core imperfect repeats.

- 30 The current invention includes optimized T-DNA vectors such that vector backbone integration in the genome of a eukaryotic cell is minimized or absent. With "optimized T-DNA vector" is meant a T-DNA vector designed either to decrease or abolish transfer of vector backbone sequences to the genome of a eukaryotic cell. Such T-DNA vectors are

known to the one familiar with the art and include those described by Hanson et al. (1999) and by Stuiver et al. (1999 - WO9901563).

The current invention clearly considers the inclusion of a DNA sequence comprising the promoter sequence of the present invention encoding a LRR receptor-like kinase, homolog, derivative or immunologically active fragment thereof as defined supra, in any T-DNA vector comprising binary transformation vectors, super-binary transformation vectors, co-integrate transformation vectors, Ri-derived transformation vectors as well as in T-DNA carrying vectors used in agrolistic transformation.

With "binary transformation vector" is meant a T-DNA transformation vector comprising:

a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in the eukaryotic cell to be transformed; and  
a vector backbone region comprising at least origins of replication active in *E. coli* and *Agrobacterium* and markers for selection in *E. coli* and *Agrobacterium*.

The T-DNA borders of a binary transformation vector can be derived from octopine-type or nopaline-type Ti plasmids or from both. The T-DNA of a binary vector is only transferred to a eukaryotic cell in conjunction with a helper plasmid. Also known in the art are multiple binary vector *Agrobacterium* strains for efficient co-transformation of plants (Bidney and Scelongo 2000 - WO0018939).

With "helper plasmid" is meant a plasmid that is stably maintained in *Agrobacterium* and is at least carrying the set of *vir* genes necessary for enabling transfer of the T-DNA. Said set of *vir* genes can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

With "super-binary transformation vector" is meant a binary transformation vector additionally carrying in the vector backbone region a *vir* region of the Ti plasmid pTiBo542 of the super-virulent *A. tumefaciens* strain A281 (Hiei et al. 1994 - EP0604662, Hiei et al. 1995 - EP0687730). Super-binary transformation vectors are used in conjunction with a helper plasmid.

With "co-integrate transformation vector" is meant a T-DNA vector at least comprising:

a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in plants; and  
a vector backbone region comprising at least origins of replication active in *Escherichia coli* and *Agrobacterium*, and markers for selection in *E. coli* and *Agrobacterium*, and a set of *vir* genes necessary for enabling transfer of the T-DNA.

The T-DNA borders and said set of *vir* genes of a said T-DNA vector can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

With "Ri-derived plant transformation vector" is meant a binary transformation vector in which the T-DNA borders are derived from a Ti plasmid and said binary transformation vector being used in conjunction with a 'helper' Ri-plasmid carrying the necessary set of *vir* genes.

As used herein, the term "selectable marker gene" or "selectable marker" or "marker for selection" includes any gene which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells which are transfected or transformed with a gene construct of the invention or a derivative thereof. Suitable selectable marker genes contemplated herein include the ampicillin resistance (*Amp<sup>r</sup>*), tetracycline resistance gene (*Tc<sup>r</sup>*), bacterial kanamycin resistance gene (*Kan<sup>r</sup>*), phosphinothricin resistance gene, neomycin phosphotransferase gene (*nptII*), hygromycin resistance gene,  $\beta$ -glucuronidase (GUS) gene, chloramphenicol acetyltransferase (CAT) gene, green fluorescent protein (*gfp*) gene (Haseloff *et al*, 1997), and luciferase gene, amongst others.

With "agrolistics", "agrolistic transformation" or "agrolistic transfer" is meant here a transformation method combining features of *Agrobacterium*-mediated transformation and of biolistic DNA delivery. As such, a T-DNA containing target plasmid is co-delivered with DNA/RNA enabling in planta production of VirD1 and VirD2 with or without VirE2 (Hansen and Chilton 1996; Hansen *et al.* 1997; Hansen and Chilton 1997 - WO9712046). The present invention further describes an approach to remove from transformed cells a stably integrated foreign DNA sequence by recombination involving a recombinase and recombination sites.

With "foreign DNA" is meant any DNA sequence that is introduced in the host's genome by recombinant techniques. Said foreign DNA includes e.g. a T-DNA sequence or a part thereof such as the T-DNA sequence comprising the selectable marker in an expressible format. Foreign DNA furthermore include intervening DNA sequences as defined supra.

With "recombination event" is meant either a site-specific recombination event or a recombination event effected by transposon 'jumping'.

With "recombinase" is meant either a site-specific recombinase or a transposase.

With "recombination site" is meant either site-specific recombination sites or transposon border sequences.

With "site specific recombination event" is meant an event catalyzed by a system generally consisting of three elements: a pair of DNA sequences (the site-specific recombination sequences or sites) and a specific enzyme (the site-specific recombinase). The site-specific recombinase catalyzes a recombination reaction only  
5 between two site-specific recombination sequences depending on the orientation of the site-specific recombination sequences. Sequences intervening between two site-specific recombination sites will be inverted in the presence of the site-specific recombinase when the site-specific recombination sequences are oriented in opposite directions relative to one another (i.e. inverted repeats). If the site-specific recombination  
10 sequences are oriented in the same direction relative to one another (i.e. direct repeats), then any intervening sequences will be deleted upon interaction with the site-specific recombinase. Thus, if the site-specific recombination sequences are present as direct repeats at both ends of a foreign DNA sequence integrated into a eukaryotic genome, such integration of said sequences can subsequently be reversed by interaction of the  
15 site-specific recombination sequences with the corresponding site specific recombinase. A number of different site specific recombinase systems can be used including but not limited to the Cre/lox system of bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, the PinB, PinD and PinF from *Shigella*, and the R/Rs system of *Zygosaccharomyces rouxii*. Recombinases generally  
20 are integrases, resolvases or flippases. Also dual-specific recombinases can be used in conjunction with direct or indirect repeats of two different site-specific recombination sites corresponding to the dual-specific recombinase (Baszczynski et al. 1999 - WO9925840). The preferred site-specific recombinase systems are the bacteriophage P1 Cre/lox, the yeast FLP/FRT and the *Z. rouxii* R/Rs systems. In these systems a  
25 recombinase (Cre, FLP or R) interact specifically with its respective site-specific recombination sequence (lox, FRT, or RS respectively) to invert or excise the intervening sequences. The site-specific recombination sequences for each of these two systems are relatively short (34 bp for lox and 47 bp for FRT). Some of these systems have already been used with high efficiency in plants such as tobacco (Dale et al. 1990,  
30 Onouchi et al. 1991, Sugita et al. 2000) and *Arabidopsis* (Osborne et al. 1995, Onouchi et al. 1995). Site-specific recombination systems have many applications in plant molecular biology including methods for control of homologous recombination (e.g. Hodges et al. 1996 - US5527695), for targeted insertion, gene stacking, etc.

(Baszoczynski et al. 1999 - WO9925821) and for resolution of complex T-DNA integration patterns or for excision of a selectable marker (Ow et al. 1999 - WO9923202).

Although the site-specific recombination sequences must be linked to the ends of the DNA to be excised or to be inverted, the gene encoding the site specific recombinase  
5 may be located elsewhere. For example, the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified recombinase protein could be introduced directly into the eukaryotic cell, e.g. by micro-injection or particle bombardment. Typically, the  
10 site-specific recombinase coding region will be operably linked to regulatory sequences enabling expression of the site-specific recombinase in the eukaryotic cell.

With "recombination event effected by transposon jumping" or "transposase-mediated recombination" is meant a recombination event catalyzed by a system consisting of three elements: a pair of DNA sequences (the transposon border sequences) and a  
15 specific enzyme (the transposase). The transposase catalyzes a recombination reaction only between two transposon border sequences which are arranged as inverted repeats. A number of different transposon/transposase systems can be used including but not limited to the Ds/Ac system, the Spm system and the Mu system. These systems originate from corn but it has been shown that at least the Ds/Ac and the Spm system  
20 also function in other plants (Fedoroff et al. 1993, Schlappi et al. 1993, Van Sluys et al. 1987). Preferred are the Ds- and the Spm-type transposons which are delineated by 11 bp- and 13 bp- border sequences, respectively.

Although the transposon border sequences must be linked to the ends of the DNA to be excised, the gene encoding the transposase may be located elsewhere. For example,  
25 the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified transposase protein could be introduced directly into cells, e.g. by microinjection or by particle bombardment.

30 As part of the current invention, transposon border sequences are included in a foreign DNA sequence such that they lie outside said DNA sequence and transform said DNA into a transposon-like entity that can move by the action of a transposase.

As transposons often reintegrate at another locus of the host's genome, segregation of the progeny of the hosts in which the transposase was allowed to act might be necessary to separate transformed hosts containing e.g. only the transposon footprint and transformed hosts still containing the foreign DNA.

5 In performing the present invention, the genetic element is preferably induced to mobilize, such as, for example, by the expression of a recombinase protein in the cell which contacts the integration site of the genetic element and facilitates a recombination event therein, excising the genetic element completely, or alternatively, leaving a "footprint", generally of about 20 nucleotides in length or greater, at the original  
10 integration site. Those hosts and host parts that have been produced according to the inventive method can be identified by standard nucleic acid hybridization and/or amplification techniques to detect the presence of the mobilizable genetic element or a gene construct comprising the same. Alternatively, in the case of transformed host cells, tissues, and hosts wherein the mobilizable genetic element has been excised, it is  
15 possible to detect a footprint in the genome of the host which has been left following the excision event, using such techniques. As used herein, the term "footprint" shall be taken to refer to any derivative of a mobilizable genetic element or gene construct comprising the same as described herein which is produced by excision, deletion or other removal of the mobilizable genetic element from the genome of a cell transformed previously with  
20 said gene construct. A footprint generally comprises at least a single copy of the recombination loci or transposon used to promote excision. However, a footprint may comprise additional sequences derived from the gene construct, for example nucleotide sequences derived from the left border sequence, right border sequence, origin of replication, recombinase-encoding or transposase-encoding sequence if used, or other  
25 vector-derived nucleotide sequences. Accordingly, a footprint is identifiable according to the nucleotide sequence of the recombination locus or transposon of the gene construct used, such as, for example, a sequence of nucleotides corresponding or complementary to a *lox* site, *frt* site or RS site.

The term "cell cycle" means the cyclic biochemical and structural events associated with  
30 growth and with division of cells, and in particular with the regulation of the replication of DNA and mitosis. Cell cycle includes phases called: G0, Gap1 (G1), DNA synthesis (S), Gap2 (G2), and mitosis (M). Normally these four phases occur sequentially, however, the cell cycle also includes modified cycles wherein one or more phases are absent



resulting in modified cell cycle such as endomitosis, acytokinesis, polyploidy, polyteny, and endoreduplication.

The term "cell cycle interacting protein", "cell cycle protein" or "cell cycle control protein" as denoted herein means a protein which exerts control on or regulates or is required for the cell cycle or part thereof of a cell, tissue, organ or whole organism and/or DNA replication. It may also be capable of binding to, regulating or being regulated by cyclin dependent kinases or their subunits. The term also includes peptides, polypeptides, fragments, variant, homologs, alleles or precursors (eg preproteins or preproteins) thereof.

Cell cycle control proteins and their role in regulating the cell cycle of eukaryotic organisms are reviewed in detail by John (1981) and the contributing papers therein (Norbury & Nurse 1992; Nurse 1990; Ommrod & Francis 1993) and the contributing papers therein (Doerner et al. 1996) which are incorporated by reference. (Elledge 1996; Francis, Francis & Halford 1995; Hirt et al. 1991; Mironov et al. 1999)

The term "cell cycle control genes" refers to any gene or mutant thereof which exerts control on or are required for: chromosomal DNA synthesis and for mitosis (preprophase band, nuclear envelope, spindle formation, chromosome condensation, chromosome segregation, formation of new nuclei, formation of phragmoplast, duplication of microtubule-organizing center, etc) meiosis, cytokinesis, cell growth, endoreduplication, cell cycle control genes are also all genes exerting control on the above: homologs of CDKs, cyclins, E2Fs, Rb, CKI, Cks, and also any genes which interfere with the above, cyclin D, cdc25, Wee1, Nim1, MAP kinases, etc.

More specifically, cell cycle control genes are all genes involved in the control of entry and progression through S phase. They include, not exclusively, genes expressing "cell cycle control proteins" such as cyclin dependent kinases (CDK), cyclin dependent kinase inhibitors (CKI), D, E and A cyclins, E2F and DP transcription factors, pocket proteins, CDC7/DBF4 kinase, CDC6, MCM2-7, Orc proteins, cdc45, components of SCF ubiquitin ligase, PCNA, DNA-polymerase.

The term "cell cycle control protein" include cyclins A, B, C, D and E including CYCA1;1, CYCA2;1, CYCA3;1, CYCB1;1, CYCB1;2, CYC B2;2, CYCD1;1, CYCD2;1, CYCD3;1, and CYCD4;1 (Evans et al. 1983; Francis et al. 1998; Labbe et al. 1989; Murray & Kirschner 1989; Renaudin et al. 1996; Soni et al. 1995; Sorrell et al. 1999; Swenson, Farrell, & Ruderman 1986) cyclin dependent kinase inhibitor (CKI) proteins such as ICK1

- (Wang, Fowke, & Crosby 1997), FL39, FL66, FL67 (PCT/EP98/05895), Sic1, Far1, Rum1, p21, p27, p57, p16, p15, p18, p19 (Elledge 1996; Pines 1995), p14 and p14ARF; p13suc1 or CKS1A (De Veylder et al. 1997; Hayles & Nurse 1986) and nim-1 (Fantes 1989; Russell & Nurse 1986; Russell & Nurse 1987b; Russell & Nurse 1987a; Russell & Nurse 1986) homologues of Cdc2 such as Cdc2MsB (Hirt et al. 1993) CdcMs kinase (Bogre et al. 1997) cdc2 T14Y15 phosphatases such as Cdc25 protein phosphatase or p80cdc25 (Bell et al. 1993; Elledge 1996; Kumagai & Dunphy 1991; Russell et al. 1986) and Pyp3 (Elledge 1996) cdc2 protein kinase or p34cdc2 (Colasanti, Tyers, & Sundaresan 1991; Feiler & Jacobs 1990; Hirt et al. 1991; John, Sek, & Lee 1989; Lee & Nurse 1987; Nurse & Bissett 1981; Ormrod et al. 1993) cdc2a protein kinase (Hemerly et al. 1993) cdc2 T14Y15 kinases such as wee1 or p107wee1 (Elledge 1996; Russell et al. 1986; Russell et al. 1987b; Russell et al. 1987a; Sun et al. 1999) mik1 (Lundgren et al. 1991) and myt1 (Elledge 1996); cdc2 T161 kinases such as Cak and Cln (Elledge 1996); cdc2 T161 phosphatases such as Kap1 (Elledge 1996); cdc28 protein kinase or p34cdc28 (Nasmyth 1993; Reed, Hadwiger, & Lorincz 1985) p40MO15 (Fesquet et al. 1993; Poon et al. 1993) chk1 kinase (Zeng et al. 1998) cds1 kinase (Zeng et al. 1998) growth-associated H1 kinase (Labbe et al. 1989; Lake & Salzman 1972; Langan 1978; Zeng et al. 1998) MAP kinases described by (Binarova et al. 1998; Bögre et al. 1999; Calderini et al. 1998; Wilson et al. 1999).
- Other cell cycle control proteins that are involved in cyclin D-mediated entry of cells into G1 from G0 include pRb (Huntley et al. 1998; Xie et al. 1996) E2F, RIP, MCM7 and potentially the pRb-like proteins p107 and p130.
- Other cell cycle control proteins that are involved in the formation of a pre-replicative complex at one or more origins of replication, such as, but not limited to, ORC, CDC6, CDC14, RPA and MCM proteins or in the regulation of formation of this pre-replicative complex, such as, but not limited to, the CDC7, DBF4 and MBF proteins.
- For the present purpose, the term "cell cycle control protein" shall further be taken to include any one or more of those proteins that are involved in the turnover of any other cell cycle control protein, or in regulating the half-life of said other cell cycle control protein. The term "protein turnover" is to include all biochemical modifications of a protein leading to the physical or functional removal of said protein. Although not limited to these, examples of such modifications are phosphorylation, ubiquitination and proteolysis. Particularly preferred proteins which are involved in the proteolysis of one or

more of any other of the above-mentioned cell cycle control proteins include the yeast-derived and animal-derived proteins, Skp1, Skp2, Rub1, Cdc20, cullins, CDC23, CDC27, CDC16, and plant-derived homologues thereof (Cohen-Fix & Koshland 1997; Hochstrasser 1998; Krek 1998; Lisztwan et al. 1998) and Plesse et al in (Francis et al. 1998)).

For the present purpose, the term "cell cycle control genes" shall further be taken to include any one or more of those gene that are involved in the transcriptional regulation of cell cycle control gene expression such as transcription factors and upstream signal proteins. Additional cell cycle control genes are not excluded.

For the present purpose, the term "cell cycle control genes" shall further be taken to include any cell cycle control gene or mutant thereof, which is affected by environmental signals such as for instance stress, nutrients, pathogens, or by intrinsic signals such as the animal mitogens or the plant hormones (auxins, cytokinins, ethylene, gibberellic acid, abscisic acid and brassinosteroids).

The term "cell cycle progression" refers to the process of passing through the different cell cycle phases. The term "cell cycle progression rate" accordingly refers to the speed at which said cell cycle phases are run through or the time spans required to complete said cell cycle phases.

With "pathogen" is meant those organisms that have a negative effect on the physiological state of the plant or a part thereof. Some pathogens are for instance viruses, bacteria, fungi, and parasitic plants. with plant "pests" is meant the group of nematodes as well as insects, which are able to attack and to damage the plant.

"Plant cell" comprises any cell derived from any plant and existing in culture as a single cell, a group of cells or a callus. A plant cell may also be any cell in a developing or mature plant in culture or growing in nature.

"Plants" comprises all plants, including monocotyledonous and dicotyledonous plants.

With "yeast two-hybrid assay" is meant an assay that is based on the observation that many eukaryotic transcription factors comprise two domains, a DNA-binding domain (DB) and an activation domain (AD) which, when physically separated (i.e. disruption of the covalent linkage) do not effectuate target gene expression. Two proteins able to interact physically with one of said proteins fused to DB and the other of said proteins fused to AD will re-unite the DB and AD domains of the transcription factor resulting in target gene expression. The target gene in the yeast two-hybrid assay is usually a

reporter gene such as the  $\beta$ -galactosidase gene. Interaction between protein partners in the yeast two-hybrid assay can thus be quantified by measuring the activity of the reporter gene product (Bartel and Fields 1997). Alternatively, a mammalian two-hybrid system can be used which includes e.g. a chimeric green fluorescent protein encoding reporter gene (Shioda et al. 2000). Yet another alternative consists of a bacterial two-hybrid system using e.g. *HIS* as reporter gene (Joung et al. 2000).

The term "fragment of a sequence" or "part of a sequence" means a truncated sequence of the original sequence referred to. The truncated sequence (nucleic acid or protein sequence) can vary widely in length; the minimum size being a sequence of sufficient size to provide a sequence with at least a comparable function and/or activity or the original sequence referred to, while the maximum size is not critical. In some applications, the maximum size usually is not substantially greater than that required to provide the desired activity and/or function(s) of the original sequence. Typically, the truncated amino acid or nucleotide sequence will range from about 5 to about 60 amino acids in length. More typically, however, the sequence will be a maximum of about 50 amino acids in length, preferably a maximum of about 60 amino acids. It is usually desirable to select sequences of at least about 10, 12 or 15 amino acids or nucleotides, up to a maximum of about 20 or 25 amino acids or nucleotides.

Furthermore, folding simulations and computer redesign of structural motifs of the protein of the invention can be performed using appropriate computer programs (Olszewski et al. 1996, Hoffman et al. 1995). Computer modeling of protein folding can be used for the conformational and energetic analysis of detailed peptide and protein models (Monge et al. 1995, Renouf et al. 1995). In particular, the appropriate programs can be used for the identification of interactive sites of the LRR receptor-like kinase of the present invention by computer assisted searches for complementary peptide sequences (Fassina and Melli 1994). Further appropriate computer systems for the design of protein and peptides are described in the prior art, for example in Berry and Brenner (1994), Wodak (1987), Pabo and Suchanek (1986). The results obtained from the above-described computer analysis can be used for, e.g. the preparation of peptidomimetics of the protein of the invention or fragments thereof. Such pseudopeptide analogues of the natural amino acid sequence of the protein may very efficiently mimic the parent protein (Benkirane et al. 1996). For example, incorporation of easily available achiral  $\omega$ -amino acid residues into a protein of the invention or a

fragment thereof results in the substitution of amino bonds by polymethylene units of an aliphatic chain, thereby providing a convenient strategy for constructing a peptidomimetic (Banerjee et al. 1996). Superactive peptidomimetic analogues of small peptide hormones in other systems are described in the prior art (Zhang et al. 1996). Appropriate peptidomimetics of the protein of the present invention can also be identified by the synthesis of peptidomimetic combinatorial libraries through successive amine alkylation and testing the resulting compounds, e.g., for their binding, kinase inhibitory and/or immunological properties. Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh et al. (1996) and Dörner et al. (1996).

Furthermore, a three-dimensional and/or crystallographic structure of the protein of the invention can be used for the design of peptidomimetic inhibitors of the biological activity of the protein of the invention (Rose et al. 1996, Ruterber et al. 1996).

To align sequences we used the Clustal V method described in Higgins and Sharp (Higgins DG & Sharp PM, *Comput Appl Biosci.* 1989 Apr;5(2)151-3). The Clustal Method groups sequences into clusters by examining sequence distances between all pairs. Clusters are aligned as pairs then collectively as sequence groups to produce the overall alignment. After the multiple alignment is completed, a neighbor-joining method is employed to reconstruct phylogeny for the putative alignment.

#### 1. Residue weight table

The Clustal method uses weight tables to construct multiple alignments. Residue weight tables are used in scoring protein and nucleotide alignments so that slightly mismatched residues, such as Ile vs. Val amino acids or G vs. R nucleotides, are not scored the same as total mismatches.

If your sequences are protein, or a mixture of protein and DNA, choose between PAM250, PAM100, Structural and Identity.

1. iPAM250 is recommended for the Clustal method and is ideal for longer sequences or highly diverged sequences.
2. PAM100 is also used with the Clustal method and is most accurate for shorter sequences or sequences having little divergence.
3. iStructural is recommended for the Jotun Hein method and gives higher scores to residues that are chemically and spatially similar.

4. Identity may be used with Clustal or Jotun Hein methods, but is recommended only for the Clustal method. The table scores non-identities as zero, and identities as a default value of 10.

## 2. Method Parameters

*A. Pairwise alignment parameters may include K-tuple, Gap Penalty, Window and Diagonals Saved:*

1. Specify the K-tuple, expressed as the unit's length in nucleotides. Any stretch of K or more adjacent nucleotides that form an exact match in both sequences is a K-tuple. A lower K-tuple will find smaller matching regions; a higher value will find fewer, but better, matching regions of similarity.
2. Type in the Gap Penalty, the amount deducted from the score for each gap in the alignment. Gaps of different sizes carry the same penalty.
3. Specify the Window, the range the program searches for all possible or alternate K-tuples. An extremely high value may slow an alignment by considering subalignments far from the main diagonal.
4. Type in the number of Diagonals Saved. In the pairwise calculation, the best scoring diagonals for each pair in the matrix are saved and used in the clustering and scoring of ancestors.

*B. Multiple alignment parameters include Gap Penalty and Gap Length Penalty.*

1. Specify the Gap Penalty, the amount deducted from the score for each gap in the alignment. Gaps of different sizes carry the same penalty.
2. Specify the Gap Length Penalty, the value which will be deducted from the alignment score after first multiplying it by the number of gaps. Longer gaps have a greater penalty than shorter gaps.

Whenever in the current description reference is made to a "phylogenetic tree", this phylogenetic tree is used to view phylogenetic relationships compatible with multiple sequence alignments. The length of each pair of branches represents the distance between sequence pairs, while the units at the bottom of the tree indicate the number of substitution events.

The phylogenetic tree gives a first estimate of the relationships between homologous sequences. It is highly recommended that your data be evaluated using additional methods before final conclusions are made about evolutionary relationships.

You can switch the view between Balanced Branches (phenogram) or Unbalanced Branches (cladogram). A phenogram averages the distances between ancestors in the tree. Dotted lines indicate a negative branch length, which may be a byproduct of averaging. In the cladogram, branch distances correspond to sequence divergence.

The compounds to be obtained or identified in the methods of the invention can be compounds that are able to bind to any of the nucleic acids, peptides or proteins of the invention. Other interesting compounds to be identified are compounds that modulate the expression of the genes or the proteins of the invention in such a way that either the expression of said gene or protein is enhanced or decreased by the action of said compound. Alternatively the compound can exert his action by directly or indirectly enhancing or decreasing the activity of any of the proteins of the invention.

Said compound or plurality of compounds may be comprised in, for example, samples, e.g., cell extracts from, e.g., plants, animals or microorganisms. Furthermore, said compound(s) may be known in the art but hitherto not known to be capable of suppressing or activating cell cycle interacting proteins. The reaction mixture may be a cell free extract of may comprise a cell or tissue culture. Suitable set ups for the method of the invention are known to the person skilled in the art and are, for example, generally described in Alberts et al., *Molecular Biology of the Cell*, third edition (1994), in particular Chapter 17. The plurality of compounds may be, e.g., added to the reaction mixture, culture medium or injected into the cell.

If a sample containing a compound or a plurality of compounds is identified in the method of the invention, then it is either possible to isolate the compound from the original sample identified as containing the compound capable of acting as an agonist, or one can further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. Depending on the complexity of the samples, the steps described above can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances or similar chemical and/or physical properties, and most

preferably said substances are identical. Preferably, the compound identified according to the above described method or its derivative is further formulated in a form suitable for the application in plant breeding or plant cell and tissue culture.

5 The invention, now being generally described, will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.



## DESCRIPTION OF FIGURES

### Figure 1. Models for stem-cell maintenance in the root and shoot meristem.

(a) In the root meristem, the quiescent center (QC) functions as an organizing center. Those cells in immediate contact with the QC are stem cells (dark shading), which are initials for all cell files: endodermis (end), cortex (cort), epidermis (epi), lateral root cap (lrc) and columella (col). (b) In the shoot meristem, WUS is functionally equivalent to the QC in the root meristem in maintaining the presence of stem cells (dark shading). Long-range signaling for stem cell maintenance is also postulated for ZLL, which is expressed in the vascular cells. CLV3 is expressed in the presumptive stem cells that the initials for the L1 (epidermis), L2 (subepidermal tissues, i.e. mesophyll cells) and L3 (subepidermal tissues, i.e. vascular cells) tissue layers (Schnittger, Schellmann, & Hulskamp 1999). CLV1 is found in the L3 tunica layer of the central zone.

**Figure 2.** List of the degenerative primers used for RT-PCR of root-specific members of the LRR receptor kinase family.

### Figure 3. Differential RT-PCR on hypocotyl/cotyledon (C) and roottip (R) RNA.

The PCR fragments obtained upon amplification of R and C cDNAs with degenerate primers are separated on agarose gel. DNA fragments of different size can be detected in the R<sub>22</sub> compared to C<sub>22</sub> PCRs.

M represents the 100 bp marker (Boehringer). C and R indicated the origin of the cDNA sample used in the PCR and the numbers below indicate the primer combination used, e.g. C<sub>23</sub> represents hypocotyl/cotyledon cDNA amplified with primers LRR2 and KIN3.

### Figure 4. RT - PCR analysis of *RCH1* expression.

Total RNA was isolated from flower (F), hypocotyl/cotyledon (C) and roottips (R) and *RCH1* mRNA was amplified for 26 cycles with RT-PCR using the RCH1F and RCH1R primers. As a control for cDNA quality and quantity as well as relative expression level, ubiquitin (Ubi) mRNA was amplified for 20 cycles under the same conditions using the UBIF and UBIR primers.

**Figure 5. The nucleotide sequence of the RCH1 operon.**

This figure shows the nucleotide sequence of SEQ ID NO 1 that contains the genomic RCH1 clone.

**Figure 6. The nucleotide sequence of the RCH1 promoter region.**

This fragment of the genomic RCH1 clone is about 3498 base pairs and is identified as SEQ ID NO 2. Two putative ATG start codons of the RCH1 protein on position 3479-3481 and on position 3491-3493 could be identified and are represented in bold. The underlined sequence corresponds to SEQ ID NO 18.

It is also known that there is a gene in the reverse orientation in this promoter region. This gene's open reading frame starts at the position 1759-1757 with the ATG (bold), contains two introns, and ends at position 350-348 with a TGA codon (bold). A homology search of the protein sequence of this antisense gene in the RCH1prom against the protein database with the program BLASTP 2.0.8 (Altschul et al. 1997) produced a hypothetical protein (Accession number AC002392, clone T20K24.8) that is a putative anthranilate N-hydroxycinnamoyl/benzoyltransferase from *Arabidopsis thaliana*. Both sequences show 38% identity and 54% of their amino acids are homologous.

**Figure 7. Promoter activity of the RCH1 promoter – GUS fusion construct in the *Arabidopsis thaliana* roots.**

Seedlings of different age were grown and stained with Glucuronidase (GUS). The expression of the GUS reporter gene can be seen in the root meristem of the main and the lateral roots, as well as in the vascular tissue of the root and the lateral root primordia. (A) 1 day old seedling stained over night, (B) higher magnification of A, (C) 2 days old seedling stained for 5 hours, (D) main root from 1 day old seedling stained for 1 hour, (E) lateral root primordia from 1 week old seedling stained for 5 hours, (F, G, H) lateral roots from an 8 days old seedling stained for 5 hours.

**Figure 8. The nucleotide sequence of the RCH1 gene.**

This fragment of the genomic RCH1 clone is about 3960 base pairs long and is identified as SEQ ID NO 3. Two putative ATG start codons of the RCH1 protein on position 1-3 and on position 13-15 could be identified and are represented in bold. Also the TAA

stopcodon on position 3958-3960 is represented in bold. There is a putative intron estimated to range from position 2900 (CAG^GTA) to position 3452 (CAG^AAT) with the program <http://www.CBS.DTU.DK/services/NetGene2>. The putative splice-sites are underlined.

5

**Figure 9. The amino acid sequence of the open reading frame encoding RCH1 protein.**

Two putative start methionins of the RCH1 protein on position 1 and on position 4 could be identified. An ORF of 1135 amino acids is identified and represented in SEQ ID NO 4.

10 The ORF of 1131 amino acids and starting at the methionine on position 4 (M) corresponds with SEQ ID NO 19.

The boxed peptide RYKIILGAAQGLAYLHHDCVPPIVH corresponds with SEQ ID NO 5.

**Figure 10. Amino acid alignment of the RCH1 protein the *Arabidopsis thaliana* CLV1 receptor kinase like protein.**

15 The amino acid sequences of the RCH1 and of CLV1 (Genbank accession number AAB58929) were aligned using the program GAP as described in example 3. Match display thresholds for the alignment(s) are: | = IDENTITY, : = 2, . = 1.

20 **Figure 11. Phylogenetic tree of CLV1 homologues.**

The novel RCH1 sequence was aligned with the CLV1 (Genbank accession number U96879.1), the known CLV1 homologues BRI1 (Genbank accession number AF017056), HAE (GenPept accession number S27756) and ER (Genbank accession number D83257) as well as with 4 other related proteins that were retrieved from the public database searched with the RCH1 sequence using the BlastP program. The multiple alignment and phylogenetic tree of said sequences were calculated with the Clustal V method as described in example 4. Based on the sequence, RCH1 is more homologue to CLV1 than other known CLV-homologues.

30 **Figure 12. Phylogenetic tree of LRR receptor like kinases**

The novel RCH1 sequence was aligned with CLV1, its homologues BRI, HAE, ER and other LRR receptor like kinases retrieved from the public database. Also plant disease resistance genes, which comprise a leucine-rich repeat and which are receptor-like were

included in the alignment (RPS5, RPS2, RPP8, RPM1, RPS4, N, RPP5, Meyers *et al.*, 1999, *Plant J.*, Nov20(3): 317-32). The multiple alignment and phylogenetic tree of said proteins were calculated with the Clustal V method as described in example 4. Because of the close proximity of RCH1 to CLV1 in the phylogenetic tree, there is more evolutionary relationship between RCH1 and clavata, than with other kinds of LRR receptor like proteins such as plant disease resistance genes.

**Figure 13. The nucleotide sequence of a minimal promoter sequence of RCH1**

This figure shows the nucleotide sequence of SEQ ID NO. 18 that contains a minimal promoter sequence of the RCH1 clone. This fragment of the genomic RCH1 clone is about 842 base pairs. This sequence is deduced from the large promoter region identified as SEQ ID NO 2 (see example 7). The minimal promoter region correlates with the base pairs ranging from -1 of the first ATG defined in SEQ ID NO. 2 to -842 of the first ATG defined in SEQ ID NO. 2.

**Figure 14. Outprint of GenBank accession number AB017061 on December 4, 2000.**

**Figure 15. RCH1prom::GUS expression in embryo's**

Arabidopsis embryo transformed with RCH1prom::GUS. Embryo's were stained with beta glucuronidase. The RCH1 promoter is active very early in plant development and is only active in the root in the embryonic phase (A). On a stronger magnification (B) it is visible that the root specific expression is limited to the meristematic zone of the embryonic root.

**Figure 16. RCH1::GFP expression in roots**

RCH1 promoter is strongly active in endodermis, cortex, epidermis, lateral root cap, in short: in the "division zone". The RCH1 promoter is also active but in a low manner, in the quiescent center and in the vascular tissue.

**Figure 17. Amino acid sequence of RCH1 homologues of Table 4**

## EXAMPLES

Unless stated otherwise in the Examples, all recombinant DNA techniques are performed according to protocols as described in Sambrook et al. (1989), Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, NY or in Volumes 1 and 2 of Ausubel et al., (1994), Current Protocols in Molecular Biology, Current Protocols. Standard materials and methods for plant molecular work are described in Plant Molecular Biology Labfax (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd. (UK) and Blackwell Scientific Publications (UK).

### Example 1: Isolation of the RCH1 operon from root tissue of *Arabidopsis thaliana*.

Plant organ development depends on the presence of distally positioned groups of continuously dividing cells, the shoot and root apical meristems. Laser ablation studies suggest that a balance between signals for proper differentiation and short range signals from the mitotically inactive quiescent centre (QC), required to keep cells less differentiated, is important to maintain the root meristem (van den Berg et al. 1995; van den Berg et al. 1997). In the shoot meristem a balance between the *WUSCHEL* and *CLAVATA* (*CLV*) genes has been implicated to play an essential role in regulation cell division and differentiation (Clark et al. 1993; Fletcher et al. 1999; Schoof et al. 2000). The *CLV1* gene encodes a leucine rich repeat (LRR) receptor kinase (Clark, Williams, & Meyerowitz 1997), a member of a large gene family (Lease, Ingham, & Walker 1998). Other members are *ERECTA* (*ER*) (Torii et al. 1996), *BRASSINOLIDE INSENSITIVE* (*BRI1*) (Li et al. 1997) and *HAESA* (*HAE*) (Jinn et al. 2000), genes which have a function in plant development

We investigated whether root specific *CLV1* homologs exist using a differential RT-PCR based approach. PCR products were generated from root tip and hypocotyl/cotyledon cDNA pools using degenerate primers, resulting in the isolation of the root specific *ROOT CLAVATA HOMOLOG 1* (*RCH1*). The promoter region of this gene was translationally fused to  $\beta$ -glucuronidase-green fluorescent protein (*GUS/GFP*) conferring root specific expression of these reporter genes (see Example 2).

### Designing degenerate primers

To RT-PCR root specific members of the LRR receptor kinase gene family we compared the sequences of the *CLV1*, *ER* and *HAE* genes and their encoded proteins. The degenerate LRR primers were designed against the NxLxGxIP encoding region of the xLxxNxLxGxIPxxLxxLxxLxxL consensus for the LRR receptor-like kinases. Degenerate kinase primers were designed for different homologous regions of the kinase domain. The primers are listed in Figure 2.

### RNA Isolation

For each RNA isolation the roottips and the hypocotyl/cotyledon part including the shoot apical meristem of *Arabidopsis thaliana* (Col0) plants, 4 days after germination, were collected.

Total RNA was isolated (Pawlowski et al. 1994) and chromosomal DNA contamination was removed upon Dnase I (Promega, Madison, WI) treatment, following the instructions of the suppliers. The amount and quality of RNA was determined using spectrophotometer and agarose gel electrophoresis (data not shown).

### Differential RT-PCR

Reverse transcription (RT) was performed using 5 µg of root tip (R) and hypocotyl/cotyledon (C) total RNA in a 20 µl reaction volume with 0.5 µg oligodT<sub>12-18</sub> (Pharmacia, New Jersey, USA) and 200 U Superscript II (LifeTechnologies, Paisley, UK), according to the instructions of the supplier. Following first strand cDNA synthesis the cDNA samples were treated with RNase H (LifeTechnologies) as instructed by the supplier and subsequently diluted to a total volume of 100 µl with water.

Amplification reactions were performed with all degenerate LRR and KIN primer pairs (Figure 2). For the polymerase chain reaction (PCR) 2.5 µl of the R or C cDNA sample was amplified in a total volume of 50 µl reaction volume containing 100 µM dNTPs, 100 ng LRR and KIN primer, 1 U Taq polymerase (Boehringer, Ingelheim, Germany) and its accompanying buffer. The cDNAs were amplified during 40 cycles (1 min at 94°C, 1 min at 45-55°C, 3 min at 72°C).

### Cloning and sequencing

The amplified R and C cDNAs were separated and compared with agarose gel electrophoresis (Figure 3). The root specific fragments, e.g. fragments present in R<sub>22</sub> and not in C<sub>22</sub>, were isolated from gel, cloned into the pGEM-T vector (Promega) and transformed into *E. coli* (strain DH5 $\alpha$ ), using standard cloning procedures.

Several PCR fragments of the same size but from different members of the LRR receptor kinase gene family may be generated using degenerate primer pairs. Therefore, 24 randomly picked colonies obtained after transformation were used in a "colony PCR". Aliquots of the amplified inserts were separated on agarose gel followed by Southern transfer to a Supercharge N+ membrane (Schleicher & Schuell, Dassel, Germany) to create a "colony blot". An aliquot of one insert was labeled with <sup>32</sup>P-dCTP and hybridized to the colony blot to determine the number of clones it represented in the pool of 24. This procedure was repeated with other labeled inserts until all clones were identified.

A single representative of each pool of clones was sequenced and the sequence was analyzed using the BLAST program. One of the clones sequenced was named *RCH1* and showed strong homology to plant LRR receptor kinases.

### RT-PCR

The relative expression levels of *RCH1* in root tip (R), hypocotyl/cotyledon (C) and flowers (F) were determined using RT-PCR. Primers specific for the *RCH1* sequence were designed (RCH1F and RCH1R, Figure 2). The PCR samples were analyzed with agarose gel electrophoresis. *Ubiquitin (Ubi)* mRNA was used as an internal control (Horvath et al. 1993). Polyubiquitin genes consist of multiple units (Callis et al. 1995) and the *Ubi* primers hybridize with the ends of each single unit.

Independently isolated total RNA was reverse transcribed to obtain cDNA and for the PCR reaction 2  $\mu$ l of the cDNA sample was amplified in a total volume of 50  $\mu$ l as described above for the differential RT-PCR. The *RCH1* and *Ubi* cDNAs were amplified during 26 or 20 cycles (30 sec at 94°C, 30 sec at 56°C, 30 sec at 72°C), respectively (Figure 4). *RCH1* expression was only observed in root tip (R).

### Example 2: Promoter isolation and promoter fusion construct.

To determine if the *RCH1* promoter region is sufficient to confer root specific expression

of a "gene of interest" we decided to fuse this promoter to the Glucuronidase/Green Fluorescent Protein (*GUS/GFP*) reporter gene. The *RCH1* clone was used to isolate a phage containing the corresponding genomic region from a  $\lambda$ GEM (Promega, Madison, WI) based genomic library. The  $\lambda$ -GEM11 phage containing an insert of about 14550 nucleotides comprising the *Arabidopsis thaliana* *RCH1* promoter and *RCH1* gene and also the surrounding genomic sequences was deposited on October 25, 2000 at the BCCM-LMBP plasmid collection and was named "lambda phage *RCH1* genomic". This deposit was given the accession number LMBP 5582CB by the international depositary authority. A restriction map was constructed of the genomic clone and a 7 kb *Xho*I fragment, containing 3.5 kb promoter and part of the *RCH1* coding sequence, was subcloned into the expression vector pBS (Stratagene, La Jolla, CA). The 3.5 kb promoter region including the first amino acids of the putative *RCH1* ORF is set forth in Figure 5 and is identified as SEQ ID NO 2. This promoter region was amplified using the primers M13F and PRCH1R and Pfu Taq polymerase (Stratagene, La Jolla, CA) in the manufacturer specified conditions. The promoter region was subsequently cloned into the binary pCambia3383Xb vector (Roberts et al. 2000) creating a translational fusion with the *GUS/GFP* reporter gene.

#### **GUS histochemical assay.**

*Arabidopsis thaliana* plants were transformed with this construct according to the vacuum transformation method (Bechtold & Pelletier 1998). Transformants were selected for phosphinothricin-tripeptide (PTT, Duchefa, NL) resistance. Histochemical localization of  $\beta$ -Glucuronidase activity (GUS) was performed using the substrate 5-bromo-4-chloro-3-indolyl glucuronide (X-gluc, Biosynth AG, Staad, Switzerland). X-gluc was dissolved in *n*-dimethyl-formamide and diluted to 0.5 mg/ml in 50 mM NaPO<sub>4</sub>, pH 7.2 with 0.1% Triton X-100. Oxidative dimerization of the produced indoxyl derivative was enhanced by adding the oxidation catalyst K<sup>+</sup> ferricyanide/ferrocyanide to a final concentration of 0.5 mM (0.5 mM K<sub>4</sub>Fe(CN)<sub>6</sub>·H<sub>2</sub>O/0.5 mM K<sub>3</sub>Fe(CN)<sub>6</sub>). The GUS reaction was incubated at 37°C over several time periods, and pictures were taken from the cloured plant-tissues (Figures 7 and 15). In Figure 7 the very specific expression pattern of the *RCH1* promoter is visualized by the GUS reporter gene activity. Persons skilled in the art will also agree to the fact that the expression level of the *RCH1* promoter is very high, since after one hour the saturation



**Example 5: Root-specific reporter gene expression in Rice mediated by the *Arabidopsis thaliana* RCH 1 promoter.**

CLAVATA homologs are found in many other plant species from both the groups of monocots and dicots. Therefore one can assume that also the promoter of RCH1 is functional in those other plant species, and possibly exerts the same specific expression-pattern. To investigate the feasibility of root meristem-specific activation of the *Arabidopsis thaliana* RCH1 promoter in monocots, the aforementioned promoter-reporter gene construct, wherein the GUS/GFP chimeric gene is operably linked to the RCH1 promoter, will be transformed to rice using the standard transformation procedures well known to the persons skilled in the art and outlined in the following paragraph. After several time periods ranging from 1 day to 1 or more weeks, the seedling will be checked for the expression of the GUS reporter gene. This will be done by growing the seedlings in organogenesis medium, and staining them with glucuronidase for several time periods ranging from 1 hour to 5 hours.

***Agrobacterium*-mediated rice transformation**

The Promoter-GUS hybrid gene outlined in Example 2 can be transformed to *Agrobacterium tumefaciens* strain LBA4404 or C58 by means of electroporation and subsequently transformed bacterial cells can be selected on a solid agar medium containing the appropriate antibiotics.

For demonstration of root-specific GUS expression with the promoter of the current invention, 309 mature dry seeds of the rice japonica cultivars Nipponbare or Taipei are dehusked, sterilised and germinated on a medium containing 2,4-dichlorophenoxyacetic acid (2,4-D). After incubation in the dark for four weeks, embryogenic, scutellum-derived calli are excised and propagated on the same medium. Selected embryogenic callus is then co-cultivated with *Agrobacterium*. Co-cultivated callus is grown on 2,4-D-containing medium for 4 to 5 weeks in the dark in the presence of a suitable concentration of the appropriate selective agent. During this period, rapidly growing resistant callus islands develop. After transfer of this material to a medium with a reduced concentration of 2,4-D and incubation in the light, the embryogenic potential is released and shoots develop in the next four to five weeks. Shoots are excised from the callus and incubated for one week on an auxin-containing medium from which they can be transferred to the soil. Hardened shoots are grown under high humidity and short days in a phytotron. Seeds

level of staining is reached in the meristem of the main root from a 1 week old seedling (Figure 7 D).

RCH1::GFP expression in roots is shown in Figure 16.

### 5 **Example 3: Alignment.**

Sequence alignment of one complete register sequence against a target sequence as in figure 10 was done using the program GAP of the GCG package. The algorithm of Needleman and Wunsch is applied here to find the alignment of two complete sequences (Needleman and Wunsch, JMB 48(3): 443-453; 1970). The used parameters during the alignment were Gap Weight: 8, Average Match: 2.912, Length Weight: 2, Average Mismatch: -2.003, Quality: 1388, Length: 1149, Ratio: 1.416, and Gaps: 25.

### **Example 4: Phylogenetic tree.**

For sequence comparison of proteins as in figure 11 and 12, the Clustal V method to align all the sequences was used (Higgins DG & Sharp PM, *Comput Appl Biosci.* 1989 Apr;5(2):151-3). The Clustal Method groups sequences into clusters by examining sequence distances between all pairs. Clusters are aligned as pairs then collectively as sequence groups to produce the overall alignment. After the multiple alignment is completed, a neighbor-joining method is employed to reconstruct phylogeny for the putative alignment.

The Clustal method uses weight tables to construct multiple alignments. Residue weight tables are used in scoring protein and nucleotide alignments so that slightly mismatched residues, such as Ile vs. Val amino acids or G vs. R nucleotides, are not scored the same as total mismatches. For the alignment and pylogenetic trees in figure 11 and 12 the PAM250 weight table was used. Other method parameters for pairwise alignment were K-tuple: 1, Gap Penalty: 3, Window: 5 and Diagonals: 5. Other method parameters for multiple alignment include Gap Penalty: 10 and Gap Length Penalty: 10.

To view phylogenetic relationships compatible with multiple sequence alignments a Phylogenetic Tree was calculated. The length of each pair of branches represents the distance between sequence pairs, while the units at the bottom of the tree indicate the number of substitution events. The phylogenetic tree gives a first estimate of the relationship between homologous sequences. A cladogram presentation of the phylogenetic tree is shown in figures 11 and 12.

can be harvested three to five months after transplanting. The method yields single locus transformants at a rate of over 50 % (Chan et al. 1993; Hiei et al. 1994)

**Example 6: Expression of a cytokinin oxidase cDNA under the root-specific RCH1 promoter in transgenic Tobacco and Rice leads to increased root production.**

The AtCKX1 gene (located in the database with the accession number: AC002510, *Arabidopsis thaliana* chromosome II section 225 of 255 of the complete sequence, sequence from clone T32G6) can be cloned under the control of the RCH1 promoter of *Arabidopsis*, in a binary vector. Also the ZmCKX1 from maize (accession number AF044603, (Morris et al. 1999) can be operably linked to the RCH1 promoter a binary vector. These genes can be transfected into Tobacco or Rice plants, using agrobacterium-mediated gene transfer. The transgenic plants expressing the ATCKX1 or the ZmCKX1 specifically in roots are expected to show increased root production without negatively affecting shoot development and without inducing premature leaf senescence.

**Example 7. Minimal promoter sequence.**

The length of the RCH1 promoter region (SEQ ID NO 2), which confers to root specific expression patterns, can be reduced to a "minimal promoter" without altering the root specific expression patterns or other features of the promoter such as constitutive and strong activity. As an example, the length of the promoter region was reduced to a minimal promoter sequence ranging from the -1 base pair just before the start codon at position 3479-3481 of SEQ ID NO. 2 to -842 base pairs before said start codon. Said minimal promoter is represented in SEQ ID NO. 18. This minimal promoter was PCR amplified and translationally fused to the GUS reporter gene. With a GUS histochemical assay (example 2) the same expression pattern as the large promoter region was demonstrated (results not shown). The difference is that the sequence before the HindIII site at the beginning of the minimal promoter is not necessary for the root specific expression pattern.

**Example 8. RCH1 prom::GUS expression in embryo's.**

Siliques of Arabidopsis plants transformed with RCH1prom::GUS were opened at different stages of development and the embryo's were surgically taken out of the ovules and transferred to X-gluc staining solution. Embryo's were stained for 0.5h – 4h before pictures were taken using light microscopy, Fig 15 A and B are at 25X and 40X magnification respectively, stained for 0.5 h. Longer staining did not reveal expression outside the root promeristem region.

RCH1 is active very early: RCH1 expression was first observed in torpedo stage embryo's in the root promeristem and was maintained in this region during the later stages of development. This figure shows that as soon as a meristem is formed RCH1 is expressed and provides further evidence that RCH1 expression is correlated with the appearance and presence of a root meristem from the embryonic stage onwards.

**Example 9. Isolation of cDNA of other Arabidopsis RCH1 homologues****Table 4: % identity with RCH1**

	Database entry genomic DNA	Database entry protein	% sequence identity with RCH1 on aa level
<b>RCH1.2</b>	AB028621 + AP002037	BAB03091	59.8%
<b>RCH1.3</b>	AB011476	BAB09286	40.1%
<b>RCH1.4</b>	AL022223	CAA1826 or T05050	38.9%
<b>RCH1.5</b>	AC015446	AAG12526	42.1%

To discover the exact function of RCH1, the inventors want to circumvent the redundancy of the gene.

AB028621 (see figure 12, phylogenetic tree) is determined to be RCH1.2 protein and is also used by the inventors for further experiments. Double knock-outs of RCH1 and RCH1.2 were made and also other closely related sequences (AB011476.pro,

T05050.pro and AC015446.pro) are used in knock out experiments or in antisense expression.

### **Ectopic expression of RCH1**

The expression of RCH1 in normal cells correlates with the zone of rapidly dividing cells and is involved in the maintenance of meristem cells and dividing cells. Improved activity of RCH1 in a host cell results in the maintenance of cell division in that host cell and thus works against the differentiation of that cell. On the contrary diminished RCH1 activity in a host cell results in a diminished cell division in that host cell and thus in the loss of the amount of dividing cells in the meristem. Particular examples to influence these processes by altering the expression level of the RCH1 gene are described in the following paragraphs:

### ***Overexpression***

In order to obtain higher activity of the RCH1 protein in the host cell the coding region of the RCH1 gene is fused to a strong promoters that is inducible or constitutive and/or ubiquitous or tissue specific.

To influence the RCH1 activity in a particular set of host cells, the RCH1 coding region is cloned under control of other meristemic promoters such as the promoters of WUSCHEL, CLV1, CLV2, CLV3, STM. The obtained effect is the control of the expression of RCH1 in a particular area of the meristem or in a particular time period of the development of the plant. Also the promoters of other RCH genes (such as AB028621.pro, AB011476.pro, T05050.pro and AC015446.pro) are used for this purpose.

In a particular application of the present invention, the RCH1 gene and the RKN gene (= the rice homologue of RCH1 as described in WO/04761) are operationally linked to the root-specific promoter PYK10, a constitutive ubiquitin or GOS 2 promoter and Cdc2a promoter and expressed in Arabidopsis or in rice.

### ***Downregulation***

In order to obtain lower activity of the RCH1 protein in the host cell, the coding region of the RCH1 gene cloned in a sense and antisense orientation in order to form a hairpin that induces gene-silencing in the host cell.

5

### ***Co-expression***

Downregulation or upregulation of the RCH1 in combination with downregulation or upregulation of other genes e.g.:

#### With WUSCHEL:

10 In the shoot meristem a balance between the *WUSCHEL* and *CLAVATA* (*CLV*) genes has been implicated to play an essential role in regulation cell division and differentiation (Clark et al. 1993; Fletcher et al. 1999; Schoof et al. 2000). The whole meristem maintenance is controlled by these genes and they seem to have an opposite function keeping the balance between cell division and differentiation. Therefore, *WUSCHEL* or a  
15 root *WUSCHEL* homologue is a preferred partner to be co-expressed with RCH1.

In an application of the invention RCH1 is upregulated in combination with the downregulation of *WUSCHEL* in a root specific manner. The obtained effect of this co-expression is a better control on the meristem maintenance.

#### With other RCH1 family members

20 Possibly the function of RCH1 is redundant and other closely related genes can take over its function. Therefore the RCH1 gene ectopically expressed in combination with closely related genes in order to downregulate or upregulate the activity of the encoded protein. The RCH1 homologues used for co-expression are for example AB028621.pro, AB011476.pro, T05050.pro and AC015446 (see figure 12 and table 4). The obtained  
25 effect is a better effect as described for the ectopic expression of RCH1 alone.

#### With other Clavata homologues:

Clavata 1 is a receptor that is involved in a signal pathway probably through the binding of a ligand (e.g. Clavata 3). A particular useful application of the present invention is the co-expression of RCH1 with the root Clavata 3 homologue RCH3. The obtained effect is  
30 a better control on meristem maintenance.

**Example 10: Domain swapping**

The N-terminal LRR domain of RCH1 functions as an extracellular perception domain and a C-terminal kinase domain functions in an intracellular signal transduction pathway. Previously it was described by He et al. (Science 2000, 288(5475):2360-3; Perception of  
5 brassinosteroids by the extracellular domain of the receptor kinase BRI1) that such domains can be swapped with the domains of other proteins. Following this methodology in a particular application of the present invention, the domains of RCH1 are swapped with the domains of another protein. In one example the LRR domain of RCH1 is replaced by the LRR domain of BRI. The effect obtained is the activation of a different  
10 pathway, which is normally not activated by RCH1, because the kinase domain of RCH1 now responds to another external signal that normally is not recognized by RCH1.

It will be clear that the invention may be practiced otherwise than as particularly described in the forgoing description and examples. Numerous modifications and  
15 variations of the present invention are possible in light of the above teaching and, therefore, are within the scope of the appended claims.

## REFERENCES

- Alberts, B. et al. (1994) *Molecular Biology of the Cell*, Garland Publishing Inc..
- Altschul S.F. et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389-3402.
- 5 An G. et al. (1985) New cloning vehicles for transformation of higher plants. *EMBO J.* **4**, 277-284.
- Armstrong C.L. et al. (1990) Factors affecting PEG-mediated stable transformation of maize protoplasts. *Plant Cell Reports* **9**, 335-339.
- Banerjee A. et al. (1996) Omega amino acids in peptide design: incorporation into  
10 helices. *Biopolymers* **39**, 769-777.
- Baron M.H. & Baltimore D. (1982) Antibodies against the chemically synthesized genome-linked protein of poliovirus react with native virus-specific proteins. *Cell* **28**, 395-404.
- Bartel, P.L. & Fields, S. (1997) *The Yeast Two-Hybrid System*, Oxford University Press.
- 15 Baszczynski, C. L., Fallis, L., Bellemare, G., and Boivin, R. Brassica regulatory sequence for root-specific or root-abundant gene expression. WO9007001. 2000.
- Bechtold N. & Pelletier G. (1998) In planta *Agrobacterium*-mediated transformation of adult *Arabidopsis thaliana* plants by vacuum infiltration. *Methods Mol.Biol.* **82**, 259-266.
- 20 Bell M.H. et al. (1993) Tobacco plants transformed with *cdc25*, a mitotic inducer gene from fission yeast. *Plant Mol.Biol* **23**, 445-451.
- Benkirane N. et al. (1996) Exploration of requirements for peptidomimetic immune recognition. Antigenic and immunogenic properties of reduced peptide bond pseudopeptide analogues of a histone hexapeptide. *J.Biol Chem.* **271**, 33218-33224.
- 25 Berry A. & Brenner S.E. (1994) A prototype computer system for de novo protein design. *Biochem.Soc.Trans.* **22**, 1033-1036.
- Binarova P. et al. (1998) Treatment of *Vicia faba* root tip cells with specific inhibitors to cyclin-dependent kinases leads to abnormal spindle formation. *Plant J.* **16**, 697-707.
- Bögre L. et al. (1997) The *cdc2* kinase is differentially regulated in the cytoplasm and  
30 in the nucleus. *Plant Physiol.* **113**, 841-852.
- Bögre L. et al. (1999) A MAP kinase is activated late in plant mitosis and becomes localized to the plane of cell division. *Plant Cell* **11**, 101-113.



- Borg S. et al. (1997) Identification of new protein species among 33 different small GTP-binding proteins encoded by cDNAs from *Lotus japonicus*, and expression of corresponding mRNAs in developing root nodules. *Plant J.* **11**, 237-250.
- Calderini O. et al. (1998) A cell cycle regulated MAP kinase with a possible role in cytokinesis in tobacco cells. *J. Cell Sci.* **111** ( Pt 20), 3091-3100.
- Callis J. et al. (1995) Structure and evolution of genes encoding polyubiquitin and ubiquitin-like proteins in *Arabidopsis thaliana* ecotype Columbia. *Genetics* **139**, 921-939.
- Chan M.T. et al. (1993) Agrobacterium-mediated production of transgenic rice plants expressing a chimeric alpha-amylase promoter/beta-glucuronidase gene. *Plant Mol. Biol.* **22**, 491-506.
- Chaubet N. et al. (1996) Identification of cis-elements regulating the expression of an *Arabidopsis* histone H4 gene. *Plant J.* **10**, 425-435.
- Christou P., McCabe D.E., & Swain W.F. (1988) Stable transformation of soybean callus by DNA-coated gold particles. *Plant Physiol.* **87**, 671-674.
- Clark S.E., Running M.P., & Meyerowitz E.M. (1993) CLAVATA1, a regulator of meristem and flower development in *Arabidopsis*. *Development* **119**, 397-418.
- Clark S.E., Williams R.W., & Meyerowitz E.M. (1997) The CLAVATA1 gene encodes a putative receptor kinase that controls shoot and floral meristem size in *Arabidopsis*. *Cell* **89**, 575-585.
- Cohen-Fix O. & Koshland D. (1997) The metaphase-to-anaphase transition: avoiding a mid-life crisis. *Curr. Opin. Cell Biol.* **9**, 800-806.
- Colasanti J., Tyers M., & Sundaresan V. (1991) Isolation and characterization of cDNA clones encoding a functional p34cdc2 homologue from *Zea mays*. *Proc. Natl. Acad. Sci. U.S.A* **88**, 3377-3381.
- Crossway A. et al. (1986) Integration of foreign DNA following microinjection of tobacco mesophyll protoplasts. *Mol. Gen. Genet.* **202**, 179-185.
- Dale E.C. & Ow D.W. (1990) Intra- and intermolecular site-specific recombination in plant cells mediated by bacteriophage P1 recombinase. *Gene* **91**, 79-85.
- de Pater B.S. & Schilperoort R.A. (1992) Structure and expression of a root-specific rice gene. *Plant Mol. Biol.* **18**, 161-164.
- De Veylder L. et al. (1997) The *Arabidopsis* Cks1At protein binds the cyclin-dependent kinases Cdc2aAt and Cdc2bAt. *FEBS Lett.* **412**, 446-452.

- Dodds, J.H. (1985) Plant genetic engineering, Cambridge University Press.
- Doerner P. et al. (1996) Control of root growth and development by cyclin expression. *Nature* **380**, 520-523.
- Dolferus R. et al. (1994) Differential interactions of promoter elements in stress responses of the Arabidopsis Adh gene. *Plant Physiol* **105**, 1075-1087.
- Doerner B. et al. (1996) The synthesis of peptidomimetic combinatorial libraries through successive amide alkylations. *Bioorg.Med.Chem.* **4**, 709-715.
- Elledge S.J. (1996) Cell cycle checkpoints: preventing an identity crisis. *Science* **274**, 1664-1672.
- Ellis J.G. et al. (1987) Maize Adh-1 promoter sequences control anaerobic regulation: addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco. *EMBO J.* **6**, 11-16.
- Ely, S., Evans, I. J., and Schuch, W. W. Root-specific promotor. 288630. 2000. 8-10-1994. Ref Type: Patent
- Evans T. et al. (1983) Cyclin: a protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell* **33**, 389-396.
- Fantes P. (1989) Yeast cell cycle. *Curr.Opin.Cell Biol* **1**, 250-255.
- Fassina G. & Melli M. (1994) Identification of interactive sites of proteins and protein receptors by computer-assisted searches for complementary peptide sequences. *Immunomethods.* **5**, 114-120.
- Fedoroff N.V. & Smith D.L. (1993) A versatile system for detecting transposition in Arabidopsis. *Plant J.* **3**, 273-289.
- Feller H.S. & Jacobs T.W. (1990) Cell division in higher plants: a cdc2 gene, its 34-kDa product, and histone H1 kinase activity in pea. *Proc.Natl.Acad.Sci.U.S.A* **87**, 5397-5401.
- Fesquet D. et al. (1993) The MO15 gene encodes the catalytic subunit of a protein kinase that activates cdc2 and other cyclin-dependent kinases (CDKs) through phosphorylation of Thr161 and its homologues. *EMBO J.* **12**, 3111-3121.
- Fletcher J.C. et al. (1999) Signaling of cell fate decisions by CLAVATA3 in Arabidopsis shoot meristems. *Science* **283**, 1911-1914.
- Fowler T.J., Bernhardt C., & Tierney M.L. (1999) Characterization and expression of four proline-rich cell wall protein genes in Arabidopsis encoding two distinct subsets of multiple domain proteins. *Plant Physiol* **121**, 1081-1092.

- Francis D. & Halford N.G. (1995) The plant cell cycle. *Physiol.Plant.* **93**, 365-374.
- Francis,D., Dudits,D., & Inzé,D. (1998) Plant Cell Division, Portland Press London.
- Fromm M., Taylor L.P., & Walbot V. (1985) Expression of genes transferred into monocot and dicot plant cells by electroporation. *Proc.Natl.Acad.Sci.U.S.A* **82**, 5824-5828.
- Giritch A. et al. (1997) A root-specific iron-regulated gene of tomato encodes a lysyl-tRNA- synthetase-like protein. *Eur.J.Biochem.* **244**, 310-317.
- Goddemeier M.L., Wulff D., & Feix G. (1998) Root-specific expression of a Zea mays gene encoding a novel glycine- rich protein, zmGRP3. *Plant Mol.Biol.* **36**, 799-802.
- Gorthy P. & Pudukottah M.G. (1999) Production of silicon carbide from rice husks. *J.Am.Ceramic Soc.* **82**.
- Gubler F. & Jacobsen J.V. (1992) Gibberellin-responsive elements in the promoter of a barley high-pI alpha-amylase gene. *Plant Cell* **4**, 1435-1441.
- Hansen G. & Chilton M.D. (1996) "Agrolistic" transformation of plant cells: integration of T-strands generated in planta. *Proc.Natl.Acad.Sci.U.S.A* **93**, 14978-14983.
- Hansen G., Shillito R.D., & Chilton M.D. (1997) T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes. *Proc.Natl.Acad.Sci.U.S.A* **94**, 11726-11730.
- Hansen G., Shillito R.D., & Chilton M.D. (1997) T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes. *Proc.Natl.Acad.Sci.U.S.A* **94**, 11726-11730.
- Hanson B. et al. (1999) A simple method to enrich an Agrobacterium-transformed population for plants containing only T-DNA sequences. *Plant J.* **19**, 727-734.
- Harlow,E. & Lane,D. (1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press.
- Haseloff J. et al. (1997) Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic Arabidopsis plants brightly. *Proc.Natl.Acad.Sci.U.S.A* **94**, 2122-2127.
- Hayles J. & Nurse P. (1986) Cell cycle regulation in yeast. *J.Cell Sci.Suppl* **4**, 155-170.
- Helariutta Y. et al. (2000) The SHORT-ROOT gene controls radial patterning of the Arabidopsis root through radial signaling [In Process Citation]. *Cell* **101**, 555-567.
- Hemerly A.S. et al. (1993) cdc2a expression in Arabidopsis is linked with competence for cell division. *Plant Cell* **5**, 1711-1723.

- Herrera-Estrella L. et al. (1983) Chimeric genes as dominant selectable markers in plant cells. *EMBO J.* **2**, 987-995.
- Herrera-Estrella L. et al. (1983) Expression of chimaeric genes transferred into plant cells using a Ti-plasmid-derived vector. *Nature* **303**, 209-213.
- 5 Hiei Y. et al. (1994) Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**, 271-282.
- Hirt H. et al. (1991) Complementation of a yeast cell cycle mutant by an alfalfa cDNA encoding a protein kinase homologous to p34cdc2. *Proc.Natl.Acad.Sci.U.S.A* **88**,  
10 1636-1640.
- Hirt H. et al. (1993) cdc2MsB, a cognate cdc2 gene from alfalfa, complements the G1/S but not the G2/M transition of budding yeast cdc28 mutants. *Plant J.* **4**, 61-69.
- Hochstrasser M. (1998) There's the rub: a novel ubiquitin-like modification linked to cell cycle regulation. *Genes Dev.* **12**, 901-907.
- 15 Hoffman D.L. et al. (1995) Rapid protein structure classification using one-dimensional structure profiles on the bioSCAN parallel computer. *Comput.Appl.Biosci.* **11**, 675-679.
- Horvath B. et al. (1993) Lipo-oligosaccharides of *Rhizobium* induce infection-related early nodulin gene expression in pea root hairs. *Plant J.* **4**, 727-733.
- 20 Huntley R. et al. (1998) The maize retinoblastoma protein homologue ZmRb-1 is regulated during leaf development and displays conserved interactions with G1/S regulators and plant cyclin D (CycD) proteins. *Plant Mol.Biol.* **37**, 155-169.
- Hwang I. & Goodman H.M. (1995) An *Arabidopsis thaliana* root-specific kinase homolog is induced by dehydration, ABA, and NaCl. *Plant J.* **8**, 37-43.
- 25 Jaglo-Ottosen K.R. et al. (1998) *Arabidopsis* CBF1 overexpression induces COR genes and enhances freezing tolerance. *Science* **280**, 104-106.
- Jeong S., Trotochaud A.E., & Clark S.E. (1999) The *Arabidopsis* CLAVATA2 gene encodes a receptor-like protein required for the stability of the CLAVATA1 receptor-like kinase. *Plant Cell* **11**, 1925-1934.
- 30 Jinn T.L., Stone J.M., & Walker J.C. (2000) HAESA, an *Arabidopsis* leucine-rich repeat receptor kinase, controls floral organ abscission. *Genes Dev.* **14**, 108-117.

- John P.C., Sak F.J., & Lee M.G. (1989) A homolog of the cell cycle control protein p34cdc2 participates in the division cycle of *Chlamydomonas*, and a similar protein is detectable in higher plants and remote taxa. *Plant Cell* **1**, 1185-1193.
- John, P.C.L. (1981) *The cell cycle*, Cambridge University Press.
- 5    Joung J.K., Ramm E.I., & Pabo C.O. (2000) A bacterial two-hybrid selection system for studying protein-DNA and protein-protein interactions. *Proc.Natl.Acad.Sci.U.S.A* **97**, 7382-7387.
- Keller B. & Lamb C.J. (1989) Specific expression of a novel cell wall hydroxyproline-rich glycoprotein gene in lateral root initiation. *Genes Dev.* **3**, 1639-1646.
- 10   Kikuchi H. et al. (1999) Molecular characterization of a gene for alanine aminotransferase from rice (*Oryza sativa*). *Plant Mol.Biol.* **39**, 149-159.
- Krek W. (1998) Proteolysis and the G1-S transition: the SCF connection. *Curr.Opin.Genet.Dev.* **8**, 36-42.
- Krens F.A. et al. (1982) *In vitro* transformation of plant protoplasts with Ti-plasmid DNA.
- 15   *Nature* **296**, 72-74.
- Kumagai A. & Dunphy W.G. (1991) The cdc25 protein controls tyrosine dephosphorylation of the cdc2 protein in a cell-free system. *Cell* **64**, 903-914.
- Labbe J.C. et al. (1989) MPF from starfish oocytes at first meiotic metaphase is a heterodimer containing one molecule of cdc2 and one molecule of cyclin B. *EMBO J.* **8**, 3053-3058.
- 20   Lagarde D. et al. (1996) Tissue-specific expression of Arabidopsis AKT1 gene is consistent with a role in K<sup>+</sup> nutrition. *Plant J.* **9**, 195-203.
- Lake R.S. & Salzman N.P. (1972) Occurrence and properties of a chromatin-associated F1-histone phosphokinase in mitotic Chinese hamster cells. *Biochemistry* **11**, 4817-4826.
- 25   Langan T.A. (1978) Methods for the assessment of site-specific histone phosphorylation. *Methods Cell Biol* **19**, 127-142.
- Lauter F.R. (1996) Root-specific expression of the LeRse-1 gene in tomato is induced by exposure of the shoot to light. *Mol.Gen.Genet.* **252**, 751-754.
- 30   Lease K., Ingham E., & Walker J.C. (1998) Challenges in understanding RLK function. *Curr.Opin.Plant Biol.* **1**, 388-392.
- Lee M.G. & Nurse P. (1987) Complementation used to clone a human homologue of the fission yeast cell cycle control gene cdc2. *Nature* **327**, 31-35.

- Lerner R.A. (1982) Tapping the immunological repertoire to produce antibodies of predetermined specificity. *Nature* **299**, 593-596.
- Lerner R.A. et al. (1981) Chemically synthesized peptides predicted from the nucleotide sequence of the hepatitis B virus genome elicit antibodies reactive with the native envelope protein of Dane particles. *Proc.Natl.Acad.Sci.U.S.A* **78**, 3403-3407.
- Li J. & Chory J. (1997) A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* **90**, 929-938.
- Liddle, J.E. & Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, Wiley New York.
- Lisztwan J. et al. (1998) Association of human CUL-1 and ubiquitin-conjugating enzyme CDC34 with the F-box protein p45(SKP2): evidence for evolutionary conservation in the subunit composition of the CDC34-SCF pathway. *EMBO J.* **17**, 368-383.
- Llewellyn D.J. et al. (1987) Structure and expression of an alcohol dehydrogenase 1 gene from *Pisum sativum* (cv. "Greenfeast"). *J.Mol.Biol.* **195**, 115-123.
- Loffer F.D. (1994) The need to monitor intrauterine pressure-myth or necessity? [editorial; comment]. *J.Am.Assoc.Gynecol.Laparosc.* **2**, 1-2.
- Long J.A. & Barton M.K. (1998) The development of apical embryonic pattern in *Arabidopsis*. *Development* **125**, 3027-3035.
- Lundgren K. et al. (1991) mik1 and wee1 cooperate in the inhibitory tyrosine phosphorylation of cdc2. *Cell* **64**, 1111-1122.
- Magyar A. & Mihalik R. (1997) Characterisation of the swine swC1 antigen. *Acta Vet.Hung.* **45**, 17-31.
- Malamy J.E. & Benfey P.N. (1997) Analysis of SCARECROW expression using a rapid system for assessing transgene expression in *Arabidopsis* roots. *Plant J.* **12**, 957-963.
- Masuda S., Sakuta C., & Satoh S. (1999) cDNA cloning of a novel lectin-like xylem sap protein and its root- specific expression in cucumber. *Plant Cell Physiol* **40**, 1177-1181.
- Merrifield R.B. (1963) Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. *J.Amer.Chem.Soc.* **85**, 2149-2154.
- Mironov V. et al. (1999) Cyclin-dependent kinases and cell division in plants- the nexus. *Plant Cell* **11**, 509-522.

- Monge A. et al. (1995) Computer modeling of protein folding: conformational and energetic analysis of reduced and detailed protein models. *J.Mol.Biol* **247**, 995-1012.
- Morris R.O. et al. (1999) Isolation of a gene encoding a glycosylated cytokinin oxidase from maize. *Biochem.Biophys.Res.Comm.* **255**, 328-333.
- 5 Murakami T., Simonds W.F., & Spiegel A.M. (1992) Site-specific antibodies directed against G protein beta and gamma subunits: effects on alpha and beta gamma subunit interaction. *Biochemistry* **31**, 2905-2911.
- Murray A.W. & Kirschner M.W. (1989) Dominoes and clocks: the union of two views of the cell cycle. *Science* **246**, 614-621.
- 10 Nasmyth K. (1993) Control of the yeast cell cycle by the Cdc28 protein kinase. *Curr.Opin.Cell Biol* **5**, 166-179.
- Norbury C. & Nurse P. (1992) Animal cell cycles and their control. *Annu.Rev.Biochem.* **61**, 441-470.
- Nurse P. & Bissett Y. (1981) Gene required in G1 for commitment to cell cycle and in G2 for control of mitosis in fission yeast. *Nature* **292**, 558-560.
- 15 Nurse P. (1990) Universal control mechanism regulating onset of M-phase. *Nature* **344**, 503-508.
- Okada K. et al. (2000) Five geranylgeranyl diphosphate synthases expressed in different organs are localized into three subcellular compartments in arabidopsis [In Process Citation]. *Plant Physiol* **122**, 1045-1056.
- 20 Okumura N. et al. (1994) A dioxygenase gene (Ids2) expressed under iron deficiency conditions in the roots of *Hordeum vulgare*. *Plant Mol.Biol.* **25**, 705-719.
- Olson D.C., Oetiker J.H., & Yang S.F. (1995) Analysis of LE-ACS3, a 1-aminocyclopropane-1-carboxylic acid synthase gene expressed during flooding in the roots of tomato plants. *J.Biol.Chem.* **270**, 14056-14061.
- 25 Olszewski K.A., Kolinski A., & Skolnick J. (1996) Folding simulations and computer redesign of protein A three-helix bundle motifs. *Proteins* **25**, 286-299.
- Onouchi H. et al. (1991) Operation of an efficient site-specific recombination system of *Zygosaccharomyces rouxii* in tobacco cells. *Nucleic Acids Res.* **19**, 6373-6378.
- 30 Onouchi H. et al. (1995) Visualization of site-specific recombination catalyzed by a recombinase from *Zygosaccharomyces rouxii* in *Arabidopsis thaliana*. *Mol.Gen.Genet.* **247**, 653-660.

- Ormrod, J.C. & Francis, D. (1993) *Molecular and Cell Biology of the Plant Cell Cycle*, Kluwer Academic Publishers.
- Osborne, B.I., Wirtz, U., & Baker, B. (1995) A system for insertional mutagenesis and chromosomal rearrangement using the Ds transposon and Cre-lox. *Plant J.* **7**, 687-701.
- Ostresh, J.M. et al. (1996) Generation and use of nonsupport-bound peptide and peptidomimetic combinatorial libraries. *Methods Enzymol.* **267**, 220-234.
- Pabo, C.O. & Suchanek, E.G. (1986) Computer-aided model-building strategies for protein design. *Biochemistry* **25**, 5987-5991.
- Pawlowski, K. et al. (1994) Isolation of total, poly(A) and polysomal RNA from plant tissue. In: *Plant Molecular Biology Manual* (eds S.B. Gelvin & R.A. Schilperoort), pp. 1-13. Kluwer Academic Publisher, Dordrecht.
- Peralta, E.G., Hellmiss, R., & Ream, W. (1986) Overdrive, a T-DNA transmission enhancer on the A. tumefaciens tumour-inducing plasmid. *EMBO J.* **5**, 1137-1142.
- Prins, J. (1995) Cell cycle. Confirmational change [news; comment]. *Nature* **376**, 294-295.
- Poon, R.Y. et al. (1993) The cdc2-related protein p40MO15 is the catalytic subunit of a protein kinase that can activate p33cdk2 and p34cdc2. *EMBO J.* **12**, 3123-3132.
- Pysh, L.D. et al. (1999) The GRAS gene family in Arabidopsis: sequence characterization and basic expression analysis of the SCARECROW-LIKE genes. *Plant J.* **18**, 111-119.
- Reed, S.I., Hadwiger, J.A., & Lorincz, A.T. (1985) Protein kinase activity associated with the product of the yeast cell division cycle gene CDC28. *Proc. Natl. Acad. Sci. U.S.A.* **82**, 4055-4059.
- Renaudin, J.P. et al. (1996) Plant cyclins: a unified nomenclature for plant A-, B- and D-type cyclins based on sequence organization. *Plant Mol. Biol.* **32**, 1003-1018.
- Renouf, D.V. & Hounsell, E.F. (1995) Molecular modelling of glycoproteins by homology with non-glycosylated protein domains, computer simulated glycosylation and molecular dynamics. *Adv. Exp. Med. Biol.* **376**, 37-45.
- Roberts, C. S., Rajagopal, S., Smith, L. A., Nguyen, T. A., Yang, W., Nugroho, S., Ravi, K. S., Cao, M.-L., Vijayachandra, K., Patell, V., Harcourt, R. L., Dransfield, L., Desamero, N., Slamet, I., Keese, P., Kilian, A., and Jefferson, R. A. A comprehensive set of modular vectors for advanced manipulations and efficient transformation of



- plants by both *Agrobacterium* and direct DNA uptake methods. 2000.
- Rose R.B. et al. (1996) Three-dimensional structures of HIV-1 and SIV protease product complexes. *Biochemistry* **35**, 12933-12944.
- 5 Russell P. & Nurse P. (1986) cdc25+ functions as an inducer in the mitotic control of fission yeast. *Cell* **45**, 145-153.
- Russell P. & Nurse P. (1987) Negative regulation of mitosis by wee1+, a gene encoding a protein kinase homolog. *Cell* **49**, 559-567.
- Russell P. & Nurse P. (1987) The mitotic inducer nim1+ functions in a regulatory network of protein kinase homologs controlling the initiation of mitosis. *Cell* **49**, 569-576.
- 10 Rutenber E.E. et al. (1996) A new class of HIV-1 protease inhibitor: the crystallographic structure, inhibition and chemical synthesis of an aminimide peptide isostere. *Bioorg.Med.Chem.* **4**, 1545-1558.
- Sambrook, J., Fritsch, E.F., & Maniatis, T. (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press.
- 15 Schlappi M., Smith D., & Fedoroff N. (1993) TnpA trans-activates methylated maize Suppressor-mutator transposable elements in transgenic tobacco. *Genetics* **133**, 1009-1021.
- Schneider K. et al. (1998) The ROOT HAIRLESS 1 gene encodes a nuclear protein required for root hair initiation in Arabidopsis. *Genes Dev.* **12**, 2013-2021.
- 20 Schnittger A., Schellmann S., & Hulskamp M. (1999) Plant cells - young at heart? *Curr.Opin.Plant Biol.* **2**, 508-512.
- Schoof H. et al. (2000) The stem cell population of Arabidopsis shoot meristems in maintained by a regulatory loop between the CLAVATA and WUSCHEL genes. *Cell* **100**, 635-644.
- 25 Semler B.L. et al. (1982) A membrane-associated precursor to poliovirus VPg identified by immunoprecipitation with antibodies directed against a synthetic heptapeptide. *Cell* **28**, 405-412.
- Shioda T. et al. (2000) A green fluorescent protein-reporter mammalian two-hybrid system with extrachromosomal maintenance of a prey expression plasmid: Application to interaction screening. *Proc.Natl.Acad.Sci.U.S.A* **97**, 5220-5224.
- 30 Soni R. et al. (1995) A family of cyclin D homologs from plants differentially controlled by growth regulators and containing the conserved retinoblastoma protein interaction motif. *Plant Cell* **7**, 85-103.

- Sorrell D.A. et al. (1999) Distinct cyclin D genes show mitotic accumulation or constant levels of transcripts in tobacco bright yellow-2 cells. *Plant Physiol* **119**, 343-352.
- Sugita K. et al. (2000) Technical advance: A transformation vector for the production of marker- free transgenic plants containing a single copy transgene at high frequency [In Process Citation]. *Plant J.* **22**, 461-469.
- Sun Y. et al. (1999) Characterization of maize (*Zea mays* L.) Wee1 and its activity in developing endosperm. *Proc.Natl.Acad.Sci.U.S.A* **96**, 4180-4185.
- Suzuki H., Fowler T.J., & Tierney M.L. (1993) Deletion analysis and localization of SbPRP1, a soybean cell wall protein gene, in roots of transgenic tobacco and cowpea. *Plant Mol.Biol.* **21**, 109-119.
- Swenson K.I., Farrell K.M., & Ruderman J.V. (1986) The clam embryo protein cyclin A induces entry into M phase and the resumption of meiosis in *Xenopus* oocytes. *Cell* **47**, 861-870.
- Tamura R.N. et al. (1991) Cell type-specific integrin variants with alternative alpha chain cytoplasmic domains. *Proc.Natl.Acad.Sci.U.S.A* **88**, 10183-10187.
- Torii K.U. et al. (1996) The Arabidopsis ERECTA gene encodes a putative receptor protein kinase with extracellular leucine-rich repeats. *Plant Cell* **8**, 735-746.
- Trieu A.T. et al. (2000) Technical Advance: Transformation of *Medicago truncatula* via infiltration of seedlings or flowering plants with *Agrobacterium*. *Plant J.* **22**, 531-541.
- Uribe X. et al. (1998) Maize alpha-tubulin genes are expressed according to specific patterns of cell differentiation. *Plant Mol.Biol.* **37**, 1069-1078.
- van den Berg C. et al. (1995) Cell fate in the Arabidopsis root meristem determined by directional signalling. *Nature* **378**, 62-65.
- van den Berg C. et al. (1997) Short-range control of cell differentiation in the Arabidopsis root meristem. *Nature* **390**, 287-289.
- van der Zaal E.J. et al. (1991) Promoters of auxin-induced genes from tobacco can lead to auxin- inducible and root tip-specific expression. *Plant Mol.Biol.* **16**, 983-998.
- van Haaren M.J. et al. (1987) Overdrive is a T-region transfer enhancer which stimulates T-strand production in *Agrobacterium tumefaciens*. *Nucleic Acids Res.* **15**, 8983-8997.
- Van Sluys M.A., Tempe J., & Fedoroff N. (1987) Studies on the introduction and mobility of the maize Activator element in *Arabidopsis thaliana* and *Daucus carota*. *EMBO J.* **6**, 3881-3889.

- Vidmar J.J. et al. (2000) Cloning and characterization of a root specific high-affinity sulfate transporter from *Arabidopsis thaliana*(1) [In Process Citation]. *FEBS Lett.* **475**, 65-69.
- Wanapu C. & Shinmyo A. (1996) cis-regulatory elements of the peroxidase gene in *Arabidopsis thaliana* involved in root-specific expression and responsiveness to high-salt stress. *Ann.N.Y.Acad.Sci.* **782**, 107-114.
- Wang H., Fowke L.C., & Crosby W.L. (1997) A plant cyclin-dependent kinase inhibitor gene [letter]. *Nature* **386**, 451-452.
- Wang K. et al. (1987) Sequence context of the T-DNA border repeat element determines its relative activity during T-DNA transfer to plant cells. *Mol.Gen.Genet.* **210**, 338-346.
- Webb K.J. et al. (2000) *Mesorhizobium loti* increases root-specific expression of a calcium-binding protein homologue identified by promoter tagging in *Lotus japonicus* [In Process Citation]. *Mol.Plant Microbe Interact.* **13**, 606-616.
- Wilson A. et al. (1999) Evidence for the activation of a MAP kinase upon phosphate-induced cell cycle re-entry in tobacco cells. *Physiol.Plant.* **102**, 532-538.
- Winicov I. & Bastola D.R. (1999) Transgenic overexpression of the transcription factor *alfin1* enhances expression of the endogenous *MsPRP2* gene in alfalfa and improves salinity tolerance of the plants. *Plant Physiol* **120**, 473-480.
- Winicov I. (2000) *Alfin1* transcription factor overexpression enhances plant root growth under normal and saline conditions and improves salt tolerance in alfalfa. *Planta* **210**, 416-422.
- Wodak S.J. (1987) Computer-aided design in protein engineering. *Ann.N.Y.Acad.Sci.* **501**, 1-13.
- Woo H.H. & Hawes M.C. (1997) Cloning of genes whose expression is correlated with mitosis and localized in dividing cells in root caps of *Pisum sativum* L. *Plant Mol.Biol.* **35**, 1045-1051.
- Woulfe J. et al. (1994) Post-translational processing of the neurotensin/neuromedin N precursor in the central nervous system of the rat-II. Immunohistochemical localization of maturation products. *Neuroscience* **60**, 167-181.
- Xie Q. et al. (1996) Plant cells contain a novel member of the retinoblastoma family of growth regulatory proteins. *EMBO J.* **15**, 4900-4908.

- Xu Y. et al. (1995) Characterization of a rice gene family encoding root-specific proteins [published erratum appears in *Plant Mol Biol* 1995 Mar;27(5):1060]. *Plant Mol.Biol.* **27**, 237-248.
- Yamamoto E., Karakaya H.C., & Knap H.T. (2000) Molecular characterization of two  
5 soybean homologs of *Arabidopsis thaliana* CLAVATA1 from the wild type and  
fasciation mutant. *Biochim.Biophys.Acta* **1491**, 333-340.
- Yamamoto Y.T. et al. (1991) Characterization of cis-acting sequences regulating root-  
specific gene expression in tobacco. *Plant Cell* **3**, 371-382.
- Zeng Y. et al. (1998) Replication checkpoint requires phosphorylation of the  
10 phosphatase Cdc25 by Cds1 or Chk1. *Nature* **395**, 507-510.
- Zhang Y.L. et al. (1996) A superactive peptidomimetic analog of a farnesylated  
dodecapeptide yeast pheromone. *Biochem.Biophys.Res.Comm.* **224**, 327-331.

## CLAIMS

1. An isolated nucleic acid comprising a transcriptional regulatory root promoter or a functional fragment of such a promoter and/or comprising a nucleic acid sequence encoding a novel LRR receptor-like kinase (protein) or an immunologically active and/or functional fragment of such a protein (kinase) selected from the group consisting of:
- (a) a nucleic acid comprising the DNA sequence as given in any of SEQ ID NOs 1, 2, 3 or 18, or the complement thereof,
- (b) a nucleic acid comprising the RNA sequences corresponding to any of SEQ ID NOs 1, 2, 3 or 18, or the complement thereof,
- (c) a nucleic acid specifically hybridizing with the nucleotide sequence as defined in (a) or (b),
- (d) a nucleic acid encoding a protein comprising the amino acid sequence as given in SEQ ID NO 4, SEQ ID NO 5 or SEQ ID NO 19,
- (e) a nucleic acid which is degenerated, as a result of the genetic code, to a nucleotide sequence as defined in any of (a) to (d), and which sequence codes for a protein having the amino acid sequence as given in SEQ ID NO 4, SEQ ID NO 19,
- (f) a nucleic acid which is diverging, due to the differences in codon usage between the organisms, to a nucleotide sequence as defined in any of (a) to (d), and which sequence codes for a protein having the amino acid sequence as given in SEQ ID NO 4, SEQ ID NO 19,
- (g) a nucleic acid which is diverging, due to the differences between alleles, to a nucleotide sequence as defined in any of (a) to (d), and which sequence codes for a protein having the amino acid sequence as given in SEQ ID NO 4, SEQ ID NO 19,
- (h) a nucleic acid encoding a fragment of a protein encoded by a DNA sequence as given in SEQ ID NO 1 or SEQ ID NO 3 or encoding a fragment of a protein encoded by a nucleic acid as defined in any one of (a) to (g), wherein said fragment comprises the sequence as represented in SEQ ID NO 5, and,

- (k) a nucleic acid encoding a protein as defined in SEQ ID NO 4, SEQ ID NO 19 or a nucleic acid as defined in any one of (a) to (h), said nucleic acid interrupted by intervening DNA,  
provided that said nucleic acid is not one of the nucleic acids as deposited under the  
5 GenBank Accession numbers AB017061, AV538507, AV520681 or AQ966419.
2. An isolated nucleic acid comprising a transcriptional regulatory root promoter comprising the sequence as given in any of SEQ ID NOs 1, 2 or 18.
3. An isolated nucleic acid comprising a transcriptional regulatory root promoter consisting of the sequence as given in SEQ ID NO 2 or 18, or a functional part of  
10 said sequence which is able to regulate gene expression in a root specific manner, provided that said nucleic acid is not one of the nucleic acids as deposited under the GenBank Accession numbers AV538507 or AV520681.
4. An isolated nucleic acid comprising a transcriptional regulatory root specific promoter comprising:
- 15 (a) at least a functional part of the DNA sequence as given in any of SEQ ID NO 1, 2 or 18, and,  
(b) a second transcriptional regulatory sequence, optionally a regulatory promoter sequence not normally exhibiting root-specificity.
5. An isolated nucleic acid according to claim 4 wherein said second transcriptional regulatory promoter sequence is the ubiquitin promoter.
- 20 6. An isolated nucleic acid selected from the group consisting of:
- (a) a nucleic acid consisting of at least part of the DNA sequence as given in SEQ ID NO 1 or 3, or the complement thereof,
- (b) a nucleic acid encoding a protein as given in SEQ ID NO 4 or 19 or encoding a  
25 fragment of said protein, wherein said fragment comprises the sequence as represented in SEQ ID NO 5, and
- (c) a nucleic acid encoding a protein with an amino acid sequence which is at least 65 % identical to the protein as given in SEQ ID NO 4 or 9, wherein said amino acid sequence comprises the sequence as represented in SEQ ID NO 5,

or a homologue or a derivative of said protein, or an immunologically active and/or functional fragment thereof.

16. A polypeptide as defined in claim 15 being a root CLAVATA homologue or a functional homologue thereof
- 5 17. An isolated polypeptide comprising the sequence represented in SEQ ID NO 5 encodable by a nucleic acid of any of claims 1 to 7, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.
18. A method for producing a polypeptide according to any of claims 15 to 17 comprising culturing a host cell of claim 13 or 14 under conditions allowing the expression of the  
10 polypeptide and recovering the produced polypeptide from the culture.
19. An antibody specifically recognizing an LRR receptor-like kinase (protein) of any of claims 15 to 17 or recognizing immunologically active parts or specific epitopes thereof.
20. A method for the production of transgenic plants, plant cells or plant tissues  
15 comprising the introduction of a nucleic acid molecule according to any of claims 1 to 7 in an expressible format or a vector according to any of claims 10 to 12 in said plant, plant cell or plant tissue.
21. The method of claim 20 further comprising regenerating a plant from said plant cell.
22. A transgenic plant cell comprising a nucleic acid sequence of any of claims 1 to 7  
20 which is operably linked to regulatory elements allowing transcription and/or expression of said nucleic acid in plant cells or obtainable by a method of claim 20 or 21.
23. The transgenic plant cell of claim 22 wherein said nucleic acid of any of claims 1 to 7 is stably integrated into the genome of said plant cell.
- 25 24. A transgenic plant or plant tissue comprising plant cells of claim 22 or 23.
25. A harvestable part of a plant of claim 24.
26. The harvestable part of claim 25 which is selected from the group consisting of seeds, leaves, fruits, stem cultures, rhizomes, tubers and bulbs.
27. The progeny derived from any of the plants or plant parts of any of claims 24 to 26.

characterised in that said nucleic acid encodes a novel LRR receptor-like kinase protein or an immunologically active and/or functional fragment of such a protein, and further provided that said nucleic acid is not one of the nucleic acids as deposited under the GenBank Accession numbers AB017061 or AQ966419.

- 5 7. An isolated nucleic acid according to any of claims 1 to 6 which is DNA, cDNA, genomic DNA, synthetic DNA, or RNA wherein T is replaced by U.
8. A nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with a nucleic acid of any of claims 2 to 7.
9. A nucleic acid molecule of at least 15 nucleotides in length specifically amplifying a  
10 nucleic acid of any of claims 2 to 7.
10. A vector comprising a nucleic acid sequence according to any of claims 1 to 7.
11. A vector according to claim 10 which is an expression vector wherein said nucleic acid sequence encoding a novel LRR receptor-like kinase or an immunologically active and/or functional fragment thereof, is operably linked to one or more control  
15 sequences allowing the expression in prokaryotic and/or eukaryotic host cells.
12. A vector according to claim 10 wherein said vector comprises at least part of a nucleic acid according to any of claims 2 to 7 and wherein said transcriptional regulatory root promoter or functional parts thereof is operably linked to one or more genes of interest.
- 20 13. A host cell containing a nucleic acid molecule according to any of claim 1 to 7 or a vector according to any of claims 10 to 12.
14. The host cell according to claim 13, wherein said host cell is a bacterial, insect, fungal, plant or animal cell.
15. An isolated LRR receptor-like kinase comprising one of the polypeptides selected  
25 from the group consisting of:
  - (a) a polypeptide as given in SEQ ID NO 4 or SEQ ID NO 19,
  - (b) a polypeptide comprising the amino acid sequence as given in SEQ ID NO 5, and,
  - (c) a polypeptide encoded by a nucleic acid as given in SEQ ID NO 1 or 3,



28. A method for conferring root-specificity, root-meristem-specificity, root-vascular-tissue-specificity, root-endodermis-specificity, root-cortex-specificity, root-epidermis-specificity, root-quiescent-center-specificity, and/or abundant expression in root and/or early expression in root to other promoter sequences comprising the fusion at  
5 least a functional part of the DNA sequence as given in SEQ ID NO 1, 2 or 18 to a second transcriptional regulatory promoter sequence normally not exhibiting root-specificity.
29. A method for root-specific expression of a gene(s) of interest comprising operably linking of said gene(s) of interest to a transcriptional regulatory root-specific promoter  
10 as defined in any of claims 2 to 5.
30. A method for modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology comprising the modification of expression in the meristem of the main and the lateral roots, in the vascular tissue of the root or in the lateral root primordia of a gene(s) of interest operably linked to a  
15 transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
31. A method for phytoremediation comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
32. A method for environmental remediation comprising the expression of a gene(s) of  
20 interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
33. A method for enhancing plant growth and crop yield comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
- 25 34. A method for conferring enhanced resistance to pathogens which attack the belowground plant tissue comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
- 30 35. A method for enhancing freezing tolerance in plants comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.

36. A method for enhancing resistance of a plant to drought and/or high salt comprising the expression of a gene(s) of interest under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
37. A method for the production of light-sensitive proteins comprising the expression of a gene encoding said light-sensitive protein under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
38. A method for root specific gene silencing comprising the expression of an antisense nucleotide sequence to the gene of interest, a gene silencing construct or a ribozyme under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
39. A method for stimulating root meristem formation comprising the expression of a gene that influences the proliferation and/or differentiation of the root meristem under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
40. A method for stimulating root meristem formation comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase operably linked to a plant-operable promoter sequence.
41. A method stimulating root meristem formation comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase as defined in any of claims 15 to 17 operably linked to a plant-operable promoter sequence.
42. A method stimulating root meristem formation comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.
43. A method for stimulating root meristem formation comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.

- 44 A method for root-meristem maintenance comprising the expression of a gene that influences the proliferation and/or differentiation of the root meristem under the control of a transcriptional regulatory root-specific promoter as defined in any of claims 2 to 5.
- 5 45 A method for root-meristem maintenance comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase operably linked to a plant-operable promoter sequence.
- 46 A method for root-meristem maintenance comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like  
10 kinase as defined in any of claims 15 to 17 operably linked to a plant-operable promoter.
- 47 A method for root-meristem maintenance comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase under the control of transcriptional regulatory root promoter as defined in any  
15 of claims 2 to 5.
- 48 A method for root-meristem maintenance comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.
- 20 49 A method for enhancing root formation and/or root growth comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase operably linked to a plant-operable promoter sequence.
- 50 A method for enhancing root formation and/or root growth comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR  
25 receptor-like kinase as defined in any of claims 15 to 17 under the control of a plant operable promoter sequence.
- 51 A method for enhancing root formation and/or root growth comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase under the control of a transcriptional regulatory root promoter as  
30 defined in any of claims 2 to 5.

52. A method for enhancing root-formation and/or root growth comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.

5 53. A method for enhancing overall growth and yield comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase operably linked to a plant-operable promoter sequence.

54. A method for enhancing overall growth and yield comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR  
10 receptor-like kinase as defined in any of claims 15 to 17 operably linked to a plant operable promoter sequence.

55. A method for enhancing overall growth and yield comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase under the control of a transcriptional regulatory root promoter as  
15 defined in any of claims 2 to 5.

56. A method for enhancing overall growth and yield comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.

20 57. A method for modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase operably linked to a plant-operable promoter sequence.

58. A method for modifying cell fate and/or plant development and/or plant morphology  
25 and/or plant biochemistry and/or plant physiology comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase as defined in any of claims 15 to 17 operably linked to a plant-operable promoter sequence.

59. A method for modifying cell fate and/or plant development and/or plant morphology  
30 and/or plant biochemistry and/or plant physiology comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding

an LRR receptor-like kinase under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.

60. A method for modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology comprising the modification of  
5 expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.
61. A method to confer pathogen resistance to a plant comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR  
10 receptor-like kinase operably linked to a plant-operable promoter sequence.
62. A method to confer pathogen resistance to a plant comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase as defined in any of claims 15 to 17 operably linked to a plant-operable promoter sequence.
- 15 63. A method to confer pathogen resistance to a plant comprising the expression in particular cells, tissues or organs of a plant, of a nucleic acid encoding an LRR receptor-like kinase under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.
64. A method to confer pathogen resistance to a plant comprising the expression in  
20 particular cells, tissues or organs of a plant, of a nucleic acid encoding a polypeptide as defined in any of claims 15 to 17 under the control of a transcriptional regulatory root promoter as defined in any of claims 2 to 5.
65. A method for identifying and obtaining proteins interacting with a (LRR receptor-like kinase) polypeptide of any of claims 15 to 17 comprising a screening assay wherein  
25 a polypeptide of any of claims 15 to 17 is used or expressed.
66. A method according to claim 65 comprising a two-hybrid screening system wherein a nucleic acid encoding a polypeptide of any of claims 15 to 17 as a bait and a cDNA library as prey are expressed.

67. A method for modulating the interaction between an LRR receptor-like kinase (protein) as defined in any of claims 15 to 17 and interacting proteins obtainable by a method according to claim 65 or 66.
68. A method for identifying and obtaining compounds interacting with an LRR receptor-like kinase (protein) comprising the steps of:
- 5 (a) providing a two-hybrid screening system wherein a polypeptide of any of claims 15 to 17 and a protein interacting with said LRR receptor-like kinase (protein) or an interacting protein obtainable by a method of claim 65 or 66 are expressed,
- (b) interacting said compound with the complex formed by the expressed proteins as defined in a),
- 10 (c) detecting a second complex, wherein the presence of said second complex identifies a compound which specifically binds to one of said polypeptides or said second complex, and
- (d) identifying the compound.
69. A method for identifying compounds or mixtures of compounds which specifically bind to a polypeptide of any of claims 15 to 17, comprising:
- (a) combining a polypeptide of any of claims 15 to 17 with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,
- 20 (b) detecting complex formation, wherein the presence of a complex identifies a molecule which specifically binds said polypeptide.
70. Use of a molecule identifiable by means of a method of claim 68 or 69 as a plant growth regulator or herbicide.
71. A method for the production of a plant growth regulator or herbicide composition comprising the steps of the method of claim 68 or 69 and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture
- 25 72. Use of a nucleic acid molecule of any of claims 1 to 7, a vector of claim 10 to 12, a polypeptide of claim 15 to 17 or an antibody of claim 19 for modifying cell fate and/or


plant development and/or plant morphology and/or plant biochemistry and/or plant physiology.

73. Diagnostic composition comprising at least one nucleic acid molecule of any of claims 1 to 7, vector of claim 10 to 12, polypeptide of claim 15 to 17 or antibody of claim 19.

5

10

Figure 1





## Primers used in experiments described

SEQ ID No.	Primer	Sequence (5'-3')	Details
SEQ ID No. 6	LRR1	AAYAAYTTMASYGGTKMDWTHCC	Degenerate LRR primer
SEQ ID No. 7	LRR2	CARYTHAMCGGWGAAATHCC	Degenerate LRR primer
SEQ ID No. 8	LRR3	MAYYTSIAYGGGAHCTATHCC	Degenerate LRR primer
SEQ ID No. 9	KIN1	THCCWTTWGSCHGTANTCATA	Degenerate kinase primer
SEQ ID No. 10	KIN2	AYTCDGGDGGCHAHGTADCC	Degenerate kinase primer
SEQ ID No. 11	KIN3	MAYHCCRAARCTRTAVACATC	Degenerate kinase primer
SEQ ID No. 12	RCH1F	CGATCAGACACAAGAACAT	<i>RCH1</i> specific primer
SEQ ID No. 13	RCH1R	AGCAATGGTGTGGAAGAA	<i>RCH1</i> specific primer
SEQ ID No. 14	UBIF	TGCAGATCTTYGTGAAGAC	Compl. to 5' end of repeating ubiquitin unit
SEQ ID No. 15	UBIR	GACTCCTCTGGATGTTG	Compl. to 3' end of repeating ubiquitin unit
SEQ ID No. 16	M13F	TGTAAAACGACGGCCAGT	Compl. to pBS sequence
SEQ ID No. 17	PRCH1R	CACACAGGATCCAATCGGCATT GCAAAGACATA	Compl. to <i>RCH1</i> promoter, includes small coding sequence and <i>Bam</i> HI site

N=A+C+G+T; Y=C+T; M=A+C; S=C+G; K=T+G; W=A+T; R= A+G; D=A+T+G; H=A+T+C; V=A+C+G;

I=deoxyinosine

Figure 3




Figure 4




Figure 5-1 SEQ ID NO 1 : 7438 bp

[illegible]

GTTCTAGCGGTATTACGGCTGTTGGCGGTTATACGAGCTAAACAAATGATTCGAGATGATACAGATTACAGAGACTGGAGA  
 AAATCTATGGACATGGCAATTCACACCTTTTCAGAACTCAACTTCACAGTCGAACATGTACTCAAGTGTTCGGTAGAAG  
 GTAATGTTTATAGGAAAGGTTGCTCCGGGATAGTGTACAAAGCTGAAATGCCCTAACAGAGAAGTCATCGCGTGAAAAAG

Figure 5-2

CTCCTGGCCAGTGACAGTACCTAATCTGAATGAGAAAACTAAGTCATCAGGAGTTCGAGACTCATTCTCAGCTGAAGTAAA  
 AACACTTGGATCGATCAGACACANGAACATTGTACGCTCTCTGGGATGTTGTGGAAACAGAACACTAGACTCTTATGT  
 ATGATTATATGTCAAATGGAGTTTGGCAAGTTTCTCTCCGAAAGAGCCGCTATATGTACCTTAGGATGGAAATGAGG  
 TACAAGATTATACCTTGGGTGAGCTCAGGGTTTGGCTTACTTGCACCATGACTGTGTCTCTCCCATGTCTCATAGAGACT  
 CAAGGCAAAATAATTTCTGATTGGCCCTGATTTTGAACCTTACNTTGGAGATTTCCGACTCGCTAAGCTGTGTGATGATG  
 CCGACTTTGCTGCTTCTCCAAACCATTCGCTGGTTCCTATGGTTACATAGCTCCAGTACTCTCCCAAATTTCTGTCTTG  
 CTGCTAATTAGTTAGTTAAGTATGATACATCATATAGTTACTAGTAGTCTTAGGCTAGTAATGGTAGAAATTCAGGTTTATAGTTT  
 GOTTTCGTTTGGTTTTGGCATTTTGAAAGAATAGTATTTTGGTTCAGTTGATTTTCGATTTGGTTTAAATTTTAGAAAAAA  
 AATGGTAGAAGAAATAATTTTGAATTAATATATATCATCTACTGTATACGATTTAAAAAAAATTTGATTTGGTTTGGT  
 CTCGATATTTGCTTGTAGCTTGGTTCATATTTTACTATATATATGATTCATGACTTTTCATACTATATCATATTTTGGT  
 CGGTTTCGTTTGTATAGTCTGAAGTGAATCATATAGTTAAATATTTGACCAATCTATCATATATCATACAGCTACNAG  
 ATTTTCGATGTAATTTTGGGTGATGATTTTGGTTTTATATGTTTCATCCTTAGTAACATAACAAATATCATATTTGGGA  
 AGCATAACGTTATNTATCCNTAGGCTATACACTATAATTTAGCAGATACGATACCTCAATGAGATACAGAGAA  
 AAGCGACGTGTACAGCTATGGAGTCGTATGCTCGAGTACTAACGGGTAAAGCAACCAATCGATCCACGATACCAATG  
 GACTCCACATAGTGGACTGCGTCAAGAAAAFCAGAGACNTACAGTAACTCGACCAAGGACTACAGCAAGACCAAGATCA  
 GAGGTTGAAGAGATGATGCAAAACGCTAGGAGTCGCGCTTTTATGCAATTAATCCAAATACCCGAAAGACAGCGCTTACAATGAA  
 AGATGTGGCTGCTATGCTTATGTAGATATGTCAAGAGAGAGAGGAAATCGATGAAAGTTGATGTTTCTCGGGAAAGTTGTA  
 ACAATGGAAGAGAACGTGGCAAGAGATGATTCGACTTCATCGGTTATGCAACAAACCGCTAAGTATTTGAGAGATGATGAGC  
 ACGAGTTTCTCGGCTCTCTTPTGCTTTACTCTTCTCTCTCTGCTACTTCTAAGTGTAGACCAAACTCTTAAATAA

Figure 6 SEQ ID NO 2

CTCGAGTCTTCGATGTCCTTCTATTAATAGAGTATATGAGTCAAGCT  
AAGATATGAGAACCGAGGATTTCTTCGATGGAAGCTGAAATAT  
AGCCATCTGCTTTCTTAPAGAACACCACTTATCATCTGTATTAAT  
AGCAGCTAAGACACACTCTATGATTTCTTGATCCCTCTAA  
TGAGGGGAGGAGAACCGAGGCGCTTAAGCGCTCTCATATCTCAAC  
CAATGGCAAGGCTTAGCTCGTCCATCATATGATAGACATGTG  
CCCATAAAGATGAGAGCAGCCCAACCAAGCTCGCATCAT  
TGAGTCTTAGATTTCTGTACCGCGGACAGAGGCTGACAG  
AGTATGATCATCTCATAGCAACCAAGAAATCATGATATCTGTC  
AGCCAGCAAGTGTGTGAGAGGAATATCATGATTCOCAAATG  
ATGTCGATCTACAGTTTGCTGCTTCTGTGGTGTGTGGAAG  
ACATCTGTGATGAGCTGTGAGTGCAGAGTGTACCCATCTG  
ATTTGAATATAGAGACAGTGTGATTTCTGAGCTCTGATTT  
ATATTCAGATATGAGAACGAGGCTGTGTGGGSGTCCCTA  
GTCTAAGACCCAGAGCGCATCTCAGACAGCTGTGTGATTAAT  
CTGCTCCCGGATGTGAGCTGTGAGTGTGAGTGTGAGTGTG  
TGTGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG  
CTCTTAATTTCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT  
GAGCGAGATTAAGGTGTAGAGCAAGATCCAAATATCTG  
GCACCGCTTACAATCGATCTCAATAGCACCATCATGCTCT  
TTTGGAAAGAGCTCTCTTCATGAGCTGAGGTTCAAGAACT  
TTTGAAGATCTGGGATGTGAAGTGTGAGCTGTGAGTGTGAG  
GTGGAATCTCTGATGTAAATTTCTTATCATTTTGAAGAA  
AAATTTCTTGTGAGAGCGGAGGGGTGAGTATTAAGATGTGT  
CCGACGATCCCGGCTATCTGCTTCAAGTGTGAGCAACTCTCT  
TTTTCTTCTATCCGGGTGTGTAAACCGTCTCTCTGTCTCTA  
CGAATGTAGAGAGAGGAAAGAAATTAAGCAAGATGATGT  
TGCGTCTTAATCTAATCATTACCACTAGCAGAGCAATACAA  
TAGAGAAATTAAGTCTCAGAACTCAAGAAAGTCTTTTTTC  
TACTATATATATAGAAATCAACTCAATATGATCATTAAT  
CGGATAGCTTTCTTCTGCTTCTGTTCTCAAAATATFTT  
ATAAATATCTCAATTTCTTGAAGCTCATTGTGCTTAAT  
CAAGAGCAAAATCAAGAAAGAAAGCAATCATGCGTGGT  
TTAATATGAGATTTAGTACCTTGGGTTGGTATGATTTAT  
AGAAATTCAGTAAAGATGTAGAAATTTAGAGCAATCTCT  
ATGAGATCTCAGGTTGTGTGAGTCAATGAGTGTGAGTGT  
TTATGAGTGTGAGTGTGAGTGTGAGTGTGAGTGTGAGTGT  
CTTAATGTAGGAAATCAAGTCAAAACCAATGCTCTCT  
GTGAAATCTTGATATATATCAACCAATGCTCTCTCT  
TCTATGAGTATCTCTGGGCTCTTGCAACAGATATCTCA  
AAACAGATACCGGAGTGTAAAGAAAGAACAGACAGCAT  
TTATGCTCTTCTCTTCAAGCAAAATCTCTCCAAACCTTAT  
TTTGTGATTTGAGAGCCAAATCAAGAAAGAAATCTCTAT  
CTGATCTTAATGATAGAGAGAGTGTGTGAGTGTGAGTGT  
GTGAGTGTGATGTGATGTGATGTGATGTGATGTGATGT  
TCTATGAGTATGTGAGTGTGATGTGATGTGATGTGATGT  
AAAGTAAAGTTAATGACTGTATTAATTTCTGTAGTGT  
AGGTTTAGTACTTTTATATATATATATATCATCTCTCT  
GCCAAGACACTGATGATGTGAGTGTGAGTGTGAGTGT  
TTTGTGATTTGAGAGCCAAATCAAGAAAGAAATCTCTAT

Figure 7




Figure 8 SEQ ID NO 3 : 3960 bp

1 ATGCTCTTTCGAAATGCGGATTCACAGAAAAAGGCTTAA  
 2 TTCTTTCACTCTCTCGACCTCTGCATCAACCAATGAAGTTC  
 3 ACCGGTCAGTCTTCTCGCTGGGAATCCTCTGATTTCTGA  
 4 ACACAACTCGTTACGAGATCAATTCGCTCTCTGTTTCAGT  
 5 A TTTCCAGAACTGTTTTATCTCCAACTCAATCTCACCGGA  
 6 TATTCGATTTAAAGCTCCAAATAGTCTTTGTGGTGAATATCC  
 7 TTAACCTCCAAATGGCTCTACAGGAAGATCCCAACCGGAAC  
 8 TAACTACTTATCTCCGAGATCTCCCGTTGGAGCTCGGAATC  
 9 TCTCATGGGAAGATCCCGGAGGAGATCGGAACCTGTAG  
 10 CGTTCTTTACCTGTTTCTGTGGGTCAACTANGCAGCTCC  
 11 TCTTAAAGAGCTGGAAACTGCTCTGAACTTATCAATCTG  
 12 AACTAGGAAGCTTCAAACCTTAGGAAGATGCTCTTATG  
 13 TCTATGAAAGCTTAAACGCCATTGATCTCTCATGAACCT  
 14 AATCTCCAAGGCTTATGCTCTTAGCAGCAACAACATCACC  
 15 TTCAGTTCAGGATGTAGCGCAATCAGATTTCAGGTTTGTAT  
 16 TTTGGATTGCGAGATTAAGCTTAGAAGGCAATATCCCGAGC  
 17 AAGAAATGATCAAGCTCATTTACCTCGAGGTTTGTGTT  
 18 NATCTCCGGTGTATCCCACTTGGAGATCGGAACTGCAC  
 19 AAGAAATCCCTTAAGGATTTGGGATTTCTCTCAGAACCTTA  
 20 TTTTGTGAGATTAAGTAAATGTGAGCAACTCCAGATGCTT  
 21 TTTTATCATCTGTTAACAAGCTTCAAGTACTTGAATGCTCT  
 22 TACTCTCATTTCACTCAACCGCTCATCTTAAGTAAGAATT  
 23 CAATCTCAGCTTCTTGATCTCAGCTGATTAATACATCTCT  
 24 AATTTGCGTTTAAACTTGAGCTGGAATTCATTAGATGGCTT  
 25 TGGATATTTTCGCATTAACATGCTTTTCAGCGGACCTTTCTG  
 26 TATAACAGATTCTCAGGTTATCTTCCAGATAGTAAAGTG  
 27 AACTCTGTTCCAAGGTTTTCAGGTTCTTGCTTTTGAAGTAA  
 28 AACTCAGGATAGCAATTGGATTGCTAATCAGCGTGCAG  
 29 TAAACAAATGATTCAGATGATTAACGATTTCAGAGACTGA  
 30 TCAACTTCAAGCTCGAAGATGTACTCAAGTGTTTGGTAGA  
 31 AAAGCTGAAATGCCATAACAGGAATCAATCCGCGTGAATA  
 32 TAAGTCATCAGGAGTTTCGAGACTCATCTTCAGCTGAAGTA  
 33 TTTTGGGATGTTGTGGGAACAAGAACACTTAGACTCTTAT  
 34 TACCAAGAGAGCGGTGTATTTAGCTTGGATAGGATGGAGTGA  
 35 TTTCCACCACTATGTTGCTCCCAATGTTCTCATAGAGAC  
 36 TTTACATTTAGGATTTTCGGAATCGCTAAGCTTGTGTATGA  
 37 TATGTTTACATAGCTCCAGGTACTTCCCAATTTCTGTTT  
 38 TATGCTTTAGCTAGTAAATGTTAGAAATTACAGTTTATAGT  
 39 GGTTCGTACCTCTTGTGATTTGGTTTAAATTTTGAATAA  
 40 TCTACTGTATACGATTTTAAATAAAAAATTTGATTTGGTTTG  
 41 TATATATTATGATTCATGACTTCTCATACTAATCAATTTTG  
 42 TTAATAATTATGACCAATCTATCATATACTTACACGCTACA  
 43 ATATGTTTCATCTCTTAGTAACTTAACAACAAATATCAATTG  
 44 TAAATTATGTCAGATTAACGATACTCAATGAAGATACAGAG  
 45 TACTAAACGGGTAGCAACAATCGATTCCAAGTATCAGGA  
 46 ATACAAGTATCGAACCAAGACTACAGCAAGACAGAGT  
 47 TTTATGTTAATATCCAAATCCCGAGAGCAGGCTTACAAT  
 48 CAGAGGATTCGATGAAGTGTGATGTTGCTCGGGAAGTTG  
 49 TCGGTTATCCCAACAACGGCTAAGTATTTGAGAAATGATA  
 50 TTTCTTGTGACTTTTATATGTAGACCAAAATCTTAATAA  
 51 CTGTTTCTCATTTTTCATATACATTTGTTCTGTTTTTACG  
 52 TCAGCTTTTGATTTCTTGCGTTTCAGACTCGAATCCGCGC  
 53 OCCATGTCAATGGCCTTACATATTACTTGTTCCTCTCCAGAC  
 54 TAGCTTTTACCTTTTCCCTCTTACATATTCTTCTGTTTACTTC  
 55 GCTATATCTTCTTGAGATTGGAGATTTGTTCGGAGCTTATAG  
 56 TGTGAGTCTTAGGGAAGCTCAGAGATCTTCAAGAAGCTTTGT  
 57 TCGCGAGCTGCGTTTACCTCTCAGAGATCTTGAGATTTTCGA  
 58 ATCTCGACTCTTGTGAGCAATAGAGCCCGGAGAAATCTAG  
 59 GATCTCAAGGCTCTTAGGCTTAGCAGCAAGCAAAATTTCC  
 60 AGTCTTTATCTGTTCTATTTACATAGTCTCTCAGTGTGAGAT  
 61 TTTCTATAACGCAATGACTTGTTCGCGACGCTTCCATAAG  
 62 GCAGATTAATCTCCACGGACCTATCCCTTAGGAGATTTGGA  
 63 ATTTCTCAGGAACCATCCCTAAATCGTTGGTAACTTGTCT  
 64 GGGTCAATACCTTTCGATCTTAAGTAATCTGCACAAAGCTTG  
 65 TCCACAGAGATTTGGATTTGCTTAAGGAGCTCAACATCTTC  
 66 AGTTAGCTGGTTGTGAGAAATCTCAAGCTCTTGACTTGTCTC  
 67 CAGCTTCGTAACTTGACTAAGCTCTTGCTTATATCTAATG  
 68 TTTCTTTGTGAGATTAAGCTTTTCAATCAACAGATCAACC  
 69 GCTTCTCGACTTGTCTTAGAGCAATCTCTCCGGTCCAGT  
 70 AACTTGAACAACAACACTCTTCAAGTTATCTTCTCTCTTGT  
 71 TTTCAATGACTTTGACTGCGCAAGTAACTGTTACTTCTTGT  
 72 CTTCAATGGAGAAATCCCTTGTCTCTCGCTCACTGCGAC  
 73 GGAATATACCGGAGGAGCTTTTGTGCTCCCAAGATCTTAG  
 74 TATCCCGGAAGAGATTTCCCGGCTTAAACGATTAACGCTG  
 75 CGCTCTCCGCTCTAGAAATCTTGTCTTCTGAACTATCTG  
 76 TTTAGCAGCTGATAGGAGCAGAGTGGAGGAACAATG  
 77 CAGTTACAGATTAACTACACGCTGTTGTTGCACTCACAT  
 78 CGTTCTACCGCTATTAGGCTTTTGGCGTTTATACGAGC  
 79 GAAATCTATGGAATGCAATTCACACCTTTTCAAGAAAC  
 80 AGTAAATGTTATAGGGAAGGTTGCTCCGGATAGTGTAC  
 81 AGCTCTGGCCAGTGAACGTTACCTTATCTGAATGAGAAAC  
 82 AAACACTTTGGATCGATGACGACACAGAACATTTGACGCT  
 83 GTATGTTATATGTCAAATGGGAGATTGGGAAGTTTGTCTT  
 84 GTACAGATTAATCTGTTGTTGATGAGCTGAGGTTTGTGCTTA  
 85 ATCAAGGCAAAATATCTCTTATTTGGCCCTGTTTGTAGAC  
 86 TGGCGACTTTGCTCGTTCCTTCCAAACCACTTGTGTTCTC  
 87 TCGCTTAATATGATTAACTTACTTACTTCAATGTTACTAG  
 88 TTGTTTGTGTTTGTGGCAATTTGAAGAATATGATATTAT  
 89 AATAGTTAGAGAATATTTTGTATTTATATTTATATCA  
 90 GTCTCGATATTTTGTGCTTGTAGCTTGTGTTCAATTTTATAC  
 91 TTCGTTTCTGTTTGTGATTTAGCTTAGTGTGAAGTGAATCATATAG  
 92 AGATTTTTCGATTAATTTTGTGTAGAGCTTTTGTGTTTTT  
 93 GAACATAAAGTTTATATATTCCCAATAGGCTATAACACTA  
 94 AAGCGACGCTGTACAGCTTATGAGATCTGTATGCTCGAGG  
 95 TGTGACTCACATAGTGTGAGCTGGTCAAGAAATCAGAGAC  
 96 CAGAGTTGAAGGATGTAGTAAACAGCTTAGAGATCGCGCT  
 97 AAAGTTGTGCTGTCTATGCTTATGTAGATTAATGACAGAGA  
 98 TAAACATGGAAGAGACAGTGGCAAGATGATGTCCACTTCA  
 99 GCACGGATTTCTCTGCTGCTCTTTCGTTTACTCTTCTTC

Figure 9

```

1  MSLQMP1PRKKALAVSHFSITLSLFLAFFISSTGASTNEV  SALISGLHSSNSPPSPVFSQWNPDSQPCQWFIYTCSSPD
81  NKLVTETINVVSQALALFFPNISSFTSLQKLVISNTNLTG  AISSEIGDCSELIVIDLSSNSLVGRIPSSLGKLNKLQELC
161  LANSGLTGKIPPELGDCVSLKMLEIFDNYLSENPLELAK  ISTLESIRAGSNSELSKGIPEEIGNCRNKKVGLAAKTIS
241  GSLFVSLQLSLKLSLFFVYSTMLSGEIPKELGNSSELNL  FLTFDNLSTLPKELGKLQNEKMLLWONLHGFIPEEIG
321  FIKELAIATLSMNFYSOTIIPKSPCNLSNLQFALSSNNT  GSIPEILSDCTKLAVQFQIDANQISOLTPEIIGLKEINIF
401  LGWQNKLEGNIPDELACQNLQALDLSQNYLTGSLPAGLF  QLRNITKLILLISHAIGSVIPLFETQNTSLVLRLVNRRT
481  GEIPKGYGFLQNLSPIDLSSENLSGPPPLEISNCRQLQML  NLSNNTLQGYLPISLSGLTKKLQVLVDSSNDLTGKIPDSLG
561  HLISLARLITSKNSFNGEIPSSLGHCYNLQLLDLSGNWIS  GTIPEELFDYIQDIDIALNLWNLSLDGPPIPERISALNRLSV
641  LDIGHOMLSGDI2SALSGLENLVSLNISHNRPSGYLPDSKV  PRLIGAEMEGNNGLCCKGFRSCFVSNSSQITQRGVHSH
721  RLRIAGILLISVTAVLAVLAVLAVIRAKQMRDNDSESG  ENLWTWQFTPPQKLNFTVEHVLKCLVBSNVIKGKCSGIVY
801  KAEMPHREVIAVKKLWPTVTVNLNEKTKSSGV3RDSPSAEV  KTLGSTRHGNIVRFLGCCWNEKTRLLMYDMSNGSLGSLL
881  HERSGVCSLGMEVRY4YKTLGAAQSLAYL5RHDCVPP6TV7RD  IKANNILIGPDFEPIYIGDFGLAKLVDDGFARSSNTIAGS
961  YGYTAPEYGY8SKITEKSDVSYGVVVL9EVLTGKQPIDPT  IPDGLHIVDVVKIRDIQVIDQGLQARPESEVEEMQTLG
1041 VALLCINPIPEDRPTMKDVAAML10SRICQEREESMKVDGCS  GSCNNGRERGKDDSTSSVMQYAKYLRSSSTSPSASSLLY
1121 SSSSSATSINVRPNLK

```



Figure 10-1

```
CNV1      1 .....maamrlkktl.....lfhlhylyffspcfayt.dmevllnlkssm 38
RCH1     1 MSLQMPPIPRKKALTVSHFSITLSLFLAPFISSTASTNBVSVALISWLHSS 50
          39 igpkghglhdwihsspsdahcsfsgvsdddarvislnv..... 77
          51 NSPPPSVFGWNPDSDDPCWPYIITCSSPDNKLVTIEINVVSQQLALPFP 100
          78 ...sftp.....lfgtispeigmhtlvnltlaannftgelpiem 111
        101 NISSFTSLQKLVISNTNLTAIGASSSEIGDCSELIVIDLSSNSLVGRIPSSL 150
          115 ksltslkvltnisnngnltgtfpgeilkamvdlevldtynnfnngklppem 164
          151 GKLNKLNQELCLNSNG..LTGKIPEE.LGDCVSLKNLEIFPNLYSENLPLEL 198
          165 selkklylsfggnf.fsgaipesygdigsleylglnaglagsgkspafis 213
          199 GKISTLESIRAGGNSELSGKIPEEIGNCRNLKVLGLAATKISGSLPVSLG 248
          214 rlnklremlygyynsyttggvprefggltkileldmasctltgteiptslsn 263
          249 QLSKLQSLFV.YSTMLSGEIPKELGNCSLINLFYDNDLSGTLPKELGK 297
          264 lkhltlflhiinnltghippelsglvsksdlsinqltgeipqefinlg 313
          298 LQNLEKMLWQNHLHGPIPEEIGFMKSLNAIDLSMNYFSGTTPKFSGNLS 347
          314 nitlinlfrnnlyggiipe.....aigelpkl 339
          348 NLQELMLSNNITGTSIPSILSDCTKLVOQIDAANQISGLIPPEIGLLKEL 397
          340 evfevwennftlqlpanlgrngnlikldvdsnhlttglipkdlcrgeklem 389
          398 NIFLGWNKLEGNIPELAGCNQLQALDSLQNYLTGSLPAGLQLRNLTK 447
          390 lilsnfffgepipeelgckcksltkirivknllngtvpaglfnlpvtitie 439
          448 LLNISANAISGVIPLETGNCSTSLRLRLVNRRITGETPKGTGFIQLNSFD 497
          440 ltdnffgselpvptmsg.dvldqiylsnwwfsgeippaignfnplqtflfd 488
          498 LSENNLSGPVPLEINSCRQLQMNLNNVTLQCYPLSLSSSLTKLOVDVS 547
          489 rnrfrgnipreifehkhsrintsannttggiptsirsctslisvdlrnn 538
          548 SNDLTGKI PDSLGLHLISLNRLLSKNSFNGBIPSSLGHCNTNLQLDLSSN 597
          539 ringeipkginnvknl.g.tlnisgnltgtsiptgignmtslttdlisdnd 587
          598 NISGTIPEELFDIOBLDTALNLSNLSGGFIPERISA NRLSVLDISHNM 647
```



Figure 11




Figure 12




Figure 13 SEQ ID NO 18 : 842 bp

AAGCTTAAATGACAATTTAGTACCTTGGGTTGGTCATGATTTAGAGCGGAACAAATATACCAT  
ACATCAAACGAGGATATACAGAGAAAAATTCATGGAAGTATGGAATTTAGAGGACAATTTCTC  
TTCTGGGCTACAACGGACCGGCCATTGCTCATTTACCCAGAGGTATCGAGTTTGIGGACTTT  
TGATGCCGCTAGAGACTATTGGCATCGGATTGAAAAAATGTTTACTTCGTTGTTAACAATTTT  
CTGAATGCAATATTTTCCTTGTCATGAATATTTAAACTTGTTATTACTTTCTTTTAGCTTAGGTG  
TGGACAATTATGGAGTTTACTTCAAACGAGGAAGAATCTTAAACGCTCGGTTCAAGTCTCGAA  
AACAAACCAACTCACAATCCTGACTTAATTGAGGAAAACAATGCAAAACCACATGCATGCTT  
CCATATTTCTATCATAATCTTATAAGAAAAACACTACTAAGTGAAATGATTCTGTATATATAT  
AACCAATGCCTTTTGTTTGTGATATTTATGTATATATAACTATTGACTTTTGTCTCTATGGA  
TAGTGTCTCGGGCTCTTGGCAAACATATTTCAAAGAAAAGTTAATGACTGTAATTAATTAATC  
TGAAGCTAGAAACAGAACCCGAGGTAAAAGAAAAAGACAGAGCACATGAAGTTTAGTACTT  
TTATATATTTAATATATCAATCTTCTTATTGCTTATCTCTAAAGCAAAAACCTCCCTAAACCTT  
AAGCCAAAGGACTCAGATCGATGCAGAACCAAGAAGGCTTGTTTTGGATTTGAGAGCCAAAT  
GCAAAGAAAAAACTCTT

## NCBI Sequence Viewer

Figure 14

☒ NCBI
 ☒ Nucleotide banner

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search ☒ Nucleotide for ☐ Limits ☐ Index ☐ History ☐ Clipboard

☐ Display ☒ Default View as ☐ HTML ☐ Save ☐ Add to Clipboard ☐ Hide Brief

---

☒ 1: GI "3510337" [GenBank] Arabidopsis thaliana genom... PubMed, Protein, Related Sequences, 1

LOCUS AB017061 61712 bp DNA PLN 09-AUG-2000  
 DEFINITION Arabidopsis thaliana genomic DNA, chromosome 5, TAC clone:K19E20.  
 ACCESSION AB017061  
 VERSION AB017061.1 GI:3510337  
 KEYWORDS  
 SOURCE Arabidopsis thaliana (strain:Columbia) DNA, clone\_lib:Mitsui TAC clone:K19E20.  
 ORGANISM Arabidopsis thaliana  
 Eukaryota; Viridiplantae; Embryophyta; Tracheophyta; Spermatophyta; Magnoliophyta; eudicotyledons; core eudicots; Rosidae; eurosids II; Brassicales; Brassicaceae; Arabidopsis.  
 REFERENCE 1 (sites)  
 AUTHORS Kaneko,T., Katoh,T., Sato,S., Nakamura,Y., Asamizu,E., Kotani,H., Miyajima,N. and Tabata,S.  
 TITLE Structural analysis of Arabidopsis thaliana chromosome 5. IX. Sequence features of the regions of 1,011,550 bp covered by seventeen P1 and TAC clones  
 JOURNAL DNA Res. 6 (3), 183-195 (1999)  
 MEDLINE 29397451  
 REFERENCE 2 (bases 1 to 61712)  
 AUTHORS Nakamura,Y.  
 TITLE Direct Submission  
 JOURNAL Submitted (26-AUG-1998) to the DDBJ/EMBL/GenBank databases. Yasukazu Nakamura, Kazusa DNA Research Institute, Department of Plant Gene Research; 1532-3, Yana, Kisarazu, Chiba 292-0812, Japan (E-mail:ynakamu@kazusa.or.jp, Tel:81-438-52-3935, Fax:81-438-52-3934)  
 COMMENT Address for correspondence: kaos@kazusa.or.jp  
 For the latest information on annotation of this clone, please see [http://www.kazusa.or.jp/kaos/cgi-bin/agd\\_graph.cgi?c=K19E20](http://www.kazusa.or.jp/kaos/cgi-bin/agd_graph.cgi?c=K19E20)  
 Genes with similarity to proteins in the databases are described in 'product' or 'note' qualifiers. Genes that have no significant protein similarity are described as 'unknown protein'.  
 The software programs used to predict genes include: Grail (Informatics Group, Oak Ridge National Laboratory, <http://compbio.ornl.gov/Grail-1.3/>), GENSCAN (Chris Burge, MIT, <http://CCR-081.mit.edu/GENSCAN.html>), NetGene2 (S.M. Hebsgaard, et al., CBS, Technical University of Denmark, <http://www.cbs.dtu.dk/services/NetGene2/>) and SplicePredictor (Volker Brendel, Stanford University, <http://gremlini.zool.iastate.edu/cgi-bin/sp.cgi>).  
 Genes encoding tRNAs are predicted by tRNAscan-SE (Sean Eddy, Washington University School of Medicine, St. Louis, <http://genome.wustl.edu/eddy/tRNAscan-SE/>).  
 This sequence may not be the entire insert of this clone. It may be shorter because we remove overlaps between neighboring submissions. The 5' clone is K24G6 and the 3' clone is K20J1.  
 FEATURES Location/Qualifiers

## NCBI Sequence Viewer

source 1..61712  
/organism="Arabidopsis thaliana"  
/strain="Columbia"  
/db\_xref="taxon:3702"  
/chromosome="5"  
/clone="K19E20"  
/clone\_lib="Mitsui TAC"  
join(1745..1821,3560..4202,4287..4390,4470..4634,4781..4973)  
/note="gene\_id:K19E20.1"  
/codon\_start=1  
/evidence=not\_experimental  
/product="pectate lyase"  
/protein\_id="BAB10313.1"  
/db\_xref="GI:10177179"  
/translation="MTRSRSNETWHEHAVENPEEVAAMVDMISRNSTARRRLGYFSCSTGNPIDDCWRCDRRWQSRKKHLANCAIGFGRNAIGGRDGRYVVSDFPNDNDPNVNPFGPTLRHNAVQEEPLWVIFKRDVITLKEELIMNSFKTIDGRGVNVHLANGACITIQFVTNIIHGSHIHDCRPTGNAMVRSSPSHYGNRTMADGDGISIFGSSHIWIDHNSLNSCADGLIDAVMASTATISNNYFTHNEVMMLGHSIDTYTRDKVMQVTIAYNHFGEGLIQRMPCRHGYFHVNVNDYTHWEMYAIGGSASPTINSQGNRYLAPRNRFAKEVLSRVSTRQWRHWNWRSEGDFLNGAFFTRSGSGLGASARASSLAAKSSSLVGVTITYNAGALNCRGRRRC"  
CDS complement(9033..10973)  
/note="gene\_id:K19E20.2"  
/codon\_start=1  
/evidence=not\_experimental  
/product="selenium-binding protein-like"  
/protein\_id="BAB10314.1"  
/db\_xref="GI:10177180"  
/translation="MNPTQTFLFSPGGNSPASSPASHPSSLPFPQINNCRITRDLSQLIHAVFIKSGQMDRTLAAAEILRFCAATSDLHHRDLDYAHKIFNMQPQRNCFSWNTIIRGFSEDEFDKALIAITLFYEMMSDEFVEPNRFTFPSPVLKCAKATGKIQEGKQIHGLALKYGFSGDFEKLINLRMYVMCGFMKDARVLFFYKNTIEKDMVMVMDRRKRIDGEIVLWNVMLIDGYMRLGDKCAARMFLDKMRQSRVSVSWNTMISGYSLSNGFFKDAVEVFREMKKGDIRPNYVFLVSVLPAISRLGSLKEGLWHLHYAEDSGIRIDVLSALIDMYSKCGIIEKAITHVFERLPRENVITWSAMINGFAIHGQAGDAIDCFCKMRQAGVRPSDVAYINLLTACSHGGLVEEGRRYFVSQMVSVDGLEPRIEYHCGMVDLLGRSGLLDEAEFILNMPKIPDDVITWIKALLGACRMQGNVEMGKRVANILMDMVPHDSGAYVALSNMYASQGNWSEVSEMRLRMKEKDIRDPCGSLIDIDGVLHFEFVVEDDSHFKAKEINSMLEISDKRLAGRYRPTITQVLLNL EEDKENVLHYHSEKIATAFGLISTSPGKPIRIVKRLRICEDCHSSIKISKVYKRLITVRDRKRFHFQDGCSCCMDYW"  
11469..12086  
/note="gene\_id:K19E20.3  
unknown protein"  
/codon\_start=1  
/evidence=not\_experimental  
/protein\_id="BAB10315.1"  
/db\_xref="GI:10177181"  
/translation="MAASVEYFPPYSPPSHQHPLSPSPVPVPPPSHISPPPPPPSPPHHP PPHPSPPHQPSPSPYPHPPPPPSYPHPPHQPSPPPHVLPPPPPTPAEGHVIIVVVISLGSFLFFLAFLAALFCYLKRRKRSSTKAEITIEFDHLKVGQETIVQPGHGEQTRVVMLEEDIHLEVDIHKTEKLSRPSHLSSTGRHAIDISDPNHFEQKS"  
CDS complement(join(12718..13611,13749..14156))  
/note="gene\_id:K19E20.4"  
/codon\_start=1  
/evidence=not\_experimental  
/product="anthranilate N-benzoyltransferase"  
/protein\_id="BAB10316.1"  
/db\_xref="GI:10177182"  
/translation="MKINIRDSMTMVRPATETPITNLWNSNVDLVI PRFHTPSVYFYRPTGASNFFDPQVMKEALSKALVPFYPMAGRLKRDDDGRIEDCNGAGVLVQVADTPSVIIDFGDFAPTLNRLQLIPEVDSAGIHSFPLVLVQVTFKCGGASLGVMGHHADGFSGLHFINTWSDMARGLDLTPPFDRTLLRARPPAPAFHVEYQAPASMKIPLDPSKSGS

## NCBI Sequence Viewer

CDS

GPENTTVSIPKLTTRDQLVALKAKSKEDGNTVSYSSYEMLAGHVWRSVGCARGLPNDQGE  
TKLYIATDGRSRLRPQLPPGYFGNVIFTATPLAVAGDLLSKPTWYAAGQIHDFLVRMD  
DNYLRSALDYLEMQPDLASALVRGAHTYKCPNLGITSWVRLPIYDAFDGFWGRPIFMGPG  
GIPYEGLSFVLPSPTNDGSLSAIALQSEHMKLFEKFLPEI\*

join(15849..18529,18605..18747,19300..19808)

/note="gene\_id:K19E20.5"

/codon\_start=1

/evidence=not\_experimental

/product="receptor protein kinase-like protein"

/protein\_id="BAB10317.1"

/db\_xref="GI:10177183"

/translation="MSLQMPIPRKKALTVSHFSITLSLFLAFFISSTASTNEVSALI  
SWLHSSNSPPSPVPSGWNPSDSDPCQWPIYITCSSSDNKLVTETINVVSQALALPPFPNI  
SSFTSLQKLVISNTNLGAISSSEIGDCSELIVIDLSSNLVGEIPSSLGKLKLNQLCEL  
LNSNGLTGKIPPELGDVCSLKNLEIFDNYLSENPLELGKISTLESIRAGGNSELGSK  
IPEELGNCNRNLKVLGAATKISGSLPVSLGQLSKLQSLSVYSTMLSGEIPKELGNCSE  
LINLFLYNDGSLTGPKELGKLNLEKMLLMQNNLHGPIPEEIGFMKSLNAIDSMNY  
FSGTIPKSFGNLSNLQELMLSSNNITGSPILSNCTKLVQFQIDANQISGLIPPEIG  
LLKELNIPFLGWQNKIKGNIPDELACQNLQALDLSQNYLTGSLPAGLQFLRNLT/KLLL  
ISNAISGVIPLEIGNCTSLVRLRLVNRRTGEIPKIGLFLQNLSPFLDSENNLSGPPVP  
LEISNCRQLQMLNLSNNTLQGYLPLSLSLTKLQVLDVSSNDLTGKIPDSLGHILSLN  
RLILSKNSFNGEIPSSLGHCTNLQLDLSSNNISGTPPELFDIQDLIALNLWNLSL  
DGPPIPERTSALNRLSVLDISHNMLSGDLSALSGLENVLNLSHNKPSGYLPDSKVFPR  
QLTGAEEMGNGLCKSGFRSCFVSNSSQLTTQRGVHSHRLRIAIGLLISVTAVLAVLG  
VLAVITRAKQIMRDNDSSETGENLMTWQFTPPQKLNFTVEHVLKCLVEGNVIGKGCSCI  
VYKAEMPNREVIKLVPPVTPNLNEKTKSSGVDRDSFAEVKTLGSIHRKNIIVRFLG  
CCWNKLVLLMYDYMNSGLSGILLHERSGVCSLGEWVRD/KANNLLIGDPDFEPIYIGDF  
GLAKLVDDGDGFARSNTIAGSYGYIAPEYGYSMKITEKSDVSYGVGVLEVLTKGQIPI  
DPTIPDGLHIVLWVKKIRDIQVIDQGLQARPESEVEEMQTLGVALLCINPIPEDRFT  
MKDVAAMLSEICQEREESMKVDGCSGSCNNGRERKDDSTSSVMQQTAKYLRSSSTSF  
SASSLLYSSSSSATSNNVRPNLK"

CDS

join(22264..22578,23196..23354)

/note="gb|AAD49765.1

gene\_id:K19E20.6

similar to unknown protein"

/codon\_start=1

/evidence=not\_experimental

/protein\_id="BAB10318.1"

/db\_xref="GI:10177184"

/translation="MDPKSPFEIIDQLPKILGFVFDLSATRVSGHLT/TERCCQPFK  
VLHGGVSALIAELASLAGIASGFKRVAGIHLSTHHLRPAALGEIVFAESPVSVGK  
NIQVWEVRLWKAKTETPDNKMVMSTSRVTLFCGLPIPDHVKDAPDELKKVTSKL"  
join(25778..26072,26283..26377,26531..26650,26729..26818,  
26916..27031,27145..27244,27447..27518,27597..27644,  
27726..27803,27879..27998,28105..28210,28303..28466,  
28574..28708,28845..28975,29226..29304,29399..29446)

CDS

/note="contains similarity to 5'-nucleotidase

gene\_id:K19E20.8"

/codon\_start=1

/evidence=not\_experimental

/protein\_id="BAB10319.1"

/db\_xref="GI:10177185"

/translation="MNLCAQDVFDKLLWRDMSTKMKFPSPSAAELPDLRKSNNKRG  
LRMIKCRAGAGDGRVAVGDDVPSVTSSSYVEVDYLGQSTKGDNLKLDPLQSFSGDQ  
AYLEGPVEEVARTEQAQAAENLIRELGICQFPFSAQHSPRGIFCSRITLNLRSIAIGYD  
NTLMLHYNVMAWEGKAYDYCHENLKSMMGFPVDGLAPDPELVIRGLMDIKEGNLKYAD  
RFGYVKRAMHGTKMLSNKAVSEIYGRELVLDNRQSRWELNTFFSVSEALAYAQWVDR  
LDDGFIISADLGTLDYKGLYKAVAKALPRAHVREGQLKESMKPELFPVEPDPELPLGK  
DQKEAGKGLLLITNSDYHYTDMKMKHSPFNKFLPNDMDWRDLFDMVIVSARKPEFFQMS  
HPLYEVVTGEGLMRCPKFAETGGLYSGGSAQMIESLNVHGDEILYVGDHYTYDVS  
KVHLRWRITALICRELEEEYMALIGSRGHREELIELINQKEVVGDLFNQLRLALQRKS  
GRPAQTLAATNLDDQELTETMQKLLIVMQRLDDKIGLMLTETDGLFNKRYADIYTSRV  
SNFLNYTFPFMYFRSEQSLAHDSPDPAGIEN"

CDS

join(30274..30620,30707..32633)



## NCBI Sequence Viewer

```
/note="gene_id:K19E20.9"
/codon_start=1
/evidence=not_experimental
/product="mutator-like transposase-like protein"
/protein_id="BAB10320.1"
/db_xref="GI:10177186"
/translation="MATKKVIAICQSGGEFVTNKDGLAYSGGDAYAIDIDQDTCMSD
FKSELAENFGFSWENMTLKYFLPGNKTLITISKDKDFORMVSFSADAPNVEIFVLPE
EAEARNVSNMPSRSSRTTASEAVVWVVSAGDIVVGDDMIANDPQIEMGIMPEESSPL
PCNFVL/TDEQHIKAAQQWENAITGVDFRNSFTEFRDALHKYSIAHGFTYKYKKNDS
HRVSVKCAQGCQWRITASRLSTQLICIKKMMTRHTCERAVVWVVSAGDIVVGDDMIAND
KEKIKAFDPYKPKDIAEDIKREYGIQLNYSQAWRAKEIAREQLQGSYKKAYSQLPSPC
KKIRETNPGSIALFMTKEDDSFHRFLPSFYASISGFRQCRCPELLVLDTPADLNSKYQGV
MLVATAPDAEDGIFPVFAFVVDAAETEDMWVWFLEHLKALADPRITTFVADFQNGLKT
ALPLVFEEKQHHAAYCLRHLEAKLNMDLQAQFSHEARRFILNDFYAAAYATQPDAYYRS
IENIKSISPDAYTWVIESEPLHMANALPEGERYNHMSIFGLDFYSWVSEAHHELPITQ
MDEDLRAKLMSQSIYTHQVQSREWIVSTLPTPTNEEKQKEIELARSLQVSAPHNSLFEV
HGETINLVDINQCDQCDCKVWRITGLPCSHAVAVVEICGSPYEXCSRYFTSESRYLTY
AESINFPVNTMTMMILEEPPVEGVSVTTPPPTRLTPPGRPKSKQVEPLDMFKRQLQC
SNCKGLGHNNKTKCAVS"
CDS complement(join(33092..33310,33890..34039,34118..34270,
34394..34521,34741..34852,34989..35105,35298..35438))
/note="gene_id:K19E20.10"
/codon_start=1
/evidence=not_experimental
/product="mitochondrial carrier protein-like"
/protein_id="BAB10321.1"
/db_xref="GI:10177187"
/translation="MIPTTATVDDDEPGQIKRALIDASAGAISSGVSRVSSTPLVDIKI
RFQVQLEPTTSWGLVRGNLSGASKYTGWQATKDIFREEGFRGFWRGVNPALLMVMPPY
TSIQPTVLVHLKLSFASGSKTDEHIIHLSPYLSFVSGALAGCAATLGSYPFDLLRTLLA
SQGEPKVPTMRSAPVDIIIQSRGIRGLYNGLTPTLVEITVPYAGLQFGTYDMFKRMMW
WNRKYLSSKIPINVDITNLSSQLFCGLGAGTSAKLVCHEPLDVVKRFPQIEGLQRPR
YGARVERRAYRNMLDGLRQIMISEGMHGLYKGIVPSTVKAAPAGAVTTFVAYEFTSDWL
ESISW"
CDS 36788..37987
/note="gene_id:K19E20.11
pir||T09376
strong similarity to unknown protein"
/codon_start=1
/evidence=not_experimental
/protein_id="BAB10322.1"
/db_xref="GI:10177188"
/translation="MADSQRLSTASGVKDGQPPWKKKKLSNDTTSNPSPLEYDVILII
ARVSRSYNTNLVSVKSFRTSILTSPELYKTRTLLGKTENFLYVCLRPDEANPRWFTL
YRKPNQTL/DHHTTKKKKKKKKEKSSVNLAPISILNSHPVEWSAISVDHYLAIS
ADIEKAPYSNPVYLDCTRTHTWNEAPRMLAHTNSSEFEGIVYLPESFESPSLNCVVEY
NTMTQTRWKVPVPEPKRMFKLENLEKKIYKSFHLDGRAGKGLSLSYKSKHLTGCVLVLD
TVDSYRLSSCMENIAIFYRKGNFIWRGLDGKILVYGIKLEGLEPKFSRYSYSSQLAE
YGGKLVVLWDKYVPASGYKEKMIWCABISLEKRNKEIWNVVEWFDVAVLTVPKYSKIL
CATAATL"
CDS 38404..39522
/note="gene_id:K19E20.12
pir||T09376
strong similarity to unknown protein"
/codon_start=1
/evidence=not_experimental
/protein_id="BAB10323.1"
/db_xref="GI:10177189"
/translation="MSSPEKRRKKKQKSSPNPSLPEDLIVSILARVSRSYNTNLVSVS
KTRPSILTSPELYKTRTLLGKPEYFLVCLRPDEANPRWILYRKPNQTLTKKKKKK
EDSSVHLHAPILNPSPTVWSSSLVAVGSLYAITADIKDSCPSNVWYLDCTRHTWLD
SPRLRLAHINSDFNRTYFPGSSEKPSLNCVVEYNTNTQTWNVPVPPQKRKLKQGNME
GKIYIPPCQENHRKAVALNPVLTWEAVGLGTNLDNRGSFCMIGNIAYCYDPGSKFRWM
```

## NCBI Sequence Viewer

CDS

NCNTAEGDWSRLEGLEGLPKFARYSTVKLAAYGGKLVVLWDKYYVAASGYKEKMTWCAR  
ISLEKRNSEEIWGWVWEPDAVLTPVPSYKFVCAKSAATV"  
40591..41709  
/note="gene\_id:K19E20.13  
pir||T09371  
strong similarity to unknown protein"  
/codon\_start=1  
/evidence=not\_experimental  
/protein\_id="BAB10324.1"  
/db\_xref="GI:10177190"  
/translation="MSSPERKKRKSLESPSTFPNPSLPDDLIVSILARVSRLYYPI  
LSLVSKSSRTLVTSPELYKTRSFNRTESCLYVCLDFPPDPNPRWFTLYRKFNQNITE  
KTRNKSQFVLAPIPNHSHSSSIVAIGSNIYAIGGSIENAPSSKVSLDCRSHTWHEA  
PSMRMKRNPAAANVVDGKIYVAGGLEEFDSKSMVEVPDIKTQTWTFVLSPLAERFIYR  
SLVIEGEIYIFGDKVVTYKPKEDRWGGVGEHQSMDLGLFFHSYCVINDVLYCYRPGGI  
KWYESEKRSWRKLRLGLKGLSKLASSCVKLADYGGKMLLWDKYPICSGNKSHSISCAV  
VSLERCKNQGIRGKVEWFDMLTVPSSYNFVGALAAATL"  
join(42528..42695,42932..43054,43183..43265,43625..43734,  
43820..44100)

CDS

/note="gene\_id:K19E20.14  
unknown protein"  
/codon\_start=1  
/evidence=not\_experimental  
/protein\_id="BAB10325.1"  
/db\_xref="GI:10177191"  
/translation="MASNSEAGGSADYETLMSTSDVELLKKRAWRNEKAAPEILOVEGA  
LVDRAKEQIELVEETIEDYVENGIDPLVVSPLYOMDLDRAGQLLSYLRVRLKLIKEFM  
FHNLSSEEAERRLSEQKVFATRCADDLAKHFEETVLLKL PENYQSVLKGSLISEVDD  
MVARVVPVLVMTLVNLVQCRSRINWIPLLFEARTLSPSIYTKSEHSLIPPTYIYITQS  
FCNLSVGLPLLLTAEERAPRLWRWSAVISTSYVTRL"  
join(47405..47596,47798..47863,48017..48125,48257..48403,  
48519..48622,48732..48830,48921..48990,49159..49255,  
49331..49385,49512..49595,49681..50009,50129..50301,  
50385..50521,50676..50747)  
/note="gene\_id:K19E20.17"  
/codon\_start=1  
/evidence=not\_experimental  
/product="arginine methyltransferase-like protein"  
/protein\_id="BAB10326.1"  
/db\_xref="GI:10177192"  
/translation="MEIPSLNKQEQFTLASVTDLTSPSSSLSSSPVVATFSCVNEVKE  
LRFQPSKSDSGFDFDLSSTQLPKLGLQFTCVSDGSISSAKEKSSFSRSGVVIKFRDEK  
DSKEPCDSFEECKDDAVKQGSALPNGTVVSANKSKFDDKIEAASAKMYFHYVGGQLH  
QQNMLQDVYRTGTGYHAAMENRSDPSGRVVDVAGSGILSMFAALAGAKHVAVAEAS  
EMAEYARKLIIAGNPLLAERTIVIKGKIEDIELPEKADVLSIPEPMGTLLVNERMLETVV  
IARDRFLSPNGKMFPVGRHMAPFADEFLFVEMANKALFWQQQNYGVLDTPLYVAS  
HOGYFSQPVVDADFDRLLVAPSMEHVIDFTMMTVKLEPTLDSVPRCLKIEDFLSCY  
FVQSSSLFSEWHDPNSITVKKFSQEQEYFIEDIPLKFTASVCTRIHGLCAKDFVLDG  
STVORWFTTAPGAPTHMYQIRCVLSQPIHVMAGQEITGRHLHIAHSAQSTINITLS  
AKMWGPGANQGGLIQTSSECKLDLKEPYRNSQFQVYPTQEPFAQSQDIHHSDDLEEL  
ELLQQNANAQL"

CDS

complement(join(51155..51406,51952..52131,52447..52696,  
52806..52879,53145..53270,53408..53572,53658..53785,  
53900..54028,54123..54254,54347..54494,54771..54907,  
54997..55283,55380..55498,55559..55611,55791..55935,  
56245..56352,56467..56535,56693..56805,56925..56994,  
57106..57207,57282..57436,57516..57540))  
/note="gene\_id:K19E20.18"  
/codon\_start=1  
/evidence=not\_experimental  
/product="isoleucyl-tRNA synthetase"  
/protein\_id="BAB10327.1"  
/db\_xref="GI:10177193"  
/translation="MAAKKASGEKQEDGKYKQTVDLPKTGPGMRANSITREPQLKI,

## NCBI Sequence Viewer

```

WEENQVQPKRVSDNNNGGSFILHDGPPYANGDLHMGHALNKILKADIINRYKLLQNYKVQ
VYVPGWCHGLPIELKVLQSLDQVRKELTPLKLRAAKFAKATVKTQMESPKRFVGV
ADWNPNYLTLDPEYEAQIEVFGQMALKGYIYGRKKPVVHSPSSRTALAEAELEYPEG
HISKSIYAFKLVGGAKTSLDEFPINTYLAVWTTTPWTMPANAAVAVNAKLQYSVVE
VQSFNQOKLFVIVATDLVPALEAKWGVKLSISKTFPGSDLENCRYTHPIDNRDCPVVE
GGDYITTEESGTLVHTAPGHGQEDYATGLKYGLPLVSPVDDGKPTTEAGQFRGLSVL
GEGNTAVVSYLDENMISLMVEESYAHKYPYDWRTKKPTIFRATQWFAVSEGFRRTATMD
AINNVKWPVHQAVNRI SAMTSSRSRSDWCISRQRTWGVPIPAFYHVKTKEPLMNEETINH
VKSIISQKGSDAWNYMSVEDLLPEKYRDKAADYEKGTDTMDVWFDSGSSWAGVLSKRE
GLSFPADVYLEGTQDHRGWFQSSLLTSIATQGGAPYSAVITHGVLDEKGMKMSKSLG
NVVDPRLVIEGGKNSKDA PAYGADVMRLWVSSVDYTGDLVIGPQILRQMSDIYRKLRG
TLRYLLGNLHDMVRDNAVYPQDLPIIDQHALFQLENVVKNIQBCYENYQFFKIPQIQ
RFTIVDLSNFYFDIAKDRLYTGTSSTFRRSCQTVLSLTHLLSILRVIAPIVPHLAEDV
WQNLFPFEYRNEDGSAEAFVPELKWPTLNEQWLSFPAEDVLPWQKLELRLTEVNKVLLE
ARNEKMGSSLEAKVYLHTADAGMAAKLLEMSKAEADTLQRIFTSQVPEVLSMMEK
EMISSVOHTGEYVEGENKVVIGVSRAGGSKCERCWNYSQGVGSGSDHPTLCGRCSVI
VANPPEPAVAANSLA"
UTR complement(58740..59315)
/note="contains similarity to disease resistance response
protein
gene_id:K19E20.19"
/codon_start=1
/evidence=not_experimental
/protein_id="BAB10328.1"
/db_xref="GI:10177194"
/translation="MSKLILILTAQILLLTATALAGKNGEDFARTINRKLGLGKKKEK
LTHLRVYVHDIIVTRNPSIRIQGPVAKYSSSSYFGSITMIDNALTLDDVINSTVVGQ
AQGMVVGAAQKEIGLLMAMNLAFKTKYNGSITITLGRNTVMKSVREMPVVGSGMFR
FARGVVEARTKLFDMKTGDATVESNCYILHY"
CDS 59993..60445
/note="gene_id:K19E20.20
unknown protein"
/codon_start=1
/evidence=not_experimental
/protein_id="BAB10329.1"
/db_xref="GI:10177195"
/translation="MFRVNIQPDNNAQIEITTLQSQSKSRKLKIKKMSGSESVVVVG
VDSARHSYHALETALDLFFIFPKANPRFKLVVLHARPTATFFLGVAGPGTVDIIPHVE
EDLNKTADLVKKKCAEVCSAKSVSEISLSEVIEGDPFRINIMLEAVERHHA"
ppRNA join( complement(AB023028.1:1..96),
complement(61652..61712))
/note="gene_id:K19E20.21"
/product="U4 snRNA"
/evidence=not_experimental
PAGE COUNT 19394 a 11344 c 11242 g 19732 t
PROTIN
1 t g a t c a a c a t g a c t a a t c t t c c a c g a g a c t a g a a c c a c t a a c c c a c c t c c t c t a g a t a a a
61 g c t a a a c a a t t c t c t a c a a a a a t t t c c a t t a a t g t t t c t a a t t g t a a t t g a a a a c a t c
121 a a c t t a a a a t a c t t a a a a a a a t t t t t g t a t a a c g t g a t c a a t t t t a a c t t t t t t t t a
241 a t g a a t t c t a a t c c a a a t g t g t c t t a a t c t a c t g t c a t c a c a a c c c a c a a t a a t a g a a
261 a c a a a a c a t a t a g a c t t g g g c c a t t t t t c t g t t t a t g a t t t t c a c g a t t t t t c a c
301 c a a t a t a c t t a c a t c a c t a g t t c a g t t g a a t c t a c t a t a a a t c a t a a a g g t a c g t a a c t c a
361 a a t c a c a t a c a c g a t t c a a g t t c a t t t t t c a t t g g a a t t a g t a t t t t a g t a a a t c a c
421 c t a t t a t g a t a t t t g g a t t t t c t a g t t t t a a a a c g g g a g g a a a a a a t a t a a c g t c t t
481 t c a t a g c a c t a t a a a a t g t t a a t t t t t t t g a a t a a t a a a t a a t t c a a t a a c g a a g t
541 g t t t g t g a t t a c t t a g a g t t t a t t t g g a g a a a t a g t c a a a a a a a g g t t a a t a c c t t
601 t t t c c t g g t t c a t t t t t t t t t t t t t g c c a a c a t t t a a a t t c c a a t t g t g a g a t t t a
661 g c c c a t c a t t t a a t g t t g a g a a g a a t g a a a g t g a a a c t t a a c t a a t g a t a t a
721 c t t t g a t t a a a a c t g t g a c a a t a t g a t t a t a t a c a t a t t g t t a a a t t t a t t c a t g
781 t g a t a c a t a t a a a a t g t t t a g c a a g t a c a a a a c a t g g a a a a t g t a c t a c a a t
841 t a t a a c a t a c t a t t c t a t t a g c a t t t c c g c t t t a c t g a a a a c a a a a a c g a g t g a t c a
901 a t t t g t c t c t c t t t t t t t t t t t g g a c a c a a g t a c a a t t a t c t c t t a t t t c c t c c
961 t t t t t g t g t a a t t g t c t c t g t c g t a t c a c c a c a t t t t c a g c a t t g g a t g t t a c a a a t
1021 g c g a a c a a a c t a t a a t g t a a a a a c a t g t c a g c a g c a t c g g g c a t t t c a a a c a t a t g t a a

```

## NCBI Sequence Viewer

1081	ataatgaaga	aattatatga	tctattctaa	ataattaatt	aggccaaata	ttattctaa
1141	tgaaaaatgt	cacgtccaat	ttctcttata	taataactcc	ttcccttgaa	gccttctctt
1201	aactcttttt	atctctcttc	ttctctcttc	ttctttaaac	acaaaaaac	gacaaaaagc
1261	tctgtctcgg	acagagacaa	agaacacaga	cagagcaaga	acaaaggctg	ttacacaaat
1321	actgtgtctg	ttgtcttctg	cattactctc	atcaatgttc	ttcagtgccg	gcgattccac
1381	aaggtagtgc	aatgtttttg	agtcacaaat	atcgtgac	tggttctatg	acaaaagcat
1441	tgaagtcttc	actctccaaa	tctaagctaa	aacaagaaaa	ccctttttct	tgatttgaat
1501	tacttgatca	tgattacatg	gattttgtat	tatatagcta	aatatggagg	gagttctttt
1561	tattttattt	ttttatcata	gattatgtag	agatccatgt	gatcaaacac	taactgtgta
1621	ttagagttta	ttgaatgata	atgcaacatt	tcaatgtgta	ttagcagtg	ttaaagcgag
1681	tttgggtatg	tggttttata	atttgagttg	tgtttctggg	tttaactaat	gagttgggta
1741	attaatgact	ttcaggtcta	atgaaacatg	gcataaacat	gcagttgaga	atccagagga
1801	agtagctgcc	atggtggaca	tgtagtcttt	ctgtgtgctt	aaattttgtg	ttatatacta
1861	tatatgaata	ataatgattg	ttgtttgttc	tctaattaa	tatttttagca	tctaattgca
1921	ttggttagtt	agtaaaacaa	tttaaaaaat	acactaaact	agcatctact	gacgtttttg
1981	ttggataaat	tcattgtgac	ttcttttata	catttttaac	tactggagaa	aaagaaaaata
2041	aaaaacgaa	aactggtgaa	agttgttatt	atgtggagca	gggctccat	ggcattctgt
2101	tgaagtattt	tatacatctt	gggtcaaaag	tcaaaaggct	aaatccata	ggcgcaaaaca
2161	ccacacctcg	attccatttc	tttctactat	ttgttttatg	tttttggttt	atattaccctt
2221	aattctcata	atgataaaaa	aaatttctct	tatgcaaat	tttaactggt	tttaactaaa
2281	gtatgtcttt	gggtcccgcg	cactgacttt	ttcaacgaga	aaatcttttg	attatttaga
2341	atttttctgt	tgccgtagaa	aatgatgtga	aactttgttg	attgttctat	ttttacagat
2401	attatgtttc	cacattttat	aaatcacaaa	cgaaaaacct	aataggttat	aaaagtttga
2461	ttttcactta	tagaacaatt	cccatcgagg	ttttttttg	gtttcttcgt	ttttcttttt
2521	gtgttttcaa	gattgactac	attttgactc	cgtaagttaa	agaacttaat	gccaacagac
2581	tcgattttct	gtttgtcttc	gcttccatgt	tttaatttaa	attttttcgg	catgaacaaca
2641	atcaatcat	ttaaacaatt	tgcaaatcat	attgtaaaac	ataaaaaaaa	taaaaaata
2701	ataagttagt	aggataaagt	ctttatgctt	attcttactg	aggggtttca	gagttattgg
2761	tggtaccagc	atgaaacatg	atgcataaca	tggttctatt	agttgagtta	gattttggatg
2821	ggattctctc	tttaatcgaa	tgcaaccaaa	tcataatgat	gaataaattt	atttttgtaa
2881	aaaaacattaa	atattattgc	ttgcaaaaa	cttatggagt	tattgtagac	atttggtccaa
2941	aaacacgtgt	aatgtttctg	ttttttgttt	tgaagagaaa	catgtgtaaa	tgttccaaca
3001	ccataactttt	aaaacagtga	attgccatca	gcccacatga	ttccctctgt	gtagtgaatt
3061	gtctcttgaa	tgatttcaat	cgatattctc	gtcttatctc	gttattactt	ataactatac
3121	tagtactttta	aagatctggt	tggtcacaa	tataatgtta	tgactataatg	attaaacgaa
3181	tagtattttt	tggaattata	ttacccaaaa	acataataca	ctattataac	gcattttcta
3241	agaaagtttta	cgagtggtttt	attttttatt	ttgtgtttta	gaataatctg	gattcttgat
3301	tgaataggaa	tagttttcgt	aacaaagcaa	gatttttttc	aattttctta	caatgcgtga
3361	atagaagaca	tagaataaat	ccacccagac	aaactaattg	tcaaatagca	ccactataaa
3421	tggtgcccac	caattctctc	gatttcgggt	aatatagtaa	ccatgtttat	attttagtga
3481	taattctctc	taatttcaat	ttagcttagt	tgcattgtta	ttatattctg	gtttgtctaa
3541	accggttttc	atggttcagg	agcattcgaa	acagcacagc	gagaagaaga	ttaggatact
3601	tctcatgttc	aaccggaaat	ccaatcgatg	attttggcgt	ttgtgaccgg	agatggcaat
3661	cccggtcgaa	actcatagcc	aattgcgcaa	tgggtttcgg	tgccaatgca	atcgccggtta
3721	gagacggtcg	tactacgtga	gtgagtgcac	caaacgacga	taaccgggtt	aacctctaac
3781	cggtgaacat	acgtcacgca	gtaattcaag	agaacacatt	atggatctgc	tttaaacgtg
3841	acatggtcat	actaataaaa	gaagaactaa	tcatgaacag	tttcaaaaac	atcgatggtc
3901	gtggcgtcaa	cgtttcaata	gctaattggt	catgcataac	gattcagctc	gtgactaaac
3961	ttatcattca	tggtatacat	attcatgatt	gtagaccaac	tggttaacgt	atggttagaa
4021	gctctctctc	gcattatggt	tggtgagaact	tggtgatagg	agatgggatt	tcgattttcg
4081	ggctcgagta	tatttggatt	gatcataatt	cgcttctgaa	ttgttgctgat	gggtcttattg
4141	atgctgttat	tgcttctact	gctattacta	ttctccaaata	ttactctaca	caccacaatg
4201	aggttaagaat	ttgtctcaac	ttttatggtt	tttggttgct	tgaatcgga	gttatttttt
4261	tttttggtag	ttgtgtttta	tgacaggtta	tgacaggtta	acatagtgat	acctatacaa
4321	gggataaagt	ttatcaagtt	acaatagctt	ataatcattt	tggtggaagt	cttatttcaga
4381	gaatgccgag	gtatcatactt	tggtcttaact	tatgattctt	tggtcaaaact	tcgaaaagtct
4441	gacaaaaata	acatttttca	ttaatgacgg	tttaggcagt	gatattttca	tgtagttaac
4501	aatgatcata	cacattggga	aatgtatcgg	atcggtggca	gtgcaggttc	taccattaac
4561	agccaaggaa	atagatatct	cgccccaa	aaccggtttg	ctaaagagct	actttcaaga
4621	gttaggtcga	gttaggtcaaa	taaaagaacc	atttttctgt	acttattgac	tagtgtcaaa
4681	ctataaccct	tatacttata	gattacagtt	tcaacaaaa	ttctacagca	aaaacttattg
4741	ggatgtttca	ggtgacaaag	agagactatg	caggggcaatg	gcaatggagg	acttggaatt
4801	ggagatcaga	aggagattct	tttttaaacg	gagctttctt	cacacgtctt	ggatcaggct
4861	ttggagctag	ctacgctaga	gcttcaagtt	tagcagccaa	atcatcatct	cttgttgagg

## NCBI Sequence Viewer

4921 tcattacccta caatgcccgtt gctcttaatt gcagaggttg tgcggggtt taatagtac  
4981 taaagtctat cttcttttgg tttcttgcat tatacatggc aatagcacta ttgaatgtc  
5041 agggcttgatt taaaattgcc tcaacttgat tataaagttg taatgatttg ttccaccgc  
5101 aattattgat gtatctataa aagcaaacaa agttggaact cctccctaac tctctcatct  
5161 taatgttttc ccacacaagg ataaaagaaa catagaaatt tgcataaaaa aaaaaaata  
5221 cgagttctct cggaatgca atgtagttac aataaatatt gtgaagttta tcatctggc  
5281 atgtgactat ggttttttga tcagtttggt ccggttatat gcgcatgatg tgtttttggt  
5341 cttggtttag tttggtttaag agtttcaaaa ccaaaattaa cgaattttct tttccggcgt  
5401 aaattgggta tttttggctg aagtttggtt tattttccgg ttaatttgg tcaataaaac  
5461 tctagttaaca gttaaaagaa gtctaaaaat ttgggtgaaa cgggtctcag ttgggtataa  
5521 tctgggttatt aaaaacaaaa tacctaagga cgggttcgat cgaaccgaac caaaatttct  
5581 cgttttagcga tgcataaata aagttcttcc tagaggttca accgatagaa ctgaaaaatc  
5641 tgatttcata cgggttcggtt ttctcgggtg ggttatttat ccattggctg aaatcccgcga  
5701 taacgggtcg tcttatttat taggctaaact acaagtttgt cattcaaggt taaacctgcac  
5761 caacccgtag agtcocgctaa ccgggatatg attaagatgg gctgaccaag actaacccgc  
5821 tttgtctcag gataacatcc ttcataatga tggagaggtg tggggatgat tgcataaagg  
5881 aataagaagg gatgtgaagg tgggtgaacc acgtgagagg agcagaagac cacatgagag  
5941 catatgagag gaaataaaag aaaggacctt agtatcattg tcatgttatg tttttgatat  
6001 tgataagata gttttggcaa atggcttgct tctctttcat tacacacttt ttttcccaa  
6061 atgttctctt tatgctaate tctccttttt tattttttta atttgcaca tttagcgtgt  
6121 ttttttttac agtcgcagag actgcttcta ctgcgattca gacgacaagt aaaaacagtt tttaaagtgt  
6181 aatttaacgc cgttgagtcg ctgcgattca ctgatttctc tctcttttct gatcgtgcg  
6241 acgctgaaat tccagcgat tcccaataa catcatctct caggcttttt atttgcgcg  
6301 gcaactgcgg caectttatt gctaaacgaa catgctaatt atcttaacta aatttaattt  
6361 cagttcttgc agtcgcgact gagtaatttc ttaaaaactt tgaatagtag tttagataaa  
6421 gtatttctag tgagttaagc gttgataatc ttttttctt tgacctgaag tgcatacata  
6481 attaaaaatt aagtttagaa ttgataatc caacaatcca tctcaatttg ttttcgtggt  
6541 acccgctctt aatagcctc aagtgctaac aagtgctaac caattatatt aagcatttta taatatcat  
6601 ctgcggaagg acgtcagtag acagaaatct caagcgtaca gagggaatta ttgtacatgc  
6661 aagaataata atctttatgt cacaataaat atattttct ttgtagattt atatatgta  
6721 tccggttttag aatatttcat acaataatct taaacttttt tgcgcttttg aaaaacaaag  
6781 acatttcaaa agcatgtact ttaacttttt tcaatactat atgttactta ttcctgaggt  
6841 tgaagaagag tcaaatagag aattttgacg gattttggtt taaaatttca taacataaaa  
6901 aaacataat acttaataa gattttggtt taaaatttca taacataaaa taacttcttt  
6961 acttggagct ataccaaaaa ccacttgtat ttttagataa aagtagtaca caactttct  
7021 acttatgcga atacaacaca atatacaata atatatagtt tttagatgc aacactatca aaatttggta  
7081 cacccttcat attgtatgaa ttatgacata atatatagtt tttagatgc aacactatca aaatttggta  
7141 ttttttggct aacaataata gtctccattt atatatagtt tttagatgc aacactatca aaatttggta  
7201 aaaaatcaat agttttttacc catatagttt accacacaca ttaaggtttt tttctttgtt  
7261 agtatataag ataaaactat tgacaacaat tctacgcaat ttttttctta ttctcatgct  
7321 aatcagctaa ttaataagtc ttgcaccatg tctacgcaat ttttttctta ttctcatgct  
7381 ttggtagtttc tctctcaggt taatlaactt ttttttctta tctcttctta ttctcatgct  
7441 tctaatacgt caatgtacaa tagcttttga catttttgaa gaactttctt atattttttt  
7501 cataggaagc taatgaacta atgctatcat ttttttctta atacatattc atttttattt  
7561 aagaagcgt actaatttta atgataatc ttgtatgatt ttaattttaga tcatatatta  
7621 tgactacttta actcactatt ttgtgcagta ttaacgatta tttcttaaac tggaggcaaa  
7681 agagagtgtg gacaagaact agagccggga agctcaatgtg tgttgggcaa gtgtgaggaa  
7741 ctgtgtttcca acaaatgaa aggaattcggg agtttctatt agaagccacc aggctcttga  
7801 aagtataact gtaattgttt ctacaattgt agttccaccag gattttttta gaacgcaga  
7861 taatgtaaac tcaatgaaac aatagtttaa tttctatggt atataaaaat gaaaatttat  
7921 ctttaataata caaataatgt aatagtttaa aggtttgttc tttctgagat atatatact  
7981 attcaaaaac ttttatactt gccatattta gatgtagcca tttctctctt ctgagctctc  
8041 atttggctctt aaccttttt tgaataacgt tttctacact atcggtattc acagacacca  
8101 ctgcaataca atttaatcac aaaaaataa tttctacact atcggtattc acagacacca  
8161 tgaacaaatg atacaatcaa accccaaat tgggaagcata agacaagcta aagacataa  
8221 agatttgtac cettttgcgag atttgggagc agaaaaatag tagagacgaa gaattctctt  
8281 agacggagg aatagatagc aggaagatag gagaaagtct aaatttctgt ttccccaaa  
8341 tgtgtcaatt ttaagatat tttccgtatg agctagctca ttatgccgcg gaacaaaatt  
8401 ggtcttttga ccatatgac cgttttagtt cgggggtgac cacagaccgt tcaaatgata  
8461 ttggaactcg ctatgcccac ggttggatga ctttctgaca tttctactta ttaagcgat  
8521 acagcttgtga cagctttgac agttaccaat cacaataaat tttctactaa cgttataaaa  
8581 agtttttacc agagttttt ttgtacaaat tttttgtaca attttttcta acagctttgc  
8641 actgttccaa taaagctaac agctaaaaat cttgtacaat ttttggttac taatcagacc  
8701 cattatgta gggttctccc tactttgaga aaatcagcc caaagtctta actctggtaa

## NCBI Sequence Viewer

```
8501 taaatatttg aatgagtgtta ttctactgtg tatattttac acacataaaa cttggtttct
8511 tataaatttg aagttttgat atttttttga ataaaaactt tgagacataa ttcatcattt
8521 acccttttata ttacaattca tcaagatttt tattttatca cattttctaa ctcgtgaata
8531 aaagatttcaa ttgtatataa tgtggagtca gtaaagaaac aacagagctt ctgtatcac
8541 agggaaagag tcttcgtctt ttatgtttag ttccaccaat aatccataca cggagcatgag
8551 ccatcttgga acgttctgaaa acgttttcgg ttctgtaccg ttatttttct ttgttaact
8561 ttggagatga gttgatgga cgaatgacaa tcttcacaga tacgtaggtt cttacaatg
8571 cgtattggtt tgccagggtc cgtgcttacc aacccaacy cgtagcgat cttctcgtg
8581 ttgttaatga gtacattttc ttatctctct tctccaagt ttaacaaact cttgtgtggg
8591 attggtctat acccagctag tctcagtttg tctgagattt ctacgagcat tgagttgatt
8601 tctttgggctt ttgggtgtga atcatcttcc acaacaaact catgaagcac gccatcaatg
8611 tcatgataac tacaccagg gtcttttctt atatcttctt ccttcatctt cagcctcatc
8621 tctgaaccct ctgaccagtt tctttgagaa gcatacatgt ttgagagggc gacataagct
8631 ccatctgtat gaggaacct atccatcaga atattggcaa cgcctttgcc catctcgaca
8641 ttctctgca tcttgcaagc acccaaaagt gtttccata tcatcatcatt tggttttata
8651 ggcataatga gtataaactc ctacagctca tcaagaagac cgaacgacc caataaatca
8661 accatacacc cgtattgttc aatcctaggc tctaaacct ccaactgca catctgactg
8671 aaatctctcc ttctctctc taccacccg ccatgggtac aagcagtcac aagatataatg
8681 taggcaacat cactaggctg aactccagcc tgcctcattt tacaataaac gtcaatagat
8691 tcaactgctc ttactgtgat agcaaacccg ttaatcatcg cactcaagt atcactgttc
8701 tctctaggta acctctcaaa tacatgaatc gctttctcaa tgcateccaa ttctgaatc
10021 atatcaataa aagcagagcc aagcacatca tcaatctcaa tctcactat cttctgcatac
10031 aaatgcaacc attctcccaa ctctaaagac ccaagccgtg aaatagcagg aagaacatac
10041 accaagataa cataattcgg tctaatactc cttttctca tctacgaaa aacctcaata
10051 gcatctttga aaaaccgctt cagcgaataa cccgatctca tctgattcca cgaacaaacg
10061 ctcttttgac gcatttttatc aaacaaactc cgttcagctt tacaatcacc aagcttcata
10071 taaccatcaa tcatcaactt ccacaaactc attcaccat ctctctttct tctatcagtc
10081 ataccaccaa tctttttctc tataatgttc ttataaatac acacagcacc atctttcata
10091 caataccaca tccatacatc tctaaacaaga ttactcatca caaattcacc accaccaaat
10101 ccatacttca aagcacaacc atgaatctgc ttacctttct agttttccga atgtttccga
10111 caagccttaa gaacagaggg aaacgtaaac cgattcgggt caacaaactc atcatcattc
10121 atactgtaga taactgtgat tgcgatacaga gctttgtctt cgtcacctctc atgaaacattc
10131 ctgattatag tattccaaga gaagcaattt cgttgaggca ttgattaaa gatcttatgt
10141 gcgtagtcaa ggtccaggtg gtggagatcg gaggtagcac agaactggag aatcctctg
10151 gcggcgagag ttctctgcat ttgaccagat ttatgaataa tgaatggagt ttgagagagg
10161 tctcttatag ttgcagactt gctgtattga ggaagagtg agatgggat atctccggga
10171 gaagagcgcc gcgagtacc gccgttgagc aaaggtgtct ggtgtgggtt cattgagaat
10181 caacaaagat tctctagtggt ggctatgtaa gacgcataac aagatataat aaagattbtg
10191 gattcaaaccc gattcaaccg aaccgaataa accgatttat tctgatttag gcgctgaagc
10201 tgaaggagg agttcaaccg tcggcgaaat tattttggaa ttggtgtct ttgattcgggt
10211 ggttctggga acctctgaacc gttttgtgtt catctctaaa gttcattaga actctaacg
10221 aataaccgaa taactcttat catttgcgca gacaaagaga ttgatattt tatgagacaa
10231 gagggaagcca attctgggact aaactaatgc aaagcttagc atgagagtc cgaagcaga
10241 agcttagaag cacattccac aaatccgacc tttaactac tactatt tactacacata
10251 ggcttcaaca tagagacta tgcacttcac ctttgcaaca accattttaa tacataata
10261 ctcccaaat ggctgctctt gtggaatact ttccctatta ctacacacc tcccatcaac
10271 atctctctcc tctctcagtt ccaccctctc tctccctcca cctccctcat
10281 ttctctctcc tcatcaccca ccaccaccac atttttctc tctctcatag ccaccgccat
10291 caccatactc tctctctcat ccgcaccacc catcaccata tctcatccc ctacgcacc
10301 cgcctccacc teagtttctt ccacaccacc caaccaacc agcaccggg catcatgtta
10311 teattgtgtt tgttatctct cttggaagtt gttttttct tgcattctct gctgctgctc
10321 tctctctcta tctcaagaaa agtcttcaaa cttctacaaa gctgagatt atcgagttg
10331 atgaacacct caaagtccaa gaacccatag tgaaggacc ccatgtcgaa caaacgagag
10341 tggtaatgct taagaggac attcatttgg tagaggatc tcaacagact gaagacctta
10351 gcgctctcag tcaactcagt tcaactggaa gacatgccat tgatatttca ctgtcccaat
10361 atcattttac tgagcaaaag tcttaaatca aaaggtctca aatgtaaaaa atcatattta
10371 tteacacatc ctacttaata aaaacaattt aaatgtggtt ccaaatcttg caatcaaaaa
10381 cttgatattg attccaagat aaatgctgta catgtccgtc caatataac acttaatat
10391 ttactatgat ttctcaactt gcgaatgtaa gtaaccttat atattttaga tcatgtatc
10401 cgtttgctgg ctttaaatg agattcctag agattcctgt ttgttttctc atataacatta
10411 ttccatttaa ctcgagtttc agatgtttct attaaataag taatgaaagc agctctaat
10421 ctcccaaat tcaccaagat agaaagatag aaagatgaa acagagttat tgcattggaag
10431 ctgaaaatca tacagctgg ttgatataac caatgagaca caaacaatc acggcatgtg
10441 ttatagaaca acacttatca tctgtattaa cataagcaa aagttctcat acgacatagt
```

## NCBI Sequence Viewer

```

12601 ccaaaagtac agaccataag cacaccttat gatttgtgtt accctctaaa acacaggatg
12661 atgacaagaa gcaaatattgc ccaaatattac tgagggaaaa acgaggggctt aacgctctca
12721 tatctcaaac aaaaaacttct caaacagttt catgtgttca gatggagggg caatgggcaac
12781 ggataacgtg ccatcattag taggacttgg tagcacaana gacaaacctc cgtatggaat
12841 tccaccaggt ccataaaga taggacgacc ccaaccaaa g tctgcatac aaataggtaa
12901 tctaaccagg ctgtgtgatc ccaaatattgg gcacttgtag gtatgtgcac cgcggcaaa g
12961 ggctacagag gtccaggtga tctccaggta gtccaagagt gacctacgat agcttatcatc
13021 catacgaacc aagaaatcat gaatctgtcc tgcagcatac catgttggct tagataaaca
13081 atccccgtca acagccaatg gtgttgcagt gaatatcaca tccccaagt aaccaggagg
13141 cagctgcgga cgtagtctag accctccatc agttgcaatg tacagtttcg tctcttggtc
13201 gtttggaagc cctcgcgctc ttcccaactga tctccacaca tgcctgtcca acatctcgta
13261 tgagctgtag ctgacaggtg tcccatcctc cttggatttc gcttaagag caacaagctg
13321 gtctcgtgtt aatttgaata tagagacagt ggtattctca ggctctgatt tagacggatc
13381 aagaggatct ttcatacttg gtgcaggctg atattcaaca tgatgaaaag cgggctgttg
13441 cgggtcceta gctcggagag gtgttcgatac aatgaaaggt ggaatgggta ggtcaagacc
13501 acgagccata tccagccatg tgttgataaa atgaagacca gaaagacctc ctgcgcggtg
13561 atgttgatct caacccccac gtgaagctcc cccacattta aagaaggtca cctatctatc
13621 caaacaatat catatcataa ctaaaaacag accatacatt atccaagaat atctatcacg
13681 aactaaaaaa cagaaacaaca caagcattca ataaattaaa aaagggaaa g actttaatt
13741 aagctgacct gcagaaacgag aagcgggaaa gataatgacg aagcggagt atccattcog
13801 ggaataagct gacggagatt aagggtagga gcaaaatcac caaatatctc gataacagaa
13861 ggagtagctg ccacaacgaa gaccgtgtac aatgcaccca aaatcgacca ttggaagaa
13921 tcatctcttc tcttcaagcg accagccata gggtaaaaag ggacaaagggc ttggaagaa
13981 gttctcttca ttgacctgagg gtcaaaagaa ttggaagcgc cgggtgggtct ctgagagtag
14041 acactagggg tatggaatct ggggatgaca aggtcgacgt tggagtcca aagattagtg
14101 atgtgtgtct cgggtggcagg ccggaccagt ggtgaatctc ttgatgtaat ttcatattt
14161 ttaagaaaga tcaaacgaac aaacaatttt tgttttgagt tgactctgag aaattgtgtg
14221 aagaacggag tcgggtagat ataagtgggt gtgaacaaa ttgaaagaggt gtaactcctt
14281 caactaccgc cccacgaccc cgcctactcc atctctgcac caactctctc agaaggttta
14341 ctattgttgc atctctacac tctgtctcac tttctctatc ggggttggtgta aaccggtctc
14401 ctctgtctta aggcaccaat aataactaat ggtcttaata gtataaagtt cgaatgaaga
14461 agaagaaaaa aaaatttgaca agttgatgtt tatagctacc gtacaagtta gtcaactgaa
14521 tgccaaaagt tggcttaact taactctgta ccagctagca gacataaca gcaaccacca
14581 ataacatcta accgcgtaaa ctacagtggt aagagaatta cgtcagaact caagaaggtc
14641 tttttttcag acaccaccaa ataaattatg ttgaaattat atccgaacta tactatatta
14701 tatgaaaaac tacctcaatt agtcataaat ttccatttc ttcatatttt attactaat
14761 ggtgtgtgtt gggatagct tttcttgc tctgttcta aaattatttc ttgacctta
14821 ttgtgatgtc caacttgggt tgaggatcta ataaaatttc aaattctbtt aagcatctat
14881 tggcttaagt atccaaacct ttgtctctc cactctctaa gaaaataaag caaaagcaaa
14941 actaaaagaa aagcaactag taccgtgggt tggctcaatg taaccctatc taccaaatca
15001 caatcaagac ttaaatgaca atttagtact ttgggttgg tcatgattag agcggaacaa
15061 atataccata catcaaacga ggtatatacag agaaaattca tgggaagtat gaatttagag
15121 gcaaatctct acacttgggt acaacggacc gcccatttcg ctattatcc cagaggatc
15181 gaggtttgtt gacttttgat ccgctagaga ctatgggcat cggattgaaa aaattgttta
15241 cttcgttgtt acaaattttc tgaatgcaat atttctctg tcatgaaatt ttaactattt
15301 tattactttc ttttagctta ggtgtggaca atattggagt ttaactctga cgaggaaaga
15361 tcttaaacgc tccggttcagg tctcgaaaac aacaacactc acaactcaga ctttaatgag
15421 gaaaacaaat gaaaaccaca tgcattgtct catatttcta tctaattctt atagaaaaaa
15481 acactactaa ctgaatgat tctgtatata tataaccaat gctttttgtt ttgtgatatt
15541 ttatgtatat ataactattg acttttgtca tctatggata gttgtctcgg ctcttggcaa
15601 acatatttca aagaaaagtt aatgactgta ataatatcat ctgaagctag aaacagaaac
15661 ccgaggtaaa agaaaaagac aggtttagta cttttatata ttttaatat taacttggca
15721 cattctttat tttgtcttat ctctaaagca aaaacttccc taaacctcaa gccaaaggac
15781 tcagatcgat gcagaaacca gaaggttgtt ttgggatttg agagcaaat agcttcaact gttctcat
15841 aaactcttat atcttttgcaa atgcgcatct caagaaaaaa agcctcaact gttctcat
15901 ttctcatatt ctgtctctcg tttttgctt ttttgactc ctgcacctc gttcaacca
15961 atgaagtctc agctttgatt tcttggcttc acagctcgaa cctgcgcaca ccgtcagctg
16021 tctcggctg gatactctct gattctgacc ctctcaatg gccttacatt acttgttctt
16081 cctcagacaa caaactctgt acagagatca atgtcgtctc tcttcagtta gctttacctt
16141 tccctctcaa cattctctcg tttacttca ttcgaaacaf gtttatctc aacactaatc
16201 tcaaccggag tatattctct gagatcggag attgttcgga gcttatagtt atcgatttaa
16261 gctcaaatag tcttgttgtt gaaataacct cgagcttagg gaagctcaag aatcttcaag
16321 aactttgttt aaactccaat ggcctcacag gaaagatccc accggaactc ggcgactcgg
16381 tttagctcaa gaactcttag attttcgata actacttatc cgagaactc cgttggaggc

```

## NCBI Sequence Viewer

```
16441 tcggaaagat ctgcactctt gagagcataa gagccggagg aaactcagag ctttcaggga
16501 agatcccgga ggagatcgga aactgtagga atctcaaggt cttagggtta gcagcaacga
16561 aaatatccgg ttcttttaact gtttcgttgg gtcaactaag caagctccag tctttatctg
16621 tctattctac aatgctctca ggtgagatcc cttaaagagct tggaaactgc tctggaactta
16681 tcaactctgt ttctatcacg aatgactctg ccggcacgct tccaaaagaa ctgagaaagc
16741 ttcaaaactc agagaagatg ctctctatgg agaataatct ccacggacct atctcctgag
16801 atgttggatt catgaaaagc ttaaaaccca ttgattctct tatgaactat ttctcaggaa
16861 ccactccctaa atcgttcgtg aacttgtcga atctccaaga gcttatgctt agcagcaaca
16921 acatcacccg gtccaatact tegtattctaa gtaactgcac aaagcttggc cagtccaga
16981 ttgacgcccc tcagatttca ggtttgattc caccagagat tggattgctt aaggagctca
17041 acatctctct gggatggcag aataagctag aagggaatat tccggacag tatgctggtt
17101 gtcagaatct tcaagctctt gacttgtcac agaattatct caccggatca ttactgcag
17161 gtttctttca gcttcgtaat ctgactaagc tcttgcttat atctaagtc atctccgggt
17221 ttatcccaat ggagatccgg aactgcactt cgcttgtag attaaggctt gtcaataaca
17281 gaatcacccg agaatactct aaaggtattg gatttctca gaactattg tctctgact
17341 tgtctgagaa caatctctcc ggtccagttc ctttggagat aagtaattgt ttcccaactc
17401 agatgtcttaa cttyagcaac aacactcttc aaaggttatct tctctgtctt ttatcatcgt
17461 taacaaagct tcaagtaact gatgtctctt ccaatgactt gactggcaag atacctcga
17521 gtctaggtaga tctcaattca ctcaacccgg tcattctaa taagaattct ttcaatggag
17581 aaatcccttc gtctctcggt cactgcacga atcttcagct tcttgactct agcagtaata
17641 acatctcttg aactataccg gaggagcttt ttgacatcca agatctagac atctgcttaa
17701 acttgagctg gaattcatta gatggcttta tccgggaaag gatttccgct cttaacgcat
17761 tatccgtact cgattttctg cataacatgc ttctcaggca cctttctcgc ctgtccggtc
17821 tagaaaagct gtttctctg aacatctctc ataacagatt ctcaggttat ctccagata
17881 gtaagtggt tagacagctg ataggagcag agatggaag aaacaatgga ctctgttcca
17941 aaggtttcag gtcttctgtt gtaagtaaca gttcacagtt aactcacagc cgtgtgtgtc
18001 actcacatg actcagata gccattggat tgcataatcag cgtgcagcgc gttctagcgg
18061 tattaggcgt ttgtggcggt atacgagcta acaaaatgat tccgagatgat aacgatctac
18121 agactggaga aaactctatg acatggcaat tccacaactc tcaaaaactc atactcacag
18181 tcgaacatgt actcaagttg tctgttagaag gtaatgttat aggggaaaggt tgcctccggga
18241 tagtgtacaa tagtgcgaat cctaacacag agtcatcgc ggtgaaaaag cctctggcagc
18301 tgacagtacc taactctgat gagaaaaact agtcatcagg agttcgagac tcatctcagc
18361 ctgaaagttaa aactcttggg tegtacagac acagaacact gtacgcttc ttggagttag
18421 gtgggaacaa gaacactaga ctctctatgt atgatataat gccaatggg agtttgggaa
18481 gtttgcctca cgaaaggagc ggtgtatgta gcttaggatg gnaagtgtag tacaagatta
18541 tacttgggtc agtccagggt ttggcttact tgcaccatga ctgtgttctt cccattgttc
18601 atagagacat caaggcaaat aatattctga ttggccctga ttttgaacct taccattggag
18661 atttccgact cgctaagctt gtgatgatg cgcagtttgc tegtcttttc acacacttag
18721 ctggttccca ttggtacata gctccaggta atctccaaat ttctgttttc tcttaatta
18781 gttaaactga tacatcataa gttactagta gtcttagctt agtaattgga gaaattcagg
18841 ttttatagtt gtttttgggt ttggcattt tgaaaagaa agtattttgg tctcagttga
18901 tttcgatttg gtttaaaatt tttagaaaaa aatggtagaa gaaataatt ttgatttaata
18961 ttactatcat tactgtatac gatttaaaaa aaaaattgat tcatgtattt ctcgattatt
19021 tgcttgttagc ttggttcaat atttttacta tataatgat tcatgacttt catactaatc
19081 aatatttgtt cggttctgtt ttgattagtc tgaagtgaa tcatatagtt aaaaattga
19141 ccaatctatc atatacttac acgtacaag attttcgatg taatttttgt tgaagcattt
19201 tgggttttat atgtctatcc ttagtaacta acaacaata tcaaattgga agcataaact
19261 tatattatcc taacaagcta taacactata attatcgaga ataccgata tcaatgaaga
19321 taacagagaa aagcgacgtg tacagctatg gactcgtagt gctcgagatt ctaacgggta
19381 agcaaccacg cgatccaagc ataccagat gactccacat agtgcagttg tccaagaaaa
19441 tcagagacat acaagtaatc gaccaaagac tacaagcaag cccaagatca gagggtgaag
19501 agatgatgca acgcgtagga gtccgccttt tatgcattaa tccatacccc gagacaggc
19561 ctacaatgaa agatgtggct gctatgctta gtgagatag tcaagagaga gaggaatcga
19621 tgaaagtgtg gtygttgcgc ggaagtgtga acaatggaa agaacgtggc aaagatgatt
19681 cgactcttcc ggttatgcaa caaacggcta agtattttgag aagtatgagc acgagtttct
19741 ctgcgctctc ttgtctttac tcttctctct ctctgctac ttcttaagtt agaccaaatc
19801 ttaataataa gcatgggtta ttatgttttt actggaacaa gtttctatgt ctaagttttt
19861 tggttttatc atatagcag ttgagatgaa ccgtgtgttt tttaaccaga caaacagatt
19921 gagactaatc tctgtcttacc ttttacattt tcttctgtca aggcgtcaag ttgattttt
19981 tttgaaatac aaataaagaa aaaactgatt aaaaacagata tgaggaggtc ataatcata
20041 actgtgactc acgagataaa tcatgtatcg tggctgcagg tgcacgaacc ttttctgtt
20101 taccattttt ttccgagttt gtccaggaaag gcggaataat agttgacctt cactcataac
20161 tctataactt aataaattta tatcaaatgt atcaataatg taattataaa aactaataag
20221 caaaccttta accaagacca gtgtgtatat taagttaacga aataaaattg aacagaattt
```



## NCBI Sequence Viewer

20281 gttgaccaga taaataataa gtatagattc atgtttttagt ttttttttaa cttcaatttt  
20341 cagatttttt ttgtagtcta aagtttggtt gattgaaata aaataaaaaa aagaaaaaat  
20401 ctatattttt ggaatatatat ctcttacaata taattctttg caaatgtgga agacctatta  
20461 tacatataga catatatatt aacacaaatca ttatgagtaa gtatcttaab attataaatt  
20521 gatgaaaagt ttaaaaaagg tataaatcta tataattttg tctatttaaa aatggttgaa  
20581 aatcttcttt tcaaaaaatc aataattggt ttatgaaac agtgaagctt aatctttaag  
20641 aagctcccca accaattttt accataaaat aagatttcaa aattaaacct acctaatagt  
20701 tttatctata ttctctaact atccatgttc ctatggtata ataattctt ctaacattat  
20761 ttttgaaact tgtttctctc tagttttttt ttttctcaa agcgagatgc aagtttattt  
20821 caagagttac caacataaac aatggcaaca aataaatttt tatctatttt gtgtcttcg  
20881 ttgatggcgt ttgtgctgat tctgctgcca atgatttccag gtaacacatc tattatctat  
20941 taaaaatttg tttgtaattg tatgataaat aattctactt taaatatgaa agttaccgtg  
21001 tgtttctatg aaaaacaaaa cctaacctaa tttttgtgtg aaggttttaa tttagctcatg  
21061 ggtgttaaac ataaactatc aaaaaatata accaaaaaaa taaagttttt aagttgtata  
21121 tttactttta ttgaaaaata ttctacttcc actacgcaa atgtactcatt ttgtaggttt  
21181 tttgtgaagg tcacaaattt tagattacga gtgcaattag agtaacacaa attttgtatt  
21241 ttgaactcaa cagctcgtgt tttaaaaaata gtgtctcaaa tcagaagaag aattgtcac  
21301 tatgtagtaa gatttaattg ttttattgaa atgtatcaat tttattttaa gagtcgttta  
21361 atttgtttg ttataaacta aacaggacaa atcataacat tttaccggg ggtaccgtta  
21421 aacccatgct aatgcaatgc agcttgcaaa tcaaatggat acaaaggagg ggcttgggta  
21481 agcatgagca taggctcaac aaccgggtgt tgtgtctgta aacctaaatt caaatctcaa  
21541 gatctcttca gattctaatg tatcattatt aataataatt agtactcat tattatctatg  
21601 tatcttgata tgaataaaaa tatattcaaa tatgtttcga tattgaacca tctataatat  
21661 ggataatcat tctgaagtta tagtagcatc caaacatttg atcgtatata ttaataaagc  
21721 atacgaatgc tgaattttat gtttaactata ctcatagaaa tttatttata tctgcacatc  
21781 caatacaatt taaaagattt ttcaatttaa tgagagata ataatttatt ttgatatttt  
21841 ttttgtttct gttttgtcaa caatagttaa tatgccatg gtgccgatt ctataaaaaa aatcatactc  
21901 tatatcata atgtgtcaac ctctagttaa gtctcaacaa cataagtaag cataacagaa  
21961 ttaaaaaata atttaaatga ttttgatata atatatgaa ttaaaatttc aaaaattcat  
22021 aaaaattcat attaaattaa cacaatcaaa tgcgcaaaaa aaaaaaaaaa caaatataca  
22081 catccaaaaa ttgtcacaaa tctgtactaa gcccagtaa gtccttttgt caataagtc  
22141 attaagccca acaaacctct tcttctcttt cgttttatga catcaatcca tttagaaca  
22201 acaaatcaga cgttaaccga caaaaaaaa caaattgttt gactgatcgt ttaaacctca  
22261 gaaatggatc caaaatcgcc ggaatttatc tctgtcagc cgtctaaaaa cttaggattt  
22321 gtattcgatc agactctccg caccgagtc tccgccaac ctaacctaac tgaaaattgt  
22381 tgcgcagcgt tcaagatctt gcacggcgcc gtatcggtct tgcgtcgtga ggcactcgcc  
22441 agtctcggcg ccggaatcgc ctctgtttt aaacgcgctc cgtgaatcca tctctctatt  
22501 caccactctc gtctgtctgc tctcggcgaa atgtctctgc cgaatatttt tccagttcca  
22561 gtcggaaaaa atcttcaggt aacctataca taacctgac caatacacat atcacatcat  
22621 ctatctttag actttaatgt atataaaatt atattaacta gaaaagtttt tagtaaatat  
22681 aaaaatttaat atttgttcaa agtaattgtaa aataaatttt tttttaaagt ttaaaaaaaa  
22741 aattaattatt ttgttgaa ga ttaataat aattctgcaa tttgagagat aacataagat  
22801 aaaaatttca gttattagta gtaaaagtga caagttatgt ttataacca ttgtattttt  
22861 aaaaaatgaa atttgttttg atgcaatata gtgtgaaaag taagaaaact aatatatgca  
22921 ataccattat taaattgtga acctatttaa ttccactcaa aatgaaaatt gaaaaacttt  
22981 ttcattgatt tgatgaactt gtagagttga attgtataaa ttcccgcaa gcaaaaagca  
23041 cctgtgtacc aagttgtcgg aaacttttaat ttgttagca tgcattggac taacaaacag  
23101 caaatttttt attttattta tctgtagact aagactcaaa gagtataagt aatgtgaa  
23161 tgacacgtgt ttgtgagat ttgtatcta actagggtg tgaggtcgg ttatggaag  
23221 ctaagaaaaa cgagacaccg gacaacaaaa taatggtatc aagactcagg gttactctct  
23281 tctgcgtaatt acctatccg gaccattgta aagatgctcc agacgagctc aagaagtta  
23341 tatccaaatt atcaaatcaa atgtgatcta tctcattttg ttaggaaata gaacattttt  
23401 agtttggttt tatgtacga accaaaacgt tttaataaatt gtgtgtacca acatgagata  
23461 ttataaggtt aaaaattgt acaagttatt gtgggtttag ttgttagtc atttgaatt  
23521 atagtattaa tagactattt tgttgaaacg tccaaatcgt atagatgggg taacaaaact  
23581 aaactccttt atatgattct ttgcataaa attctgtctt tgaataact gaatcataca  
23641 tactttttta gtttgatcag cataataata ttaccacaac tatgtatgct attaagata  
23701 tctttcttca tattataaat gtgtatcaat accattttaa cttctgtata caaagaaaaa  
23761 aaaaagaagt gatattagtt ttcgatgga gatattattt ggttgtagtc tttctctgt  
23821 agacacaac acccagtggt gagtctaaat gtccacaatt accatcattt tgtataccg  
23881 tggcttaagc aaactgtatc acatacgtt gtacactcgg accgatacca catggcgta  
23941 tgaacgtcaa tagcgtttat tttagctga aaagatatca taaaattct gagattcga cagatcaaa  
24001 tagatcatca tttatgttta tatttagaga gctctacatt gagattcga cagatcaaa  
24061 cggtaagggtt acctcacgat aaactaacga agatagtga ttttgcgtga ttgattgaa

## NCBI Sequence Viewer

```
24121 tatattaaaa aaaatatata ggtatatata taacacaaagc aaggtaaatat ggtttagatc
24181 aagaaaaatcac cctaatagaa atcaaaaaagt ttgtgtatat gttttatgaa ttgtgaaatt
24241 gacttaccggc ccactcgaaa caagagaaaca attgttatggc cagtcagtat atgtgaaatgt
24301 ttctgtgtctt gcatctttttt gttttgaggt gtcaactagct atagactatt accttgtgtt
24361 ctcttttgttt tttagatgtat ccattgagat taaagtgtct ttatataggg gtttggtttt
24421 ccgcagataaaa ataatatcaat tttagctggat ttattcttat accccaaaaa taaatttata
24481 aagactgaaga tctaattttg atatgatggg tcatgaaagt tccaactttg gttagattta
24541 aacctgcata acgcattatg aaattgacta tatatatatt aagaaagaga taattctcat
24601 ttaatttttta gtaaaaataa tactcaattt tatattatac cagaaaacaaa gtgtgttcaa
24661 aaattatttaa ggtctatatca tatgtaaata caattaaaga acattttatg aagaaaatta
24721 gaaactgaaa ttcatatca agaattaaaca aagaaaacat taataaaatt tcttgttacc
24781 ttaatatatcac catttttatt aaattgaacat taaatcatag ttttttagat ttaagaaaaa
24841 agaactcac agaaatttct ttaataaatc taatatatat gaaaaataat aagaagcact
24901 actgaaagaa gatccatata aattaagaat agtttgggtc atgccaatca tcaatttgaa
24961 ccaaatccac atttaaggaa aattgatggt attctttaac ccgaaaacat gttcttgtct
25021 ttgcacatgt gtctgcacct attttttatt tcttcattca acagtgctaa atgaaatcat
25081 ttactacatg gatctctata aacgaatgtc ataaagattt tatggccttc aaacttggct
25141 tcaacatcga tcatgtatat gattatctga tctatccata tagaaggagg ctaaccggct
25201 ccttgttacct tgcatttggcc ttataaaaggt ttccaagaat ttaatttctt tcttatttga
25261 ttttaactttt gtaaaatgcca catttttccg tttaattaat tcttgcgtga gtgacaataa
25321 tgtttaatatg acatttatca tcaagatata taltttatca tattttttag acatgactat
25381 caaacaaacg tcacactatt ggtcagagat tattcgacat aatccataag aagttgttaa
25441 tatatcatca agtctgtaata aatgtatata aatatatta aaaaaaaaaa tttagagttg
25501 ctttttccgc attgtgcgga atagattctg atgccaaatt tggatattat caagagcaca
25561 tgtttttctct ttgtgtcgtgt cttttattta agagagaatc atcatctctc actctctact
25621 tctagaagttt aaaaatcac gccatcttcc gaattcaaac cgcgtatgtt tgtcgtttct
25681 tgtctcacat atttgaagca gcgtaaagat gcttaatttc tcaagttaacg ttccccaact
25741 ctcatctctt tacagtaaaa aaacgcccaa agtgagaatg aactcatgtg ccgactcaggt
25801 gttcgtataaa aagcttctgt ggagagatat gtcaacgaag atgaaattgt cttctttttc
25861 tgtctgcggaa ttacctgatt tgaggaaaag taacaagagg agggggatct tttagatgat
25921 caagtcgcaga gcccgcgagg gtgcagggtg gtgtgtggct gtttggtgat atgtgttttc
25981 ggttactact tcttctaagt atgaagtga tctctgggtt caaagtaata agggagataa
26041 gaatctcaag ctgcacctc ttcagtcatt ttgtatatat caaatcttcc ttttgttttt
26101 gtttagtttc gaaattcaga ttggttttag tgaagtgtg taataagttt tggttcttgg
26161 ttagggaact ttatttttgt tgatttttgt taataagag ctaaaaagat
26221 cgaaccttgc ttgtgaatgt cagcttgtat gagcaaaatg taacttttgt tctatctagt
26281 agggagatgg caggtcacat tggagggttc ttgaggaggt ctagcgagaa cagaggctca
26341 accggtcgaa aatttgatta gagagttggg tatccaaagt cgttatctaa tcttacaac
26401 ttctgttgta gtatttgaa tttagtatac ctttgttgc gaaaacatga atgtattgag
26461 ttaatttgtgt ctggttttat atggactttt ctttttctt ttgtgtacat tctcatgctt
26521 ggtattgcag gcccttttct cgcacacaga ctctctcgg gtgtatattt gtatgctgac
26581 attgaatctt cgttcattta gtgcaattgg atatgatagt gattacactt tgatgacta
26641 caatgtcatg tgaagttctt cctttctcag acataagtag gaacacatta tgcaccataa
26701 ttgtatgcta atttatggtc ttgttcaggc ttgtgaaagg aaggcttatg actatttgat
26761 ggaataattc aagagcatgg gtttccctgt ttgtggactt gttcttgatc ggaactgtg
26821 gatagtatta ttttctgtg aaagttagaa ggttccacat tgtagtacc tatctagttt
26881 tttagactact ttaactatgt ttctcttat taggttatc tatagggtc atgattgaca
26941 aagagaaggg ttatttagtt aaggccgata gattttgggt tglgaagaga gccatgcacg
27001 gtacaaagat gttatcaaat aaagctgtca ggttatattg acttttcaac acctattgta
27061 atgtgaagaa aaaaagagat tgcaaaagtc ttgtcttgt ttgaaaggag tctcattctc
27121 ttattgtctt tttagtgatg aaagttagat ctatggaagg gagttagtgt acctcggaag
27181 ccagagtcga tgggagtttc tcaatacatt ttcttcagtt tcagaggctc tggcttatgc
27241 acaggtacct ttctttatta caagaaatgt tagctgaaca atttgcagtt gattttggac
27301 ataattacct ttatttattt atttgaact catataattt tcaacttgt ttctcaact
27361 gatgactctt caccccaatg aactttcttt ttacaaaat ttctcgcagc gctctctgta
27421 agcaaatgaa gttcaaatca ttccagatgg tttagatgt ggaatgaga ttattttcgg
27481 cagatcttgg caactttgat tataaaggac ttgtataagg ttgtttcat agaacataa
27541 ttctattgaa ctgaaaattct tgcataata ttttttaac taatcatttt tccaggctg
27601 ttccaanaag ctctctcaga gcacatgttg aaggacaact taaggttgat agtttgcaca
27661 ttccataatc attttcttaa atatcagttt tctcggctc ctcttgaat atagtactt
27721 tgcagagtga gataatgtcc aagccggaac tatttgcga gccagacca gaactacctt
27781 tagctctttt agatcaaaag gaggtaatgg tttagtctt acacaatata tctaactatg
27841 atggttcaga ttgttaatat gttctaggc ttgtaagaag tcttctgcta
27901 tcaacaaactr ggattatcac taacagaca aatgatgaa gcactcattt aacaaattcc
```

## NCBI Sequence Viewer

```
27961 ttcccaatga catggactgg cgagatcttt ttgacatggt gagcaaacgt aaatttgttt
28021 ctgcattttg atgtgaatc tttgtttttt cagagagtag tttgcattag acttcaactgt
28081 ttttgactat atgtttccac ataggttgata gtttttcgca ggaacccaga gttcttccag
28141 atgtgcacc cttctatatg ggtttgactt ggagagggtt tgctgcgtcc atgcttcaag
28201 gctgaaacag gtaattttct atgttttcta ctgaatttaa gctgataaact ttaccgttga
28261 gaaagctaaa attttttccc tctcactggt ttgtctctat aggaggtttg tactcaggag
28321 gaagtgtctca aatgatagag agttcaactca acggttcagtt agatgcagtt ttgtatgtgt
28381 gtgaccacat ctacactgat gtcacgctat ccaaagttcca cgaactgtgg cgaactgcgc
28441 tgattttgcc tgaactggaa gaagaggttt ggataaatta tactctgaac tatatcaata
28501 gatgacagaa cttgaagggt tttcgtttcgt tgaagtaaca taaaaaccct gattttctct
28561 tgtttctttt cagtatatgg ctctaatggg cagctcgtggt acccgagaag agctaataga
28621 gcttataaat caaaaagagg ttgttgggga tctctttaac caactctcggc ttgctcttca
28681 aagacgaagc aaaggccgtc ctgctcaggt ttgtttatct ccttacaagc tctagagtct
28741 atcttctctc cctgtgtgga acacaattgt tatgaacaga taaatagata gagatttgtg
28801 acggaaattg cttataaaga tatcgttttc tcttttcttt ctagactctc gctgctacca
28861 acttggtatg tcaagaactg acagagacca tgcaaaagct tctttatcta atgcaaaagc
28921 tagatgcaca atgtgtgtcta atgtcggaaa cagatggcga gctctttaac aaaaggttaa
28981 aagaaaaaga caaacaatgt tagtaagatt actcaagatt cacagaaggg gatattgtctg
29041 agccataatt ttgggctgtt ttgcaggtgt gggcttctct ctaacgcggg gttttgggga
29101 taaaagccac ttgatgagac aaatcgaaaa gttattatgc ctctctctct catttccgag
29161 taactcataa gctcaaacca agctcacatt taaactcacac tctatttctt caaattctgc
29221 ttcaggttatg cggatatata cacatacaaga gctctcaact gctctcaact ctactcaact ccaacacctt
29281 atgtattttc ctctcaacga gcaggttaaaa agcttaaatc agattattcag tatagataca
29341 aacgaatatc taggcagatt tcaattaaac tcaactatgt ttttctctc tctcagctgc
29401 actcgtctcag gattctccgc ttccagatgc gggatagaaa aactagcggg tgtgaccaaa
29461 gttgagtcga ttgtgtgttt gtaatttttt ttgtgtgaaa aactgtggag aaagaaagat
29521 taacaataga aattctaaag ctcaaaagtc tactttatgt ataaaatgtg tattctattt
29581 ccttgaattg aattctaaag ttggtgtgtt ttggtatgtt ctgattgttt atgttattgt
29641 ctattctttt ttaagctttt tcaggctgac acagaggcca gattgtgcga atgactactt
29701 ttagaaaaac attatacttt gaattctgat cagctgttata tcaaacatac taacacatac
29761 gacactagac agtgcagact tctataataa caatogattc gttgtgagca cacaattcat
29821 tacttcaatt ccaagagaat taataagaaa tctctcaatc tctttattat tcatgtatct
29881 gatctttcgc agatacaaaa tttgtgatat tactatacga aacatctaaag acctatgatg
29941 cagaaaaacc accggttttc ttacgtcaaa gtttgaacaa taccgactgt tctctgtcgt
30001 tcttattcgc tcttctggat ttgttttctc caagcaaaaa aaactccgac ttccaacgaa
30061 aactcagtcg ctctgccttc aggtactctt tttttatcac ccagcttaag ccaacttaag
30121 ttttaggggt caattttgga tcttaggggt ccgtaatttt aaagattctt aaacattctt
30181 ttgcctttgc ttactgagg attagaatoc caatttgagt tggataccta aaagtcagat
30241 tcttttttgt tghtaaattg aaggtttttt gttattggca ggaagaaagt gatagctata
30301 tgtcagtcag gaggagaggt tghtaaattt aaagtgtgct ccttagctta tctgtgtggt
30361 gatgcttatg cagtagacat cgtcaggagc acttgatga gccatttcaa atccgaattt
30421 gctgaaaaat tggggtttag ttggggaagt atgacactca agtacttctc cctgtgaaac
30481 aagaaaaacc taattactat ctctaaggat aagaattttc aaacgcatgt tccgttttcc
30541 gcagacgcac caaatgtcga aatcttctgc tactccggaag aaacgcgaagc aaggaattgc
30601 tccaacatgc cagctagtat gttgtgtgtc tagtttcaag acctaaacgtt ttatgcacat
30661 tgtttatgat tghtaatgact gatgtgtgga ttctgttatt tgacagatga agtaggacaa
30721 ctgcatcaga agcaatggta cctgtagtgt ctgcaggaga tattgtttgc ggagatgata
30781 tgtatgacaa gatttttcag attgaaatgg tataatgcc tgaagaaagc agccotttgc
30841 cgtgtaattt tgttttgact gatgagaaa acataataaa agctgcacagc agctgtgaaa
30901 acgctattac ggggtttgat caaagggttca atagctttac tgagtttcga gttgcgttgc
30961 acaagactc atgcgcacat ggggtttactt acaagatcaa cagactacaa gatccctcgtg
31021 tttcagtcga atgcgaagct caagggttgc cactgcatcg aggcctttca
31081 ctacgcgcaa gatttgcatt aagaaaaatga atcagagcca cactgtgtag agagcgggtg
31141 tgaagctggt ttaccgtgca agtagaggtt gggttggaag tateataaag gagaagttga
31201 aagcttttcc agattacaaa ccgaaggata ttgcagagga tatacaaaaga gagtatggga
31261 ttcaactgaa ttactcgcag gcatggagag atcaaacctc ctttttctgt ctaaaagagat
31321 gttcttataa aaagccatat agtcaacttc atgttttctg caagaagactc agggagacaa
31381 atcttggaa gatttgcatt ttcatgacaa aaacgaattc cagctttccat cggcttttca
31441 tatcgtttca tgcattcaata tctgggttta gacaaggtt tctgtctctc ctttctctgt
31501 acaccgcgca cttaaacctc aagtaccagg gttttatgct ttgtctacaa gctcctgata
31561 ccgaagatga aattatttcaa gtacattttg cagttgtgga tgctgagaca gaagataaact
31621 ggggtttggt ttggaagcac ctcaaaattgg cactgtgcga tcccgcgaca atcacatttg
31681 ttgcagattt tcaaaaagggt ctgaagactg cattacctct ggtatttgaa aagcaacacc
31741 atcatgcta ttgtctgcgc ccctcgtgct agaaatttaa catggacttg caggtcagat
```

## NCBI Sequence Viewer

```
31801 tttctcatga agcaagacggt tttatactca atgatttcta tgcgtgctgt tatgcgacac
31861 aacctgatgc atattaccgt tccttggaaa atataaaaag catttccctt gatgcttata
31921 ctgtgggttat cgaagatgag cctctccatt gggcaaatgc tttgtttgaa ggagagagatg
31981 ataaccacat ggaactccatt tttgggctag atttctacag ctgggtttctt gaagcacatg
32041 agttgcccat aactcaaatg atagacgaac ttcgagcaaa gttagctgcaa tcaatatata
32101 cccatcaggt acaatcccgg gaatggattg tttcaacgtt acacaacacc aacgaagaga
32161 agctacagaa agaaatagaa cttgcaaggt cacttcaggt ctacagcgctt cacaatagct
32221 tattcgaggt tcatggtgaa accatcaact tagttgacat caatcagtgt gatctgcgact
32281 gtaaggtatg gagactaacg ggtttaccgt gtgcccacgc agttgcagtg gttgaaatgca
32341 tagggaaaaa cctctatgaa tactgtctca gatactttac ctgagagagt taccgttttaa
32401 cgtatgctga gtccattaac cctgtcccta acacaacaat gacgatgatg attttggagg
32461 agccgcccgt tgaagggggt gtttcggtta ctccaccgcc cacaagcata accaccacag
32521 ggagggcaaa gagttaagcag gtggaaccgc tggacatgtt caagctcag ctccagtgtta
32581 gcaactcgaa ggggttagga cacaacaaga agacctgcaa agctgtatct taagtattag
32641 ggagagagag agaaatagag agggagagaca tggttaggac aaattgtgat ttatatagct
32701 gaccttggcc aacttttgg ttttgggtct tctgtttgtt ggggttttga gaatgatctt
32761 aaagggttaa aatctctctt aggttgaac ccaaaagcta agagagcaaa gggcgctcaga
32821 atctctgttc ttaaatggca ttaacattga tgaagattt tacataaatt ttttgtgat
32881 taatttgttt ttttaacaaa aatttcaatg aatcttttca atatcatata ttttgtattt
32941 cgagctttac aaaaaccacc acactgttta caatgtgcgc agctgtgcgc acaaaccttt
33001 gacgtttaca ttcacaaaaa gatcgatgtt catatgagcc tataataaatt gcttagagaca
33061 gtaagatata agtatcaaaa gagttaaaat ctatccaaga caatagctcc aacaaactcg
33121 aggtaaactc ataggccaca aaggttacag caccagcagg agcagcttct acagctcgatg
33181 gcacaatgcc ttgttacaga ccatgccacc ctctgatata ccaatctgtc caagccgct
33241 ctaaacatgtt tctgtatgca cgtcgtctca ctccggctcc atactcttga tgcctctgta
33301 aacccctcaat tgcacaacaa agaactcttc aaaaactctc tggcatttct atatttcat
33361 gtttgaaggg aattgcattt tcaittcggt ggagatccac tggcaagttt gaaaagcctt
33421 tatattttaa agtcttttca cacnaattgg agtttttctt ctgaggtac gttgacagaa
33481 caagttgggtt acccggaatc agtaagtaga ctcagccagg acatatcata gggcagtaaa
33541 tgtgaaagct tgccttaaga tacataaaga tagaccocaa acatgaaac acatgaaac
33601 gataagtcac tgccttagtt cagtgttaga gcatttgaat taccttagct ggtttgtagt
33661 agatagtaga tggatttgaca actcagcaac tatagttgag taagtttagca atagagcttaa
33721 gtggctaaaaa ggaagtttgg gatgtgtaat gatgtgtaat agagattctc attagcaag
33781 tctctctatg atgtgttagtt ctgtctgaag tgaataaata aggaatttaa ttttggat
33841 cctatcacat tatcttccca gagaatccat tgggttaaga ttaactgacc tggaatcgctt
33901 ttttgaccac atcaagggga tggcagacga gcttgcgcgt agtgcgcgct ccaagccacc
33961 aaaaagaaa gtggaagcta tggaaagttg tttccacatt gatcgggatt ttggaggata
34021 atttggatcg atttccagtc ttgttataaaa taactcaatt agttgatatt catatgaaca
34081 ttgatcnacg ataaactata tattcctatg ctcttaccat ctctcaacgc ttaaacatat
34141 catatgtgccc aaattgcagg ccagcgttag gcaaatcttc aacagagttt ggtgttaatc
34201 catatataca tctcttata cctcgagact gaataatct cactcaatct gaccttaatc
34261 ttggatatac ctgttgacag tcatcagtcg gtcacttata tgtaacttaa gcatgaataa
34321 ggaagaaaaa acagaaaaat cccagtacat ggaattttaa tatatttaca gaccttaacg
34381 tcacgcaact aaccttaggc tccccctgtg agggcaatat tgcctttaa agatcaaaag
34441 gataagcttc tagagtacga gcacatcctg gcacatcctc atgcccgtcc caagcgaag gatagataag
34501 ggctcaaatg tatatgatcc ctctcattgc cataagatca caaccagcta ataatgaag
34561 aaccacattg atctctaaat tatcagagtc aattccaaaa gctattcaaa gaaatcacaa
34621 attaaccaaa agaatgtata cccctaagat agccaccaga ctacacaacaa aaatagcact
34681 ggcaattctac aaaaaccat atctagagag accgcaatgt tactagtag actaattac
34741 ctgttttagt agaaccagat gcaaaaagatt tcagettatg tagtactgta aactgaattg
34801 aagtatcagg catgaccatg agcaaatgtt gcacattacc ccgcgcaaaac ccttaagaaa
34861 aacaagtatc aacatcaaaa acagagaaca agacctcata aaaaactaatg cccacaacct
34921 gaaactacta cataactata aacaaaaaag aatgtgtttc agaaaacgga aatgttcaag
34981 agacggacc gaaaaacctt tctcttaaat ctgtctttag tgcgtctgac catcttcaga
35041 tacttagaag ctctcgacaa gtttccccga accaaacccc atgaatgtgt cgttctagc
35101 tgaacctgca atcacatata acattcttca accatctcta taatatataa agcttataag
35161 tttctacaaa tgataattct catctaaaac ctccacataa caatccaaag caccagcaac
35221 aactcaaat attcttttagc taaaacaaat aaaaactata actttatcac tcagagataa
35281 gaaaaaaaaa gccaaacctg aatcttaact ttaattgacat caagaggaga agttacagaa
35341 cgagaacac caccagagat agctccggca gaagcatcaa ctacagcag ttttaactgt
35401 cccggtctat catcaacctg cgcgtctgta ggaatcatat ttttctaggg tttttggtct
35461 ctataagcta aatatcaata aaaaagtttc ccaaacctta tcatcaatat aataaaacct
35521 aacacaaata aggaagaaaa ctaaatttca ggaacaccta aacgaacaa acgaaaactg
35581 aaactctttt taccagataa agcgagaaac gaaaagggtc gacatttgat gagtttccgg
```

## NCBI Sequence Viewer

35641 ggaaaaagca aggaactttg agatgacccct taaacgtttg ccgcggtatt gagagttggg  
35701 tggagattga aaatgatttt actcggaaaa agattataaa aaaaaaaga agctgagaga  
35761 ttctttctcc gtcaacgttc gtgttctctc cgatagagag attaaagctg gctttctacc  
35821 aaaaattgac ctcttttaga tctgttctcg ttcggtttaa tgaagtatac attcgtctta  
35881 aagtaattaa attgtttcaa aggggtttta tggttactct taaacctctc tccattacgt  
35941 tgtgtggatg gatttagtca ttactaatt tgaagatcaa cagagacctc tacaanaaga  
36001 aaattactcc ttatgttcca atatataga tgtttcttgc ttcttcttg tttcatttta  
36061 ttatattaag aaaaaacatt atgtataatt tatgcatttt taatagtttg tgaattattc  
36121 tattctgcaa tctttttctc ttgtattaaa cttttttaaa taaagttaga taagattttt  
36181 tttagagaaat acactaaaat acatttatgt aaaaataata aaatagtttc ttattttttt  
36241 taatgatttt agctaaaaca ttttctatg taaaatatga ttttcatatg tggtaactca  
36301 agtgttttga acacatcgac atgtgtcaaa tctaattttg ttaattatta aaacaaaaaa  
36361 tatagaaaaa ttttaacct atggaagaat gaaaatacat acaaaaatca caaaaagtta  
36421 tattcttatt cataaatttt taagtaataa gatttagaat aagttaatgg gatgtaacat  
36481 tgcataagat tagaatactt tctctttatt tttgggttaa agttgtagaa taattaatta  
36541 gttacaaaaa aaagttttta attttttgaa gttaaagtgt agactaatta attatttaca  
36601 aaattttttt tatttttttt ggagtaagtt gtagaataat taattagtca caaaaaaag  
36661 tttttatttt ttggagtaaa gtgttgaat aattaattag ctaaaaaagt agtatcgttt  
36721 tgttaatttt attatttatct ctaaaattgt gtccaacaag gcccaatttt tgaattgaatt  
36781 tgcctcaatg cgggattccc aaaggttgtc acagcgttcc ggcgtcaagg accgccaacc  
36841 accatggaa gagaagaagc tgcgaacga tacgactctg aatccgtcac ttccattatga  
36901 tctaatactg atcatcttag ccgcgtcttc gagatcgtag tacacaatct tctccttagt  
36961 ctccaaaagc tccgatcaa tcttactctc accggagctt tacaaaaacc gaacctctct  
37021 aggcaaaacc gagaaatttc tctatgtttg tctacgtctc cctgtagaag ctaatcctcg  
37081 ctggttcaact cctcacagga aacctaatca aacctaaacc gatcacacca ctaagaaga  
37141 gaagaagaag aagaagaagg aggagaagtc aagtgtaaat cbgttggctc caatctcaat  
37201 tctcaattct catcctgtgg aatggtcagc tatcataatc gtgtgatcatt atctctacgc  
37261 cattagtgca gacatcggaa aggcctcgta ccttaacgtt ccttaacctg aattgtgaag  
37321 tcaacatctg aatgaagctc caagaatgag attggctcac actaatagtg aatttgaagg  
37381 aatatgttat ttaccagaaa gctttgaaag tccagattct ttgaactgtg ttgaagtgt  
37441 caacacaatg actcagactt ggaiaacggg gcccccgaag aaacgttagt tcaaattgga  
37501 aaaaattgaa aaaaagattt actacaagag ctttcattta cagcttaggg ctggaaagg  
37561 tctatctctg agctataaat cgaagcattt gacatgtgga tggtagtagt tggatacgtg  
37621 ggaattccat ctctgcctct ctgtatgat agagaataa gcttattttt atctgaaaagg  
37681 aaattttatt tggagaggac ttgacggtaa gatactgggt tacggtgaaa tagaggggtt  
37741 agaagggtca cctaagtttt ctatagattc tagtgtcaa tagtgccgaat atggcggaaa  
37801 cgtgtgtgtt ttgtgggaca agtatgtacc ataatgagga tatagagaga agatgatttg  
37861 gtgtgcagag attctgcctg aaaaacgtaa cggtaaaag atttggggga acgttgagtg  
37921 gtttgatgct gtgtctacag tccctaagtc ttacaaaatc ttgtgtgcta cgtcgtctac  
37981 tctttgattc gatgactcgg gaacttgaat ttgtcaagct tttgtattgt tatttttctc  
38041 tcaagtgaac caattcatba aactcatatt coatttgatt tcatcttaa tactccttaa  
38101 tttttttcat agttcaaatc aagttagatt tctacaaat gcacacaaaat cgaaatttgt  
38161 gctttacttg atgttgatga ataggttaca aatagttgat tcaattgaaa caaatccac  
38221 tattgtttta aggcccatla agaatacgt agaatcaatg aaaaatgatt tttttatcat tctataagg  
38281 cccatgcgtt tgtgttcaat agcagtgaa agcagtgaa tgaagtgtc aaatccctaa atctccctc  
38341 cctctgcgtt ctgttatctg caagtcgga aagaagcaga agaaatcaga aatcgtccc taatctctcg  
38401 ccaatgtcgt tcccgagaaa gaagaggaag gagcatctta gccccgctct cgagatcgta ttacaaaaat  
38461 ctccagaaag atttgatagt tctcaaaaac ctcccgtag atccttactt acccggagat ttacaaaacc  
38521 ctcccgtag tctcaaaaac ctcccgtag atccttactt acccggagat ttacaaaacc  
38581 cgaacctctc taggcaaac ccgagacttt ctctatgtct gactacgctt cctctgatga  
38641 gtaactctc gttgttgat tctctaccg aagcttaate caaacctaac taagaagaag  
38701 aagaagaag aggatccaag tgttcatctt ttggctccaa ttccaaattc caattctcca  
38761 actgtggaat ggtccaagct agtcgtgttt ggttcttate tgtacatgca taaccgagac  
38821 atcaaaagat ctccatgctc taacgttttg taccgttgaat tgcgaacgcc cacttggctt  
38881 gattctccaa gatgtgcact ggctcacatc aabtagtact ttaattggag cagctatttc  
38941 ccaggaaagt ctaaaaagcc agattttatg aactgtgttg aagtgtaaaa cacaaatact  
39001 cagacttgga atcccggtcc tccccaaaaa cgaaaactca aatcggaaa tatggaagga  
39061 aagatttaca taccctcttg tcaagagaa ctgtaggaag caataggaag aaaccccaag  
39121 gttttgacat gggaaagcgt aggattgggt ttagaggggt ttatttgatg  
39181 atagggaaca taccctattg ttatgacct ataggtgag taagtgtaac  
39241 actgcccgaag gagactggag cagattagag ggtttagaag gctactacaa gtttctgata  
39301 ttaagtactg ttaaatagc ggaatatgac ggaagctgg ttgttttggt ggaactgata  
39361 gtagctgcaa gtgggataaa ggagaagatg atttggtgtg cagagatttc gcttgaaaag  
39421 cgcaacagtg aagagatttg ggggaaggtc gagtgtgttg atgcggttct tacagtctct

## NCBI Sequence Viewer

```
39481 aagtccttata aattcgtctg tgcataaatct gctactgttt gattgatgac tcaggacata
39541 ggtagcgaag tctctttatt tagtatgtca cagttgaaag aatgtcttgt atctttttcg
39601 gaatgcagtt gtcatctgtt tgcctttaca aacacctgca ctcacttgtc tgtatagatgat
39661 taagctctatt ttgtgaactgt ctcttttgata attcctttga tgcacaaactc ttctctccatc
39721 tgtttttttgc tctgaatggt gtgtgtgcaca atgcctatat ttatgtgaatc atgatgaatg
39781 cttctctcat gctttttagcg gatatttgttt cacttctaaa catatgcactc cacttgttaga
39841 ctttttcaagc ttctttttgct tatgagaata atcttttttg ttctgcatac aaaaagactgt
39901 ttatcttttac attttcacgt aagagctaga acaagatata cagttagtgc ttgtttttgt
39961 gttttaaactg aggccttatgt tcatgtcttg tacttgtagc caacctaaccc tacaagaaga
40021 ttgtcactgt aacaatttga ggatcttgtg tgaaaaatcag ggtggacttg tcaagccact
40081 cttctctttc ttctcaatcg aacactatta ttatccaatt atatactctc tattttttat
40141 gattcaatcc aagtttagatt ccattatgca cagaaaaata aattgatcat atgtccaaag
40201 cttcactgtc atttttgaaat ttgttgtaaa tttttgaagg ttaacttatt ttgtgccatt
40261 gaccactatg atgcatttga aagacttttt ttttgttgtt gaacatacgt gattactagc
40321 ttcataggaa gaaataagtg taacaaaaata tatatatttg acgaattttt ggtcctctac
40381 tggtaatttt gtgttgtgtt gtagaatttt gtggtgactc aaaactattg ttttaaggct
40441 cattaaagtt caatacaaat gaattagata atctcataag gccactcaag attcaatcgca
40501 aattattttag aattatttcta aaggcccaat aagattcaat gaaaactctc aagtcggagt
40561 aagaagcagaa gaaatcgact aatctcgcca atgtcgtctc cagagagaaa gaggaaagaaa
40621 gctgcgtctg accctgcacc ggaatcgacc cgaaccctgt tactccccga tgmtttgatt
40681 gtgagcatct tagcccgagt ctctagattg tactacccaa ttctctcgtt agtctccagg
40741 accgtctgaa cctctgttac ttccgcggag ctctacaaga ctgcattttt ttccaacga
40801 acggagagatt gtctctatgt atgcttagat ttccctcctg accctaacc accgttgttt
40861 acactctacc gaataacctaa ccaaaaacatc accgagaaga caaagaactc aagcgggttt
40921 gttttagccc caatcccaaa tcaccattca catagtctgc tatctgtctg tattgggctt
40981 aacatctatg ccatgtgtgg atccatttag aatgcacctc ctctcaaatg ctcaatttta
41041 gactgcaggt ctcatacatg gcacgaggca ccagcagagg ggatgaagcg gaattattcca
41101 gctgctaatt gtgttgtatg gaagatttat gtacgaggag gcttggaaag gtgttagtca
41161 tctaaattga tggaggtttt cgbatataag actcaaacct gggagtttgt ttttaagcct
41221 tagcgggaga gatttatata taggagcttg gtgatagaag gagaatctca cattttggg
41281 tagaaggttg tgacttacaa gcctaaagaa gtatagatgg gagggttagg agaacatcag
41341 agtatggatt tgggatttgt tttctattct tattgtcgtga tagataatgt actgtattgt
41401 gctgtcctgt gtggaatcaa atggttatga atgtgttagc gactgtggag gaagttagg
41461 ggtttgaagg gattgtctaa gcttgcctag agctgtgtta agttagcgga ttatggtgga
41521 aagaatggctc ttctgtggga caagtatatc ctttgtagtg ggaataagag tactcgatt
41581 tctgtgtcgg tgattttccct tgagagatgc aagaatcagg gtattggggc aaaggttagg
41641 tgtgttttag tatgtcttac ggtacctagc tcttataact ttgtgggttg ttctctgct
41701 actctttgat gatgattgcc atacgaggca cacacatttg attgtctagc tagctttaca
41761 ataatttgtt tctatgttgt aataaactaa ttagtgggatt ctcttaccac tatatttcca
41821 taatgcaact ggcctctatt ccttctcact ttttgttttc ttgtcaagat caaaacaaga
41881 ttatcaagtt atagcgtctt ttgttgtcta caacttgtat ggacagggaa gtgagaatct tttttgcctt
41941 accttcacgc gttgaaacaa acatttgtga tgagagtaag agcaagaatt tcaaaagaaa
42001 ctcggtctta gaagaatgt acattgtcga aattgcacat aattgtactt tgactcaact
42061 caaagaggaa aagcgaaga aattgcacat tgaggttaa ttgacaacta taacaaacta gaagatttaa
42121 gctgtgtgat taagtgttat ttgagctctt tttagactct atcatatgt gaataactaa
42181 gttgcaataa gctagcgtaa gctaaggctt atatagatac tctctgaata caacaaaatt ggacacaatg
42241 agccaagtgc caataccttt aatatataat aactaaaccg atttctcaat cacaactttg
42301 atctgagttt aatatataat aattaaacct aaatctctct tcttgcactc ttbatattgc
42361 ctaaaggctt aatatataat tttttatttc ccctcgagc aatgtgatct tgcctgagc
42421 gccaaaccgc cagttacatg gttttatttc gttagtacat tgacttaagt gcgtcgaaat
42481 ttccaagcgg gagagagaag gattacgaaa gattgatgtc gacatcggaac gtgtgactat
42541 cggaaagcgg aggatcgcg gagaagcgg cgccggagat tcttcagta cgttagagctc
42601 tgaagagggg gtccgcgtac cagatcgaaat tagtggtaaa atctctaaa ccttaaatca
42661 ttgtctgatg attccaatgg gaaattaaacc actaaatcag aaatcttttg gtctatgatc
42721 ttctgttttg tttttccca atctgttttt ttgataatgg atctgtttt agtggatttc
42781 caaattagac ttccaatttg ttagaatgtt tcatgtctt gtcactttct agtgggtgca
42841 ttaaacttag ttagagttgt taaataaaca ggaagaataa attgaagatt atgtggaaaa
42901 atcagttttt gaaattgttg tctctgtatg tctctcataa tctagatgat ttgtggaaat
42961 tggcatagat cctctgtggg ttaggttttt caaggtttt tttttttccc aagactttca
43021 actcagatca tatctcagag ttaggttttt tcaagttgtc atgtgtgctt atgtactttc
43081 tattagtttt gcgagtttat cactcccaat ggtgtttctt ttgttctgtc agtatagaaa
43141 gctttttatg catcacaaga agctcaagaagg gacttttttg atgtttgtct gcttttgtct
43201 cacaacctca gctcaagaaga tgcctgactt tatgttaatt ggcctgtctc ttccttgtat
```

## NCBI Sequence Viewer

```

41321 gcggttttggc ctaaattgta taggtttggg ttagaatctg tgtggaatag atagtgttta
41381 gagatttgct tggctactct ctctacatgt ttgagttgtg aatgggatac aatgaactta
41441 catatttaata tagttcttgt tcatggagta ttggaactgt cccgagaaat gaiaaaacttc
41501 aggtagccta ggcacttttt tcatgaatat ttgttttagg ttgttttttc agacttttaa
41561 ctcaattttt ttcaaattta cgttagtggc tgcgtttgtg agacataatc ttgaagtatc
41621 acaggtgtgc tgaatgattg gcaaaagcatt tcgaagagac tgtactgctt aagctgcgcc
41681 agaatatata cgtcggtttctc aagcaatccc tataagtgta agttagtctt atgggtacct
41741 cccataaatc tattactttg gaccataatc tcagttgggt aagattgttt gctattatat
41801 ctgagcattc ccttcacagt agctagagtt gtgtttgtgt agttagtgac tacactgtgt
41861 aacttacagt gccgcagccg cattttggata cctttgttgt ttgtagaagc aagaactttg
41921 tctccctcaa tctatacgaa gagttagcat tctcttattc ctccatacac atataatat
41981 acacaaatct tctgtaatct aagttgtggg ttgcctttac ttacottggc agaggagaga
44041 gccccgagac tgtggagatg gagcgcgggt atctctactt caactgttac aagattgtaa
44101 aaagggcaat cgaactctgc caaattgact tgatctgagc tcgctctttt ttctactcaa
44161 atctctcttc atagaataag ttccggatga tgcctaccag tatcactgtt tatgatcttg
44221 tgttacttga gttgtatagt gattttgagt ttctgtttct gatataaact atgtgttag
44281 tatcttttga gttcttttga gttgtgccta gcttagctat ccattgttct aaccataacc
44341 agtggcggag ccagaacaat gactcactgg ggtcaaaaaa aagttaaagc aacgataaaa
44401 ctttactata aataaataaa ttgtcattaca aataaaaaact ctaaagatga actctcagtt
44461 cactcaatca ttggaatcgt tttactcacga aatcagttga agctgtttaa gactactgtt
44521 ttaataaaaa caaacaagac aatcacttaa gaacacatca ctgactacggt tacgcaatgt
44581 agtcttcaca aatttcacg ccgaaaaaaa cttcttcaga gtgtccgttg cacaaggag
44641 aatcaaaagt agctttagaa gacgatatac caaaggatgt gaagatgct tttttgtctc
44701 taccatacac cgagcaagat caccagaaga ctttaaatct caactttttt catcgttttt
44761 cacattatca aggtagatct caaagttagt atcaagatga ctacgtccca caaaactaaa
44821 tgcactctgga taaaactcga tcaatctcac ggcactgac ttactcaaat aacgaaaaag
44881 atctatcgga ctacagaaat ctatgggtcc taactaaacc acatactttt ttttaaaagg
44941 gccctgtctg ttatttttta tttctctaga tcttttttga attgggtccat gatgactgag
45001 taagacccca aaagccgttaa aaagtgaagt aagattggaat ctttataagt taaaaacaaa
45061 aggtcagtac taataatctc agttacggtt acgtgtgcgc ccatagaaaa acaaaacttc
45121 tgcctagctt ggtatggaga ggttaagagt ggtatggagt tgccttttca actctcgaag
45181 ctgcttttga atctctagt tgatttcttt gaatttgcat cgtacaagtt taggtttttt
45241 ggattttggg ttctctctaa tatcttcttt gaccagtgtg ttgtgtgttt ggtatttttc
45301 ttttggatct ttgtttctcc agaataagta cagaanaattg aatcttttgc aaatttttgt
45361 tattttgagt tctttagtga actcagttgy ataacagtga cacatatgag atttgagagt
45421 aatttatttt cgtgtaaatg catagagatt attaatgttt ctactagttg tgagaaatgc
45481 aagcgtattg cttcttgagt ttgtgtttaa gttagcaaat ttggtactat ctctatgag
45541 tttactatgt tttcttggtt ttatttgtag gatttaagat ttgcctgctt ttgtattctt
45601 ctttggctct gtaattgcag aattgtttct tctttaaagt tgatttttct tgcactttct
45661 aaacagagat cgaggtgttt ttgacgcaat agaagatcgt ttctgctgtt attatccaa
45721 aactgtctta cgtttataaa catttgagca gagatgcatt tctggccact ttgtgaactt
45781 cgaaaactct tcaagagcac tagccagaag agcatatcaa ggaatgaac ttgggaacaa
45841 agtcaaaaga atcagcagaa actccttgac ggcagacaaa aactctggaag ccccgagatt
45901 tctgtctgtg ttgggttttcg tgtttgtttg agagatgttg ccactgttct tcttctgtgc
45961 tactgtctgt ttctctgtgg aggttaataa actctctga catgtttccc ttctgtctgc
46021 atctgtgtga tctctggacc atctacatct attaacttag ttggtatcag tttagaattc agattaaccc
46081 atggcacatt ctcacacatc gatatacaat ggttttctct tagtgattct tctgtctctt
46141 atgattttta tcttttagca gatatacaat gattctgatt cctctgattt gttcagcttg catttgatgag
46201 tcaataaatt cctcgattctt gtagcattca ttgaagctcc ttgtggagga tttaagtga agatttttag
46261 gaggagattt atcaatctct caaaaatgta tatattatga ttgaaatgt taacttgaat
46321 ctttagaaaa cttcttccaa atcaaaatgta ttataatga ttgaaatgt cgtatctgac
46381 tctttttcca cttcttccaa ggttaaatgta tttaaatga ttctctattt aacacaaact
46441 tttatcaaat agataagcat ggttaaatgta tttaaatga ttctctattt aacacaaact
46501 aaacttaaac agacacaaat ttgttctcat tagaatttta actattcatg gcaattagga
46561 tttagaactgt tgaattttccg tttaaaatta aatgtcattt ttaatgcat tttaaaaaa
46621 ttatttactt ttttatgact ttaattttaa tgtattttaa taagaaaaat tataaaatat
46681 attataaaat agtaaaaaat aggtttttca tatgtatata aaatttttaa atgtttttt
46741 tttataaaac gatattttatt tataaacgga aggaatatgc ttttatgagt ttttttttct
46801 tttttttttg tcatccaaaa ttaattttaa ctcaaacgag ccaattctga gggatattgt
46861 ttacacatgt gatattgtgc ggttcggtac gatttttcgc cgccaaatta ttctgctttt
46921 ctatagcact tcgagaaatt aaagataaag aaaaatttgt tggactcttc ttgtcgtctt
46981 gcaattcttc caaatcacaa gaaaatcgcc gccactcggt tggacaattt tatgaattat
47041 gagttagaaa tttagaatat ttaggaaatt taaggaaatt tagaagagca
47101 aaagtcaaaa ttactaagtg agagtattta ctgactcag ccaaccaata ataatccgac

```

## NCBI Sequence Viewer

47161 acgtcatata aaaggtccag aaaggtcaaa aaaaatcaca cacattaagag tgctttgaat  
47221 acaaaagcag actccactat ccaattaaat tacgaaaata ccacttcggt taactccact  
47281 ctttttcact tcttctctgc cgtctgctgc cggagctcgc gaagctctct catcagaagt  
47341 caaaaaccct aaaaaaagtc ctttctgatt catctataca gacacatact gtattctcaa  
47401 aacaattggag attcctcttc tgaataagca gcaagagttc actctcgccct ccgctcactga  
47461 tctctacttc tctctctcat ctctctctgt ctctctctgt tagctacgtt tttcactcgt  
47521 taatgaagtc aagggaactcc gatttcaagga atcgaatacgt tctgattgggt tcagttttaa  
47581 tcttagtttc actcaggtgg ctctcgattc gatttacaat tgcacttct tctggttgaa  
47641 gaattctact taattctcgt ttgaaacggt tctatgcttg gttaagttaat gcactaaaac  
47701 aattctctgt tagttaggat tattaataat tacgtgtggt aagtttctgt tctttgagct  
47761 agattctgat taggttattt gattttgtgc tgctcagttt tccaagttgg ggccacttca  
47821 gtccactcgt gtatctgatg gttcagtttc tcttgcaaaq gaggtaactgt aaacatttct  
47881 tttgtttcgt tctctctgag ttgtcatggg attggtttag ttgagtgtgt tgcaggagat  
47941 gacttgatgc actgcgcact ctgttttact atacttagtg tgaatacggg aatgttttgg  
48001 ctgattttgt taccagaaat cgtcattttc tcgaggagtt taataaaaat tttagagatga  
48061 gaaggatagc aaggagtttt gtgattcatt tgaggaaatg aaaaaggatg actcgtgtta  
48121 acaagattgt aactcaatgt ctaattcccc atcctagtgt tatcgttcat caaaatccca  
48181 tttctattgt cataataat tcatgaactc tgcactgtat cacttcttga tgccttttgc  
48241 tctcttttgg tcttaggatc agcctgccc aacggaacag ttgtttcagc taataaaagc  
48301 aagttcagtg ataagattga ggtcgtctca gccaaaatgt atttccatta ttagggacaa  
48361 cttttacact agccaaaatg gctacaggat tatgtgagga cgggttaagt tgcgctcttg  
48421 tgcaattgatt accctcagagg aaggataact tttttttgg ttgtgcgaac agcaatttga  
48481 cccaagataga ctcttttctg tcatattact ctcttcaggt acatatcatg ctgcagttcat  
48541 ggagaaccgt tcagattttt ctggccgtgt tctgtctgat gtaggcgtgt ctgagtgaaat  
48601 tttgtcnaatg tttgctgcac ttgtatgcac aaatttgaaga tatgtgggac cttaggtgat  
48661 actaactacta ttgtctgcga ttttgccttc acttttctg acttctcagc ataatgtct  
48721 tctcaattta ggctgggtccc aagcatgtgt acttttctg ggcctcagaa atgtgctgat  
48781 atgcctgtaa accctagttgct ggaaacccat tgcctgtcga gaggatcaca gtaagacctga  
48841 tagtccagtt tctgtttgaa ttttttcat ttatatccat ttctctgatg tcttccaact  
48901 tgacacaatt tgattttgat gtcataaagg gatatattag ttgatttgag ttgctctaga  
48961 agcgcagatg ttgtatctct gaaccaatgt gtacgtgttt cagagacgga tctctcttga  
49021 cagctctctc tcatgaaat cagaattctt cagaaggcaa atattgtct ctgagaaaaat  
49081 aatatggagaa accaccattgc tgattcaaaq ttagtatatga aagaatatct ctctatatct  
49141 atataactctg tctcttaggc acctattgg tcaatgagag gatgctggaa acatatgtta  
49201 tctgtaggga tctgttttct tctcnaaac gaaaatggt tccagcaggtt ggaaggtaaa  
49261 gctgactctc gatagtttta cttttctat cttataatgt ctcttcttc atgttaactt  
49321 tctcactgat gatccacatg gcaacttttc cagatgaatt ctattttgtt gaaatggcaa  
49381 ataaggtaca taagtcacat ctatcagga agatatgttt gagattgctt tgaacataaa  
49441 ataaaatggt tgaattaaat atgggatggt taggctttca gtaacttagg cttttctacc  
49501 tctctatgca ggtctttgtt ttgcaacac agaaactata tggagtgtgat ttgacacctc  
49561 tataattgat accccaccag ggttaatttt cccaggtaba tgcatttaca accagtgaa  
49621 actcagatatt tcttttctgt tctcactcct tacaattttt catatttact gattttgtag  
49681 cccgttgattt atgcaattga tccgagatga ttgtgtgctc cctcagatgt ccatgtgata  
49741 gatttcaata tgatgacggt aaagctattt cctacattat attcaggttcc accgtgttcc  
49801 ctaaaaatgt aggcacattt tctgtcatgc ttttctatg ttgcagctgc cctgttctag  
49861 gaattggcatg atccaaatct aataactgtc aaattttttt cacaggaaga tgaactttgc  
49921 agatgtgata tcccggtgaa gtctactcgc tctgtgtgta ccgaaattca ttgacttgag  
49981 tctgtgtttg atgttctctt tgacggggagg tttagaccat tttagaccat aagccttcaa aggtatata  
50041 tctgtgtgcta atataaagat aatattcctt tttagaccat tttagaccat aagccttcaa aggtatata  
50101 tacaacacatg atgtctttgta tccaacagaa cgggtgcaaa atgtgtttaca actgcctccg  
50161 gtgcaccaac aaccatttgg taccaaaata gatgtgttct cctcagagat attcactgca  
50221 tggcagggca agagatcact ggttagactc atttgatgct cccagctgctc cagagttaca  
50281 ctataaaact aacactctca ggtcaatttt gctttctgac atctctcct ttgaaatata  
50341 tctctactat tggacttatt tcttgattgg ctttttaaac acagctaaaa tgtggggtcc  
50401 ccgtgcacat caaggtggaa tctctcagac atcatcgtgc aaactcgatc tgaaggaaac  
50461 ctattataga atgtctcagc cacaggttta cctcacagc gaaccaccag cctagtcgca  
50521 agtaagcgtt ttgtattgct tgattataac aatcaggata tgattgtctgt cctgttttaat  
50581 catttggcgc ttttaagcca ttacattctt tgaatttgca ttatttgcaa agtatataac  
50641 agcatttaac atattgtggc attctcttgg tgcagacat acatatacat agttagtact  
50701 tggaaagacg atagtttctca caacaacacg caaacgctca gctctagaaq ttgactagaa  
50761 aatagaacagc ttttttctgt ccatgtattt gtactttgag tctgtctgca accggttagc  
50821 accaaccattt tattttcaga catattctgt gcttttctg tctagtctga cctattttca  
50881 ctacgtatca cccacatggt atttgatcct ctacacgcaa atgaactatt ctattttggg  
50941 cgctaagcca ggttaattga actttctgac aactatgttc tgacactgca attttttaa



## NCBI Sequence Viewer

51001 ttgattttgga gtgtgttttca aatctgcaat agggacctag ataattcata actggagttg  
51061 acttcaattt attgagttta ctgaacacaaa aatgaatcgg agaacgtgca aattgtgcaa  
51121 tggacaaagg aactgttaata gtatagagca aattttaggc taagctgttc acagctgcga  
51181 ctgcaggctc aggtggatta gcaacaatga cactgaagca gcgtccacag agagtaggat  
51241 ggtcagaaaa tgatccgacc tggcccaggt aattccagca cctctcacac tttagacccc  
51301 cggtcgtgta caccccaate cacactttgt tctctccctc cacatactct cctgtgtgtt  
51361 gcacactgct tatcatctcc ttctccattg aagacaatac atcaacctgt aattgtgcatc  
51421 cacatcagtt tcatggctat agtctcatag aattacactc agcatgatca aatctcaact  
51481 tcaatcgaaa attgtggata ataattgagg aatttgagat gaatcaccat caccagatgt  
51541 aactaaaaac acaatatatt gtaatttcct catacctatt tcttagggat ttaaaagggc  
51601 ttctgagagt gactctata gatctgatca gtacatgtgt ctactgtctc aaagtactct  
51661 tcttttaaca tctaaacatt tagcatttct cttggggccc aaatactata tctaggataa  
51721 agaaccagac tggattaaaa caaattccgg ttgtgccta atcttatcca cctaatact  
51781 tttacttatc tgttttggga agagcctact tgtagtgtgg cctgaacctt acacagaaaa  
51841 accacgggtc acataaaaaa ggccctactt gttgatggtc accatgattc taatttatct  
51901 taagctataa caggaatatg aattttccag ataaggagat ttagtgtctta cttgtgtagt  
51961 tatgaattat cgttgcagag tgtcagcttc gtcttttgcct tcaactatct ctgatttaatt  
52021 tgcagccagt cctgcatctg cagtatgcaa atacaccttt gcttctaaac tggaaaccaat  
52081 cattttctca ttaactgtcta gctccagcac ttgtttccac tcagtctcca actgtcacac  
52141 aaacaccata agtatttcta agatcctcat tacaacaaagc taagcccaagt taagggcagt  
52201 aaaaacaaag aataacgtaa aagtctctcc agagaiaatt cgataatgcc atatcataag  
52261 aactgattac acgtaccact tgacataaca cgagttgtaa tctgtgata ttgtgtaaag  
52321 ttctatagtg ttcatctcca ctaagtactc aaggaaactct atactcttaa ttacaagcag  
52381 agtagagtag atacaatggg ctgaattgtc gaggttttcc gaattgattt agggagagtt  
52441 tgatacctcg agagcccttt gccagaagag aacatcttca gcaggaaatt aaagccattg  
52501 ttcattcaaa gtggccattt taagttcgaa gacaaattct gctgcagagc catcttcaat  
52561 tctgtactca aatggaaagt tctgccagac atcttccgct aggtggagga cgtattggagc  
52621 aatcactctc aattgggata aaagatgtgt tgaangaact gtttgagcgc ttctctcgtt  
52681 aaaaacttga gtaaccttga gaaaaaagat ttttggcca gattacgttg aatagccagc  
52741 acgtattaa agcagggtcta cataggttct agcaaatatc taactcaaga agatgagaaa  
52801 cttaccagat atacagtcta tctttagcta tatcgaagta gaaatttgac agatcacaca  
52861 ttgtgaattt ttgtatgata tgaataaaag aaagctcatt tagtaaacaca gatgatctag  
52921 aacatgaaat taagcctcat gaggttcaaa ggttccatt tgc aaatacatca accctactct  
52981 cacttaagt gtatgatgac caattgctga actaaagttt cttgatgata ttaagagttt  
53041 taactgtata atatacctct gtaagagtga tctatgtctg ttctccgtca caggaaaaatc  
53101 aatacatcac tgaattgatgt accaaaataa gatggagagg ttactctgga tacttttaaag  
53161 aactggtagt tctcgttaaca ttctttgata tcttttcaaa catcttcaag ctgaaagaga  
53221 gcattgctgt caatgatagg taattcctgg tatgggaacc cattatacac ctacaacaca  
53281 acacaagtta attatttaaca gattgtcacc aaccccagaa atacaacaga aagcaagaag  
53341 tcttgacaag tgggaaaaga ggtgaaaaac actactctat taaagagcta agagttagaa  
53401 aacctactct ccaatcatga agatttccca agaggtatct' taagcttccc cttaactctc  
53461 tatatactat agacatttga cgaagaatct ggggacctta tagtacctcc cctgttgaat  
53521 ctacactaga aacccaaga cgcataaact cagctccgta agcaggtgagc tctgttgta  
53581 agaagacaaa ccatgtctag gaaacagatt aataacgaa gatagtctta tacaatggga  
53641 tttacaatgt tgaacaacct cctctctcaa tggccaaagc ttgggtccact tctgttgta  
53701 acgttacctt gagaacttgc cattttcata cctctctcat ccaatacaca tccatgtgtt  
53761 ataacagcag aatagggggc ttctctata tacaatcatg caaatgtgga agaaaataat  
53821 caaaagccac acaagaatag ctaaaaatct gtgaataaat ctcaatttat aaggcaaat  
53881 catggcactc acttctcacc ttgctgtgag atgcttgta acaaaagact ttggaacat  
53941 ccacgatgtc gatctgtacc ttctagatac acgtctgag ggaacttaag accttccgt  
54001 tctcttaata caccagccca agaagatctt ggacatacag attggaaaga aagtgatgta  
54061 aaataggtgt tgggtgctta ggtgtgactat ctgattttaa ttacttctt tctatcagt  
54121 acctgagta caccagagct ccactgtata agtccctct tctaatatct ctgctttatc  
54181 acgataattt tcaaggagta ggtcttccac tgacataac caccacgat cactacctct  
54241 ttgggaaatt atagctgcc acaaaaggtat tcaataaagg ttacagaaaa cacagccagt  
54301 aaatacaaac caataaaaaa caccatactc tcttataggc ctctacattt aacatgtgtt  
54361 attgtctctt cctcatgag tggttctttt gtcttaacat ttactaaagg gactaaagg  
54421 acaccctagt tcttttgtct tgagatgcac catcagagc gatcgaaggt catggcagat  
54481 attctgttta ctgctgaat gacaattgcc aaaaagatac aagtaccaaa attaaacctc  
54541 aaacaatata tctgcatctc ttaaaaaaga aaagctaagg aaagacgta agtaacgac  
54601 tacggatata tctcaacttt gaaaaaactc cacagcaagg tggatcagat catgtggttc  
54661 atgtgctatg aatctcttt cacaagctac acccacat acgactttta gccacatata  
54721 tgaagaaaaa aaatcaact tataaactca atcaggtatc atcaggtgtt ctgattggg  
54781 acccattrta catgtgtgat tgcattcatt gtggctgtgc gaaaccttc aaccgatgca

## NCBI Sequence Viewer

54841 aaccactgctc cagtegetctc aaatategtc ggtttctttag ttcgccaatc atatggggtat  
54901 ttatgagctg catgaacaaga aggaatattaa atcatgagac tctatagcaa atgtttctctc  
54961 aaatgggttcc tcagaagaaga ttggaatcaa actcaccata agattcttcc atgaccagag  
55021 acatatttttc atccaggttaa ctgacaacag cggattctcc ttctccaagg acagagagcc  
55081 ctctaaactg tccagctttct tcagtgaact ttccttcact atccaccgga gagacaagag  
55141 gcagcccgta cttccagccca gtgtcataat cctcctgacc ataccacggy gcagcttgga  
55201 caaacctcgt tccagattctc gtatgttatgt ataacccccc tatagcagctc ggacagcttc  
55261 tgtttgcaat gggatgagtg taacctacaac catgatgaca caccataaac taaggggttc  
55321 aatttatagc attacatgca aaactggaaat tctctttatt ctctaactta aaatgtaaccc  
55381 tgcagtttttc aagatctgaa ccaagggaatg ttttacttat gctaagcttc acaccccatt  
55441 tgcgtcttaa tgcgtggaca aggtcgtctg ctacaatcac gaaaagcttt tgttgattct  
55501 ttagaacctt cccgggaatt ttctttttat tgcttgtcac tgttgatca tcttctgaga  
55561 atgactgcac ttccacaaca gagtactgaa cgtttgcatt caccgcgaca gctgaaagt  
55621 agacaagaaa gaatagaaga agagtacta aatgtgctta tatagcattc aaaaaggtta  
55681 cgttatattg cttctaaagt gagaagagat atgagaaga accgatacaa atgaaagttag  
55741 gatgcaattt gcacgtctgag agaaacacat atactaatat atactaatat accctttac cgtacttgcc  
55801 tggcatagtc caaggtgttag ttgtccatcat tgccaagtat atattaggga taaactcttc  
55861 taaaaggctt gttttcgctc tttaaaaata gcataatgc ttttgaaat  
55921 atgacctctc ggaatactaca cataatggca gatcatgttg ggactgatat cagatacaaa  
55981 ggttaagtgc ttgaattgtca aaatattatt tcatgaaagc ggcatcgctt gagaacagca  
56041 atagaaaaact acatataact agggttctatt ggtgatcat aaagaattag tcttgcata  
56101 atgacaacaa tcagtcacatc tggtaaaatg tcacagggaag ccatgaagag aatagaaga  
56161 ctgttatgac tttagcgagc ttaagttttg cttgaaactg gccgtaatat tattattatt  
56221 cttagattat tgaattcaaat ttacctctag ttcagctctc atcaagcagc tctcgagatt  
56281 agggctccag tgaactggtt tctacctctc ataaatgtac cctttcaagg ccatctggcc  
56341 aaataactca atctacaata tatgataaaa atatatgttg actttatcag gctttgaaag  
56401 actctataaa tcaatctcta caccggttatt tcaatcagta aataagtgtg gacgtgagac  
56461 cattacctga atgctcttcac atctcgggac aaaggtttga taaggattgt tccagctctc  
56521 ccatactcca aatcgtctga aacaccaaga ataattgcat atgtgtcata tgtcaattga  
56581 ataccaaacg aaatactctt cttaattatt atgtcctgag agataataaga atcattatga  
56641 cggttgcatc caaaaacaaga aatcagatat atactaatat cttgatagcc accttaaatg  
56701 attccatttt tgttttgact gttgcttttg caaatttcgc cgcctttgcc ctttaattta  
56761 atgggtgaag ttctctctct acttctctgat ccagggaactg cagaactgaa acgtttgaag  
56821 gttacaccaa attacacagg caagttaagta actggaataa acaaaaagca cagttacttt  
56881 ctgtttctaa tgaactcaag gtccataacc ttattgtgtg ttactcttca actcaattgg  
56941 aaggccatgg .caatccacc caggaaactga ctgaaactta tagttttgga gcagctgcat  
57001 gaagaagcag cgaatttgat gagaaggaac tccactattg gcagtaagga aaataaagaa  
57061 tccctaaaga gagttaattg atgtccaagt aggaagaaga atactactat agcgtattat  
57121 gatactctc taagagcatg acccatatgc agatcaccat tggcataagg accacagctt  
57181 aggcacatca tgaagaatga aaactccctg aaattctaac taactatcagt acaccagctt  
57241 ctcaaaagca agttggcata agataatttt gagaactcca ctccattgtt attatcagaa  
57301 actctcttga agacctgggt ctcttccac aaactcttga gttcaggttc ccttctcaaa  
57361 gaattcgctc tcataccaaa accagctctt ggcataatcaa cggtttctt gtaacttcaa  
57421 tcttctgtct tttcccttaa ccaacacaca acggttaagt atgtgccact tgcataatag  
57481 tattaagaaga gaagtacaag caatgtgatg caaaccttca gattgttct tggctgccat  
57541 aacagcgcca cgaagacctc gttttgatga atgaccaaac tcaattgttg ttgcagttga  
57601 aaaccgtgag acatataaga atgaaaagc cttgacggga gaacttctct gttgacgaaa  
57661 agaatccaac ggcgtgttct tctttaaatt tgaacaactt ttacctgaca aaacctaaac  
57721 ggggacaaac tacactctaa aaactatttc caccgaattt ttacgcaaga cacaataact  
57781 tctatcagat tgaatagggt ggttaagaagta gttctccaaa acaagttctc taggcacaac  
57841 attttttcaa gaaagaaat agagcataga aacagactta gtttgggtta taagggcttt  
57901 aagcatatca gaggacacat atgagctcgt catgtgatgc tcaagatgtc agaactcaag  
57961 taaactactt gcaatttcac tcttcaagga taaagcaact ggataaaagt caaccaagaa  
58021 gcctctgaaa tttttcgaaa caactgggtg gcaatagtaa accttaagaa atctatgaa  
58081 aacgaagaag gaactacttt aacgaaccca atgtccaagt atgtgcaagt catgtataat  
58141 aagaacaaat tctgcacaaa agtggttcta ggaatcgaa ttgatagaaa agactatcac  
58201 tttcccagc acacaaaaca aagagtcaac agagcagaga tggattcaaa gtaatcaat  
58261 ttacattcac atactctgta agatgaagat tgaaccatcg ccaattgact agcttctctg  
58321 gggatttccc ggaaggtatt gaagaaagac gacatctgtc tcaagtctt cgtgtcgt  
58381 tctatctatc agactcatt ttcacgggca acaaaagagt gagaagagat caagaagact  
58441 aacgtcacca aattcaggga agcgtatgtg tgtatgtgtg tggataactg aaaagttaa  
58501 actaaaaaca aacctcaga ctcggtttca attttggat ttatccgtt aaactccata  
58561 tcttctgtgt tttttcttc aagctcataa tagtcaaaag taatcagctt taatacata  
58621 aataactcaa caagttaaaa gaggcgaaat tcttttaaca tgtatggtta atagattaaa

## NCBI Sequence Viewer

```

58681 tataaacaacaa aacaaaaaca accattagat aaggcgggtg aatttatagc catatatcat
58741 cagtaaatgca agatataaca gttagactca acagtagcat caccagtcct catatcgaa
58801 aacttggtcc gtgcttcgac ataacctcta gcaaacccga acattccact gccctccaac
58861 accggcattt cccataacct ggacatcacc gtgttcgacg caagaatcgt tatggctact
58921 ccgttgttact tccagtcctt aaaaagccaa gtcatcgcca tcaacaaccc tatctccctt
58981 tgggctgctc caacgtacat ccttgggctc tgaccacca cagtgcagtt aatggggaca
59041 tctaattgta aagcgtgtgc tatcatgggtg attgatccga aatatgagga acctgagtat
59101 tttgcaaccg gcccttggat cctgatggaa cttgggttcc gaccagttac aatgtcgtgc
59161 caatagacac ggagatgtgt gagttctctt ttttgccga ggccggaggtg ttttcggttt
59221 attgttcttg cgaagtcttc accgtttttt ccggctaagg cggttgcggt gaggaggagg
59281 attgtgtcgg tgaggatgag aatgagcttt gacatgttag ttttctgggt tcgtacttta
59341 attgttttat ttgttgttga tagaatgata gtttatgtaa atttgggggt tatactactgc
59401 aactgtgcaa cgtgctatat atatcgagta atagatgaga tcacgagtggt gaaaattaaa
59461 aaggggaaat agataaacga gtgtctacat atagaagaag acgtgctatt tgataaaatct
59521 tctagaaaat aaagtgaatt tggattattt gattttghta ggtgaataaa gtgtaaaata
59581 ttgtatttta tgggggaagt tacaaaagat tcaatcaata tcagtttcta aaagtttgat
59641 tggcaaaagta ttagatgag tatagctagg tgagcatttg taagtataat gacttctcta
59701 agctttgatt tcagattctt tgttgggtgt ttaattctct aaaaacaaaa caaatattat
59761 atcccttttt taattaaatc tcacgccttta cctttttatt tcatgcaaaa gatgaaataa
59821 ttgaaacttc tcacatttgg tttgttttat catataaaac caaacgaata ataaaacac
59881 ctgtgtcata accctaatag accgatataa tcgggaaggg aaaaacaaac taactattta
59941 aatgttctct taattgaaaag tattgaaact gacctgtttt agttggaaac atatgcctcg
60001 cgtgaatatc ggtccagaca acaacgctca aatagatgc actactctac aatcacaaag
60061 caaatcaaga aagctaaaga aaatcaaat ggctagtggg agtgagcgtg tgggtgttgt
60121 cggagttagt gatccagccc atagctacca tgcactcgag acggtccttg atctcttttt
60181 catacctttc aaggccaaac cacgattcaa acttgtctc ctacacgctc gccaccaacg
60241 gaccttcttc ctccgtgttg ctggcccggt aacggctgat attataccaa tgggtggaag
60301 agatttaaac aagaccgctg acttgtttaa gaagaagtgc gcggaagtgt gctcggctaa
60361 gtccgttgag atctcgtctt tggaggtgat tgaaggagat ccggaagaac taatgttggg
60421 agccgttgag agacatcatg catgagttct gtctcttggt agccatggat atggagctga
60481 cgtcaagag agtgttttgt ggaagcgtcaa gtgtgcgtaa ctatttagct cattaagcac
60541 attctccgt tcgtatcate aagaagccta aggataaac tacttctacc aataaacagc
60601 aatgaggaaa aagacgagct gattttcgta tatgttacca tctgttttcc tataagtaaga
60661 cccattctat cgaacattgt gcataactct ctccccaac attctccatt tcaacgttaa
60721 tagcttagct ctaatgtata tactcatgtt atatattcac tatattaatc gccctcttct
60781 tgtgttatta atctccttat ttttttctgt atatattgca taataattgaa gtctcttagaa
60841 catttttaca ttattcttat atgatacata ttgaaagggt ccaatataat tttgtctttg
60901 gaaacttgta gtaccttgca aagagcttga tcatgtacca agctcgaggtg gaaattaaac
60961 ctatctgta gtactcata accaatgttg ccgagaactc gtaagcaatg gctctagtaa
61021 ctctcttttc ttttctcagt ttacctttac caactttgtg gtcttgtttg cattaataagg
61081 ttgtatccga gctgttagaa gaagcccaact gtgttttatg gcgaagacat ttaaaactcc
61141 tataaccaaa taatcatggt gctgactgct gaggacttgt caatttatga atatacaatc tatgttaacat
61201 tttttcttc ttttctcagt ttacctctac cattttatga atatacaatc gatcttaacat
61261 ttccaatccg agcttgtgta agataaacat tcttcaaaact tcttcaaaact gattcaaacat
61321 tgttgtattc ctcttaagca atattttcaa acagactcca aaagatcgga taacatacat ctctccaaat
61381 aacctccaaa acataaatat cacagactcca aaagatcgga taacatacat ctctccaaat
61441 aatgactgct tactctcata ctctccagca gtctccaaat gttctcaatt tctctttcca tctccaaat
61501 atggacgctc caatcggtag caccacagcc accggttcga tgcctcaaaa ccggtgcac
61561 cgtctccaat gcaaaagagt gtcttccaca aattactgga ccgcttcgag aaagagagac
61621 acaatgccac aactaaaaat tgaactgaaa attgtagggc tttaccccaa agggagccct
61681 tccagaattt ctcgaagcc caagacaagc tt

```

11

FIGURE 15


**A****B**

FIGURE 16



## FIGURE 17

RCH1.2 AB028621+AP002037 (BAB03091)

mslshliffssssslsffifficfslsdaeqnpeasitlywhssstpslsslfnwnsidntpcnwnftitcssqgfitdiesv  
 plqislpknlpafsrslqktisganitgtlpeslgdcglkvldssnglvgdipwslsklmletilnsnqltkgkipdiskcsklksliif  
 dnlltgsiptelgksglevirngnkeisgcipseigdcnsntvlglaeatsvsgnlpslsgkklkdetisiytmisgeipsdlgnce  
 lvdfilyenslsgsipeigqitkleqiflwqnsllvggipeeigncsnlkmidslnlsgspsisgrtslfeefmisdnkfsipsipiti  
 snccsslvqlqldknqisglpselgltitklitffawsnqlsgsippgladctdlqaldlsmsltgtipsgifmlmtkllinslsgtipq  
 eigncsslvrlrfgnritgeipsigsglkkinfldfssnrllhgvkvpdeigscselqmidlssnslsgslpnpvsslsqglvdivsan  
 qfsgkipasgrlvslnklisnlfsgsipsitlsgmcsglqldgsnslsgsipselgdielalnlssnrlltgpikpskiaslnklsil  
 dshnmlegdlepianienlvslnisynsfgyipdnklfrqlspqdegnkkkcsstqdcslfityrkngnlgddgdasrtrklritl  
 allititvmlilgavavirarnmidnerdseletykwqftpfqklnfsvdqirclvepnvigkcsqsvvyradvndgeviavkkl  
 wpamvngghdektknvrdsfseavltgtirhknivrlfgocwnmrtrlmydympngslsgslherrgssidwdlryillgaa  
 qglaylthdclppivhrdikannilgldfepiyadiqlaklvdegdlgrcsntvagsygyiapeygyismkitekssdvysygvv  
 vlevitgkpldcpvgiehlvdwvnrngslsvldslsrteaeademmvgltalovnssdpderptmkdvaamleikq  
 ereeyakvdlilkksppptttmqeocrknermmipaaaassskemrreerllksnntsfassylsssssl

RCH1.3 AB011476 (BAB09286)

mpnrprcfiffllfhsllfscpsideqglallswksqnlsgdalsswkasesnpcqvwgikcnerngqvseiqqlvmdfqq  
 plpatnrlqksltsltslntgtspikelgdlselevidladnsisgelpvdfiklklkilslnnnlegvipselgnlvieltfdnkl  
 ageiprtgelknlleifraggnknrlrgelpweigncesvltglaeatslgrpasignklkvqtaiyltslsgpdpdelgncetqln  
 ylyqnsisgspvsmrglkkqslllwqnnlvgkipitelgtcpelflvidlsenltgniprsfgnlqlgelqsvnqlsgtippealanc  
 tkthleidnnqisgelppligkltstlmffawqnqltgiipesisqcelqaidlsynnlsgsipngifglefvdlnhsgnltgpgtp  
 kslqfildsdnsitlsglptglsiteltklaknrfsgeipreisscslqlnlngdngftgelpnelgripalsaislnscnhtgelpsrf  
 sslntgltdvshnklagnlnvladlqnvlslnisfnfsgelpntlfrklplsvlesnkgflstrpengiqtrhsavkvtmslvaa  
 svvlvmavylvtkaqritgkqeeldswevtlyqkldfsiddivknitsanvigtgssgvvyrvitpsgettlavkkmwkskeerna  
 fnseintlgsirhmriilrgwscsnrnklifdytlpngslssllhagkgsggadwearydvvlgvahalaylhhdclppilhgdv  
 kamnnvlgsrfeysladfqlakivsggevgtdgdsskslnrplagsygymapehasmqhitekssdvysygvvlevitgk  
 pldpdpdgahlvqvwrdhlagkkdprieldprlrgadpimhemqltlavsficvsnkasdrpmmkdivamleikqrdm  
 drsesdmikggkcekwwqpplpekevstprgssncsfaysdesv

RCH1.4 AL022223 (T05050 or CAA18216)

mppniyrslfssllcfffpcsfldqggqallswksqnlsgdafsswhvadstpcnvwgvcnrrgeveiqklgmldqgsl  
 pvtlsrslksltstlsslntgvpikegdfteileldsdnsisgdipveifrlkkkltstlnnnleghipmeignslsgivelmfdnkl  
 geipsigeklnqlvraggnknrlrgelpweigncenlvmglaeatslgrkvqtaiyltslsgpdpdelgycetqln  
 lylyqnsisgspitpiggkklqslllwqnnlvgkipitelgncpelwidfsenltgtiprsfgklenlqelqsvnqlsgtippeelntk  
 klthleidnnltgeipslmsnlrstmffawqnkltnipqslsqcelqaidlsynslsgsipkeifglefildhntslsgslgtitp  
 skldfidsnaisslppgigltitklaknrfsgeipreistrqilnlgendfsgelidpdpqslaislnscnrlgeipersf  
 dklnglvdshnqltgnlntvtdlqnvlslnisyndfsgdlnptntfrlplsdlasnrlylsnaistrpdpntssvvrtlilivv  
 avtlvmavylvtraraagkqlgeeldswevtlyqkldfsiddivknitsanvigtgssgvvyrvitpsgeslavkkmwkskeesg  
 afnselktlgsirhmriilrgwscsnrnklifdytlpngslssllhagkgcvdwearydvvlgvahalaylhhdclptilhgdv  
 kamnnvlgsrfeypylcdfglartiesgypntgidlakpnrppmagslwhgssdfdlfcllgtetehasmqriteksdvysygvv  
 llevitgkpldcpdgahlvqvwrdhlaekkdpsrlldprldgtdsimhemqltlavsficvsnkanerplmkdvamltle  
 irhidvgrsetekikagcgskpqqfmsnekinshgssncsfafsdsv

RCH1.5 AC015446 (AAG12526)

merersnffiffcswvsmacptlssissdggallsikrpspslsswdpqqdqtpcswygitcsadnrvisvsipdttlnlssipdl  
sslsisqilnlssnlsngpippsfgklthirlclssnslsgpipselgrlstlqfiilnanklsgsipsqisnlfalqvclqdnllngsips  
sfgsivslqqfrlggntnlggpipaqglgiknltldgfaasglsgsipstfgnlvnlqtlalydteisgtippqglgclselmlylhmnklt  
gsipkelgklqkltstllwgnslsgvippeisnccslvvdvsandltgdipgdlgkdvwleqlqsdnmftgqipwelsnccslla  
lqldknklsgsipsqjgnlkslqsfllwensisgtipssfgnctdlvaldlsmkltgripeelfslklrsklllgnslsggplkpsvakcq  
slvrlrvgenqlsgqipkeigelqnlvldlymnhfsgglpyeisnitvielldvhnnyitgdipaqlgnhneqldlsmstgnipl  
sfgnlsyinkliinnlilgtqpkssiknlqkltlldsynslsgeipqelgqvtsitindlsyntftgnipetfsdtlqlqslssnslhgd  
kvlgstlsaslniscnnfsgpipstpfkltisttsylqntlnichslsgitcsshtgqnnvgkspkivaltavilasitiaawliilrnn  
hlyktsqnssspstaedfsypwtfipfqkigivnnivstldenvigkgcsgivvykacipngdivavkklwktkdnneeges  
tidsfaaeiqilgnirhrnivklgycsnksvkllynyfpgnlqqllqgnrldwetrykiaigaagglaylhhdvcpailhrdvk  
cnnilldskyeailadfglakimmnsprnyhnamsvaeygymnitekssdvysygvvleilsgrsavepqigdgdlhiveww  
kkkmgtfepaivldvklqglpdlqivqemlqlglgiamicvnpspverptmkevvtllmevkcspeewgktsqpllpss

crop01lpct.ST25.txt  
SEQUENCE LISTING

<110> UNIVERSITEIT UTRECHT

<120> A novel root specific promoter driving the expression of a novel LRR recepto  
r-like  
kinase

<130> CROP-011-PCT

<150> EP 00870288.8

<151> 2000-12-04

<150> US 60/256,204

<151> 2000-12-15

<160> 19

<170> PatentIn version 3.1

<210> 1

<211> 7438

<212> DNA

<213> Arabidopsis thaliana

<400> 1

ctcgagtttc agatgtttct attaaataag taatgaaagc agctctaatt ctcccaaat	60
tcacaagtat agaaagatag aaagatagaa acagagttat tgcattggaag ctgaaaatca	120
tacagcatgg tggatataac caatgagaca caaacaatc acgcatgtg ttatagaaca	180
acacttatca tctgtattaa cataaagcaa aagttcctat acgacatagt ccaaagttac	240
agaccataag cacaccttat gatttgttgt accctctaaa acacaggatg atgacaagaa	300
gcaaatttgc ccaaattttac tgagggaaaa acgaggggctt aacgctctca tatctcaaac	360
aaaaactttc caaacagttt catgtgttca gattggaggg caatggcaac ggataagctg	420
ccatcattag taggacttgg tagcacaaaa gacaaacctt cgtatgggaat tccaccagg	480
cccataaaga taggacgacc ccaaccaaaag tctgcatcat aaataggtaa tctaaccag	540
cttgtgatcc ccaaatttgg gcactttagt gtatgtgcac cgcggacaag ggtgtgacaga	600
tcaggctgca tctccaggta gtcaagagct gacctcagat agttatcatc catacgaacc	660
aagaaatcat gaatctgtcc tgcagcatac catgttggtt tagataacaa atccccctgca	720
acagccaatg gtgttgcagt gaatatcaca ttcccaagt aaccaggagg cagctgcgga	780
cgtagtctag accttcate agttgcaatg tacagtttct tctcttggtc gtttggaaac	840
cctgcgcctt ttccactga tctccacaca tgccttgcca acatctcgta tgagctgtag	900
ctgacagtgt tccatctctc ctgtgatttc gcttaagag caacaagctg gtctcgtgtt	960
aatttgaata tagagacagt ggtattctca ggtcctgatt tagacggatc aagaggtatc	1020
ttcatacttg gtgcaggctg atattcaaca tgatgaaaag caggctgtgg cgggtcccta	1080
gctcggagga gtgttcgac aatgaaaggt ggaatggtt ggtcaagacc acgagccata	1140



## crop011pct.ST25.txt

tcagaccatg	tgttgataaa	atgaagacca	gagaaccat	ctgccgctg	atgttgcac	1200
ccaaccccaa	gtgaagctcc	cccacattta	aagaaagtca	ccatatcat	caacaatat	1260
cattacataa	ctaaaaacag	aacatacatc	atcaaagaat	atcattacag	aactaaaaaa	1320
cagaacaaca	caagcattca	ataaattaaa	aaagggaaag	actttaaat	aagctgacct	1380
gcaaaacgag	aagcgggaaa	gagtgaatgc	cagcggagt	atcaacttcg	ggaataagct	1440
gacggagatt	aagggttagga	gcaaaatcac	caaaatcatc	gataacagaa	ggagtatcag	1500
ccacaacgaa	gagaacacca	gcacogttac	aatcgatctc	aatacgaacca	tcactgtctc	1560
tcttcaacgc	accagccata	gggtaaaaag	ggacaagggc	tttggaagaa	gcttccttca	1620
tgacctgagg	gtcaaaagaaa	ttggaagcgc	cggtgggtct	gtagaagtag	acactagggg	1680
tatggaatct	ggggatgaca	aggtcgacgt	tggagttcca	aagattagtg	attggtgtct	1740
cggtggcagg	ccggaccatg	gtggaatctc	tgatgttaat	tttcataatt	ttagaaaga	1800
tcaaacgaac	aaacaaat	ttgtttgagt	ttgatctgag	aaattgtgtg	agaagcgagg	1860
aggggtagat	ataagtgggt	gtgaacaaaa	aaggaaaggt	cgtaatcctt	cacctaccgc	1920
ccacgaccc	cgcctactcc	atctgtcgac	caactctctc	agaaggttta	tcattggtcc	1980
atctcatcac	tgggtctctc	ttttctatcc	ggggtttggt	aaccggttcc	ctctgtctca	2040
aggcccaaat	aatactaatg	ggtctaaata	gtataaagtt	cgaatgaaga	agaaggaaaa	2100
aaaattgaca	agttgattgt	tatagctacc	gtacaagtta	gtcaatcgaa	tgccaaagtt	2160
tggcttaact	taactcatta	ccacgtagca	gagcataaca	acaaccacca	ataacatcta	2220
acggcgtaaa	ctcacgtggt	aagagaatta	cgtcagaatc	caagaaagtc	tttttttcag	2280
tacaccacaa	ataaattatg	ttgaatttat	atccgaacta	tactatatta	tatgaaaaca	2340
tacctcaatt	agtcataaat	ttcacatttc	attttaattt	attaactaat	ggttgttggt	2400
gcggatagct	ttctcttgc	ttctgttcta	aaattatttc	ttagccatta	tttgatgac	2460
caacttgggt	tgaggatcta	ataaaaaatc	aatttctttg	aagcattcat	tggcttaatg	2520
atccaaacct	tttgtctctc	cactctctaa	gaaaataaag	caaaagcaaa	actaaaagaa	2580
aagcaactag	taccgtgggt	tggtcaatgg	taaccocatgc	taccaaatgc	caatcaaagc	2640
ttaaatgaca	atttagtacc	ttgggttggt	catgatttag	agcgggaaca	atataccata	2700
catcaaacga	ggatatacag	agaaaaattca	tgggaagtatg	gaatttagag	gacaatttct	2760
cttctgggct	acaacggacc	ggcccatctg	ctcatttacc	cagaggtatc	gagtttgtgg	2820
acttttgatg	ccgctagaga	ctattggcat	cggattgaaa	aaaaagttaa	cttcgttggt	2880
aacaattttc	tgaatgcaat	attttccttg	tcatgaatat	ttaaacttgt	tattactttc	2940
ttttagctta	gggtgtggaca	attatggagt	ttacttcaaa	cgaggaagaa	tcttaaacgc	3000

crop011pct.ST25.txt

tcgggttcagg tctcgaaaac aaaccaactc acaatctcga cttaattgag gaaaacaatg	3060
caaaaccaca tgcgatcttc catatttcta tcataatctt ataagaaaaa acactactaa	3120
gtgaaatgat tctgtatata tataaccaat gccttttgtt ttgtgatatt ttatgtatat	3180
ataactattg actttttgtc tctatggata gtgtctcggg ctcttggcaa acataattca	3240
aagaaaagtt aatgactgta attaatatt ctgaagctag aacagaaacc ccgaggtaaa	3300
agaaaaagac agagcacatg aagtttagta cttttatata tttaatatat cattctttct	3360
tattgcttat ctctaaagca aaaacttccc taaacctaa gccaaaggac tcagatcgat	3420
gcagaaccaa gaaggcttgt tttggatttg agagccaaat gcaagaaaaa aaactcttat	3480
gtctttgcaa atgccgattc caagaaaaaa agccttaact gtttctcatt ttccattac	3540
attgtctctg tttttagctt tcttcacttc ctgcacctct gcaccaacca atgaagtctc	3600
agctttgatt tcttggcttc acagctcgaa ctgcccgcca ccgtcagtct tctccggctg	3660
gaatccttct gattctgacc catgtcaatg gccttacatt acttgttctt cctcagacaa	3720
caaacctggt acagagatca atgtcgtctc tgttcagtta gctttacett tccctcctaa	3780
catttctctg tttacttcac ttcagaaact cgttatctcc aacactaatc tcaccggagc	3840
tatatcttct gagatcggag attgttcgga gcttatagtt atcgatttaa gctcaaatag	3900
tcttgttgtt gaaataacct cgagctcagg gaagctcaag aatcttcaag aactttgttt	3960
aaactccaat ggctcacag gaaagatccc accggaactc ggcgactgcg tttagctcaa	4020
gaatcttgag attttcgata actacttata cgagaatctc ccgttgaggc tcggaaagat	4080
ctcgactctt gagagcataa gagccggagg aaactcagag ctttcaggga agatcccgga	4140
ggagatcggg aactgtagga atctcaaggt cttaggggta gcagcaacga aaatatccgg	4200
ttctttacct gtttcgttgg gtcaactaag caagctccag tctttatctg tctattctac	4260
aatgctctca ggtgagatcc ctaaagagct tggaaactgc tctgaactta tcaatctgtt	4320
tctatacgac aatgacttgt ccggcacgct tccaaaagaa ctaggaaagc ttcaaaacct	4380
agagaagatg cttctatggc agaataatct ccacggacct atccctgagg agattggatt	4440
catgaaaagc ttaaacgcca ttgatctctc tatgaactat ttctcaggaa ccattccctaa	4500
atcgttcggt aacttgtcga atctccaaga gcttatgctt agcagcaaca acatcaccgg	4560
gtcaataacct tcgattctaa gtaactgcac aaagcttggt cagttccaga ttgacgccaa	4620
tcagatttca ggtttgattc caccagagat tggattgctt aaggagctca acatcttctt	4680
gggatggcag aataagctag aagggaatat tccggacgag tttagtgggt gtcagaatct	4740
tcaagctctt gacttgtcac agaattatct caccggatca ttacctcgag gtttgttca	4800
gcttcgtaat ctgactaage tcttgcttat atctaatgac atctccgggt ttatccact	4860
ggagatcggg aactgcactt cgcttgctcag attaaggctt gtcaataaca gaatcaccgg	4920

## crop011pet.ST25.txt

agaaatccct	aaaggatttg	gattttcttca	gaaccttagc	ttctctgact	tgtctgagaa	4980
caatctctcc	ggccagttc	ctttggagat	aagtaattgt	agacaactcc	agatgettaa	5040
cttgagcaac	aacactcttc	aaggttatct	tcctctgtct	ttatcatcgt	taacaaagct	5100
tcaagtactt	gatgtctctt	ccaatgactt	gactggcaag	atacctgata	gtctaggcca	5160
ttctatttca	ctcaaccggc	tcatttcaag	taagaattct	ttcaatggag	aaatcccttc	5220
gtctctcggt	cactgcacga	atcttcagct	tcttgatctc	agcagtaata	acatctctgg	5280
aactataccg	gaggagcttt	ttgacatcca	agatctagac	attgcgttaa	acttgagctg	5340
gaattcatta	gatggcttta	tcccggaaag	gatttccgcg	cttaaccgat	tatccgtgct	5400
cgatatttcg	cataacatgc	tttcaggcga	cctttctgcg	ctgtccggtc	tagaaaactt	5460
ggtttctctg	aacatctctc	ataacagatt	ctcaggttat	cttcagata	gtaagtggtt	5520
tagacagctg	ataggagcag	agatggaagg	aaacaatgga	ctctgttcca	aaggtttcag	5580
gtcttgcttt	gtaagtaaca	gttcacagtt	aactacacag	cggtggtgtc	actcacatag	5640
actcaggata	gccattggat	tgctaatacg	cgtgacagcg	gttctagcgg	tattaggcgt	5700
gttgccgggt	atacagagcta	aacaaatgat	tcgagatgat	aacgattcag	agactggaga	5760
aaatctatgg	acatggcaat	tcacaccttt	tcagaaactc	aacttcacag	tcgaacatgt	5820
actcaagtgt	ttggtagaag	gtaatgttat	agggaaagggt	tgctccggga	tagtgtacaa	5880
agctgaaatg	cctaacagag	aagtcacgcg	ggtgaaaaag	ctctggccag	tgacagtacc	5940
taatctgaat	gagaaaaacta	agtcacacag	agttccagac	tcattctcag	ctgaagtaaa	6000
aacacttgga	tcgatcagac	acaagaacat	tgtacgcttc	ttgggatggt	gttggaacaa	6060
gaacactaga	cttcttatgt	atgatttat	gtcaaattgg	agtttgggaa	gtttgcttca	6120
cgaaaggagc	ggtgtatgta	gcttaggatg	ggaagtgagg	tacaagatta	tacttggtgc	6180
agctcaggggt	ttggcttact	tgacacctga	ctgtgttcct	cccattgttc	atagagacat	6240
caaggcaaat	aatattctga	ttggccctga	ttttgaacct	tacattggag	atttcggact	6300
cgctaagctt	gttgatgatg	gcgactttgc	tcgttcttcc	aacaccattg	ctggttccota	6360
tggttacata	gtccaggtga	cttcccaaat	ttctgttttg	ctgctaatta	gttaacttga	6420
tacatcataa	gttactagta	gtcttagcct	agtaatggta	gaaattcagg	tttatagttt	6480
ggttttgggt	tttggcattt	tgaaaagaat	agtattttgg	ttcagttgat	tttcgatttg	6540
gtttaaaatt	ttagaaaaaa	aatggtagaa	gaaataattt	ttgattaata	ttatatcatc	6600
tactgtatac	gatttataaa	aaaaattgat	ttggtttgggt	ctcgatatatt	tgcttgtagc	6660
ttggttcaat	atttttacta	tatatatgat	tcatgacttt	catactaate	aatattttgtt	6720
cggtttcggt	ttgatttagtc	tgaaagtga	tcatatagtt	aaaatattga	ccaatctatc	6780

crop011pct.ST25.txt

atatacttac acgctacaag attttcga	taattttggt gtaagcattt	tggtttttat	6840
atgttcatcc ttagtaacta acaacaata	tcaattggga agcataacgt	tatatattcc	6900
cataaggcta taacactata attatgcaga	atacggatac tcaatgaaga	taacagagaa	6960
aagcgacgtg tacagctatg gagtcgtagt	gctcgaggta ctaacgggta	agcaaccaat	7020
cgatccaacg ataccagatg gactccacat	agtggactgg gtcaagaaaa	tcagagacat	7080
acaagtaacg gacc'aaggac tacaagcaag	accagagtca gaggttgaag	agatgatgca	7140
aacgctagga gtgcgccttt tatgcattaa	tccaataccc gaagacaggc	ctacaatgaa	7200
agatgtggct gctatgctta gtgagatatg	tcaagagaga gaggaatcga	tgaaagttga	7260
tggttgctcg ggaagttgta acaatggaag	agaacgtggc aaagatgatt	cgacttcac	7320
ggttatgcaa caaacggcta agtatttgag	aagtagtagc acgagtttct	ctcgctcttc	7380
tttgctttac tcttctctct cttctgctac	ttctaattgt agaccaaatc	ttaaataa	7438

<210> 2  
 <211> 3499  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 2

ctcgagtttc agatgtttct attaataaag	taatgaaagc agctctaatt	ctcccaaat	60
tcacaagtat agaagatag aaagatagaa	acagagttat tgcattggaag	ctgaaaatca	120
tacagcatgg tggatataac caatgagaca	caaaacaatc acggcatgtg	ttatagaaca	180
acacttatca tctgtattaa cataaagcaa	aagtctctat acgacatagt	ccaaagttac	240
agaccataag cacaccttat gatttgttgt	accctctaaa acacaggatg	atgacaagaa	300
gcaaatttgc ccaaatttac tgagggaaaa	acgagggtt aacgctctca	tatctcaaac	360
aaaaacttct caaacagttt catgtgttca	gattggaggg caatggcaac	ggataagctg	420
ccatcattag taggacttgg tagcacaaaa	gacaaacct cgtatggaat	tcaccagggt	480
cccataaaga taggacgacc ccaaccaag	tctgcatcat aaataggtaa	tctaaccag	540
cttggtgattc ccaaatttgg gcaactgtag	gtatgtgcac cgcggacaag	ggctgacaga	600
tcaggctgca tctccaggta gtcaagagct	gacctcagat agttatcatc	catacgaacc	660
aagaaatcat gaatctgtcc tgcagcatac	catgttggct tagataacaa	atcccctgca	720
acagccaatg gtgttcgagt gaatatcaca	ttcccaagt aaccaggagg	cagctgcgga	780
cgtagtctag accttccatc agttgcaatg	tacagtttgc tctcttggtc	gtttggaagc	840
cctcgcgcct ttcccaetga tctccacaca	tgccctgccca acatctcgtg	tgagctgtag	900
ctgacagtgt tcccatcttc cttggatttc	gccttaagag caacaagctg	gtctcgtggt	960
aatttgaata tagagacagt ggtattctca	ggctctgatt tagacggatc	aagaggatc	1020

crop011pct.ST25.txt

ttcatacttg	gtgcaggctg	atattcaaca	tgatgaaaag	caggctgtgg	cgggctcccta	1080
gtctggagga	gtgttcgac	aatgaaaggt	ggaatgggta	ggtaacagacc	acgagccata	1140
tcagaccatg	tggtgataaa	atgaagacca	gagaaacct	ctgccgcgtg	atgttgcac	1200
ccaaccctaa	gtgaagctcc	cccacattta	aagaaagtca	cctatatcat	caaacaatat	1260
cattacataa	ctaaaaacag	aacatacatc	atcaaagaat	atcattacag	aactaaaaaa	1320
cagaacaaca	caagcattca	ataaattaaa	aaagggaaag	actttaaatt	aagctgacct	1380
gcaaaacgag	aagcgggaaa	gagtgaatgc	cagcggagtg	atcaacttcg	ggaataagct	1440
gacggagatt	aagggtagga	gcaaaatcac	caaaatcatc	gataacagaa	ggagtatcag	1500
ccacaacgaa	gagaacacca	gcaccgttac	aatcgatctc	aatacagcaa	tcatcgcttc	1560
tcttcaagcg	accagccata	gggtaaaaag	ggacaagggc	tttgaaaga	gcttctctca	1620
tgacctgagg	gtcaaaagaa	ttggaagcgc	cggtaggtct	gtagaagtag	acactagggg	1680
tatggaatct	gggatgaca	aggtcgacgt	tggagtcca	aagattagtg	attggtgtct	1740
cgggtggcagg	ccggaccatg	gtggaatctc	tgatgttaat	tttcatattt	taaagaaaga	1800
tcaaacgaac	aaacaaattt	tgttttgagt	ttgatctgag	aaattgtgtg	agaagcgagg	1860
aggggtagat	ataagtgggt	gtgaacaaaa	aaggaaaggt	cgtaatcctt	cacctaccgc	1920
cccacgacc	cgcctactcc	atctgtcgac	caactctctc	agaaggttta	tcattggtcc	1980
atctcatcac	tcggtctctac	ttttctatcc	ggggttggtg	aaccgcgtcc	ctctgtccta	2040
aggcccaaat	aatactaagt	ggcttaata	gtataaagtt	cgaatgaaga	agaaggaaaa	2100
aaaattgaca	agttgattgt	tatagctacc	gtacaagtta	gtcaatcgaa	tgccaaagtt	2160
tggttaact	taactcatta	ccacgtagca	gagcataaca	acaaccacca	ataacatcta	2220
acggcgtaaa	ctcacgtgtt	aagagaatta	cgtcagaatc	caagaaagtc	tttttttcag	2280
tacaccacaa	ataaattatg	ttgaatttat	atccgaacta	tactatatta	tatgaaaaca	2340
tacotcaatt	agtcataaat	ttcacatttc	attttaattt	attaactaat	ggttgtgtgt	2400
gcggatagct	ttcttttgct	ttctgttcta	aaattatttc	ttagccatta	tttgatgac	2460
caacttgggt	tgaggatcta	ataaaaaattc	aatttctttg	aagcattcat	tggcttaagt	2520
atccaaacct	ttgtctctcc	cactctctaa	gaaaataaag	caaaagcaaa	actaaaagaa	2580
aagcaactag	taccgtgggt	tggtcaatgg	taacctatgc	taccaaatatc	caatcaaaagc	2640
ttaaatgaca	atttagtacc	ttgggttggt	catgatttag	agcggaacaa	atataccata	2700
catcaaacga	ggatatacag	agaaaattca	tggaagtatg	gaatttagag	gacaatttct	2760
cttcgtggct	acaacggacc	ggcccatctg	ctcatttacc	cagaggatc	gagtttgtgg	2820
acttttgatg	ccgctagaga	ctattggcat	cggattgaaa	aaaaagttta	cttcgttggt	2880
aacaattttc	tgaatgcaat	attttccttg	tcatgaatat	ttaaacttgt	tattactttc	2940

## crop011pct.ST25.txt

```

ttttttttt gggtgtggaca attatggagt ttacttcaaa cgaggaagaa tcttaaacgc 3000
tcgggttttt tctcgaaaac aaaccaactc acaatcctga cttaattgag gaaaacaatg 3060
cagaacatna tgcattgttc catattttta tcataatctt ataagaaaaa acactactaa 3120
gtgaaatttt tctgtatata tataaccaat gcctttttgt ttgtgatatt ttatgtatat 3180
ataaatattt actttttgtca tctatggata gtgtctcggg ctcttggcaa acatatttca 3240
aaqaattttt aatgactgta attaattaat ctgaagctag aaacagaacc ccgaggtaaa 3300
agaaattttt agagcacatg aagtttagta cttttatata tttaatatat cattctttct 3360
tattttttt ctctaaagca aaaacttccc taaaccttaa gccaaaggag tcagatcgat 3420
gcgaattttt gaagccttgt tttggatttg agagccaaat gcaagaaaaa aaactcttat 3480
gtctttttt aaatgccgatt 3499

```

```

<210> 1
<211> 3499
<212> DNA
<213> Arabidopsis thaliana

```

```

<400> 1
atgtcttttt aaatgccgat tccaagaaaa aaagccttaa ctgtttctca tttttccatt 60
acattttctc tgttttttagc tttcttcate tctcgcacct ctgcatcaac caatgaagtc 120
tcagttttga tttcttggct tcacagctcg aactgcgcgc caccgtcagt cttctccggc 180
tgaaattttt ctgattctga cccatgtcaa tggccttaca ttacttgttc ctctcagac 240
aacaattttt ttacagagat caatgtctgc tctgttcagt tagctttacc tttccctcct 300
aacaattttt cgtttacttc acttcagaaa ctctttatct ccaacactaa tctcaccgga 360
gcataatttt ctgagatcgg agattgttcg gagcttatag ttatcgattt aagctcaaat 420
agcttttttt gtgaaatacc ttcagagtcta ggaagctca agaattctca agaactttgt 480
ttaaattttt atggcctcac aggaagatc ccaccggaac tcggcgactg cgttagcctc 540
aagatttttt agatttttga taactactta tccgagaatc tccggttgga gctcggaaag 600
attttttttt ttgagagcat aagagccgga ggaactcag agctttcagg gaagatcccc 660
gaaatttttt gaaactgtag gaatctcaag gtcttagggt tagcagcaac gaaaatatcc 720
ggtttttttt ttgttttcgtt ggggtcaact agcaagctcc agtctttatc tgtctattct 780
acaaattttt cuggtgagat ccctaaagag ctgggaaact gctctgaact tatcaatctg 840
ttctctattg acaatgactt gtccggcagc ctccaaaaag aactaggaaa gcttcaaaac 900
ctctctttta tgcctctatg gcagaataat ctccacggac ctatccctga ggagattgga 960
ttctctttta gcttaaacgc cattgatctc tctatgaact atttctcagg aaccatccct 1020
aaattttttt gtaacttgtc gaatctccaa gagcttatgc ttagcagcaa caacatcacc 1080

```

## crop011pct.ST25.txt

gggtcaatac	cttcgattct	aagtaactgc	acaaagcttg	ttcagttcca	gattgacgcc	1140
aatcagattt	caggtttgat	tccaccagag	attggattgc	ttaggagct	caacatcttc	1200
ttgggatggc	agaataagct	agaagggaat	attccggacg	agttagctgg	ttgtcagaat	1260
cttcaagctc	ttgacttgtc	acagaattat	ctcaccggat	cattacctgc	aggtttggtt	1320
cagcttcgta	atctgactaa	gctcttgctt	atatctaata	ccatctccgg	tggttatcca	1380
ctggagatcg	ggaactgcac	ttcgcttgtc	agattaaggc	ttgtcaataa	cagaatcacc	1440
ggagaaatcc	ctaaaggat	tggatttctt	cagaacctta	gcttctctga	cttgctcgag	1500
aacaatctct	ccggtccagt	tcttttggag	ataagtaatt	gtagacaact	ccagatgctt	1560
aacttgagca	acaacactct	tcaagggtat	cttctctctg	ctttatcate	gttaacaaag	1620
cttcaagtac	ttgatgtctc	ttccaatgac	ttgactggca	agatacctga	tagtctaggt	1680
catctcattt	cactcaaccg	gctcattcta	agtaagaatt	ctttcaatgg	agaaatccct	1740
tcgtctctcg	gtcactgcac	gaatcttcag	cttcttgatc	tcagcagtaa	taacatctct	1800
ggaactatac	cggaggagct	ttttgacatc	caagatctag	acattgcgtt	aaacttgagc	1860
tggaattcat	tagatggctt	tatcccggaa	aggatttccg	cgcttaaccg	attatccgtg	1920
ctcgatattt	cgcataacat	gctttcaggc	gacctttctg	cgctgtccgg	tctagaaaac	1980
ttggtttctc	tgaacatctc	tcataacaga	ttctcagggt	atcttcacaga	tagtaaagtg	2040
tttagacagc	tgataggagc	agagatggaa	ggaacaatg	gactctgttc	caaagggttc	2100
aggtcttgct	ttgtaagtaa	cagttcaccg	ttactacac	agcgtgggtg	gcactcacat	2160
agactcagga	tagccattgg	attgctaata	agcgtgacag	cggttctagc	ggatttaggc	2220
gtgttgccgg	ttatacgagc	taaacaatg	attcgagatg	ataacgattc	agagactgga	2280
gaaaatctat	ggacatggca	attcacacct	tttcagaaac	tcaacttcac	agtcgaacat	2340
gtactcaagt	gtttggtaga	aggtaatgtt	atagggaag	gttgctccgg	gatatgttac	2400
aaagctgaaa	tgcttaacag	agaagtcac	gcggtgaaaa	agctctggcc	agtgacagta	2460
cctaacttga	atgagaaaac	taagtcatca	ggagttcgag	actcattctc	agctgaagta	2520
aaaacacttg	gatcgatcag	acacaagaac	attgtacgct	tcttgggatg	ttgttggaac	2580
aagaacacta	gacttcttat	gtatgattat	atgtcaaatg	ggagtttggg	aagtttgctt	2640
cacgaaagga	gcggtgtatg	tagcttagga	tgggaagtga	ggtacaagat	tatacttggt	2700
gcagctcagg	gtttggctta	cttgccacct	gactgtgttc	ctccccattg	tcatagagac	2760
atcaaggcaa	ataatattct	gattggccct	gattttgaac	cttacattgg	agatttcgga	2820
ctcgctaagc	ttgttgatga	tggcgacttt	gctcgttctt	ccaacacctt	tgctgggttc	2880
tatggttaca	tagctccagg	tacttcccaa	atttctgttt	tgctgctaata	tagttaacct	2940

crop01lpct.ST25.txt

gatacatcat aagttactag tagtcttagc ctagtaatgg tagaaattca ggtttatagt	3000
ttggttttgg tttttggcat ttgaaaaga atagtatttt ggttcagttg attttcgatt	3060
tggtttaaaa ttttagaaaa aaaatggttag aagaataat ttttgattaa tattatatca	3120
tctactgtat acgatttaaa aaaaaaattg atttggtttg gtctcgatat tttgcttgta	3180
gcttggttca atatttttac tatatatatg attcatgact ttcatactaa tcaatatttg	3240
ttcggtttcg ttttgattag tctgaaagtg aatcatatag ttaaaatatt gaccaatcta	3300
tcataactt acacgctaca agattttcga tgtaattttg gtgtaagcat tttggttttt	3360
atatgttcat ccttagtaac taacaacaaa tatcaattgg gaagcataac gttatatatt	3420
cccataaggc tataacacta taattatgca gaatacggat actcaatgaa gataacagag	3480
aaaagcgacg tgtacagcta tggagtcgta gtgctcgagg tactaacggg taagcaacca	3540
atcgatccaa cgataccaga tggactccac atagtggact gggccaagaa aatcagagac	3600
atacaagtaa tcgaccaagg actacaagca agaccagagt cagaggttga agagatgatg	3660
caaacgctag gagtcgcgt tttatgcatt aatccaatac ccgaagacag gcctacaatg	3720
aaagatgtgg ctgctatgct tagtgagata tgtcaagaga gagaggaatc gatgaaagtt	3780
gatggttgct cgggaagttg taacaatgga agagaacgtg gcaaaagatga ttcgacttca	3840
tcggttatgc aacaaacggc taagtatttg agaagtagta gcacgagttt ctctgcgtct	3900
tctttgcttt actctctctc ttcttctgct acttctaagt ttagacaaaa tcttaataaa	3960

&lt;210&gt; 4

&lt;211&gt; 1135

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 4

Met Ser Leu Gln Met Pro Ile Pro Arg Lys Lys Ala Leu Thr Val Ser
1 5 10 15

His Phe Ser Ile Thr Leu Ser Leu Phe Leu Ala Phe Phe Ile Ser Ser
20 25 30

Thr Ser Ala Ser Thr Asn Glu Val Ser Ala Leu Ile Ser Trp Leu His
35 40 45

Ser Ser Asn Ser Pro Pro Ser Val Phe Ser Gly Trp Asn Pro Ser
50 55 60

Asp Ser Asp Pro Cys Gln Trp Pro Tyr Ile Thr Cys Ser Ser Pro Asp
65 70 75 80

Asn Lys Leu Val Thr Glu Ile Asn Val Val Ser Val Gln Leu Ala Leu
---



crop011pct.ST25.txt

85	90	95
Pro Phe Pro Pro Asn Ile Ser Ser Phe Thr Ser Leu Gln Lys Leu Val		
100	105	110
Ile Ser Asn Thr Asn Leu Thr Gly Ala Ile Ser Ser Glu Ile Gly Asp		
115	120	125
Cys Ser Glu Leu Ile Val Ile Asp Leu Ser Ser Asn Ser Leu Val Gly		
130	135	140
Glu Ile Pro Ser Ser Leu Gly Lys Leu Lys Asn Leu Gln Glu Leu Cys		
145	150	155
Leu Asn Ser Asn Gly Leu Thr Gly Lys Ile Pro Pro Glu Leu Gly Asp		
165	170	175
Cys Val Ser Leu Lys Asn Leu Glu Ile Phe Asp Asn Tyr Leu Ser Glu		
180	185	190
Asn Leu Pro Leu Glu Leu Gly Lys Ile Ser Thr Leu Glu Ser Ile Arg		
195	200	205
Ala Gly Glu Asn Ser Glu Leu Ser Gly Lys Ile Pro Glu Glu Ile Gly		
210	215	220
Asn Cys Arg Asn Leu Lys Val Leu Gly Leu Ala Ala Thr Lys Ile Ser		
225	230	235
Gly Ser Leu Pro Val Ser Leu Gly Gln Leu Ser Lys Leu Gln Ser Leu		
245	250	255
Phe Val Tyr Ser Thr Met Leu Ser Gly Glu Ile Pro Lys Glu Leu Gly		
260	265	270
Asn Cys Ser Glu Leu Ile Asn Leu Phe Leu Tyr Asp Asn Asp Leu Ser		
275	280	285
Gly Thr Leu Pro Lys Glu Leu Gly Lys Leu Gln Asn Leu Glu Lys Met		
290	295	300
Leu Leu Trp Gln Asn Asn Leu His Gly Pro Ile Pro Glu Glu Ile Gly		
305	310	315
Phe Met Lys Ser Leu Asn Ala Ile Asp Leu Ser Met Asn Tyr Phe Ser		
325	330	335

crop011pct.ST25.txt

Gly Thr Ile Pro Lys Ser Phe Gly Asn Leu Ser Asn Leu Gln Glu Leu  
 340 345 350

Met Leu Ser Ser Asn Asn Ile Thr Gly Ser Ile Pro Ser Ile Leu Ser  
 355 360 365

Asp Cys Thr Lys Leu Val Gln Phe Gln Ile Asp Ala Asn Gln Ile Ser  
 370 375 380

Gly Leu Ile Pro Pro Glu Ile Gly Leu Leu Lys Glu Leu Asn Ile Phe  
 385 390 395 400

Leu Gly Trp Gln Asn Lys Leu Glu Gly Asn Ile Pro Asp Glu Leu Ala  
 405 410 415

Gly Cys Gln Asn Leu Gln Ala Leu Asp Leu Ser Gln Asn Tyr Leu Thr  
 420 425 430

Gly Ser Leu Pro Ala Gly Leu Phe Gln Leu Arg Asn Leu Thr Lys Leu  
 435 440 445

Leu Leu Ile Ser Asn Ala Ile Ser Gly Val Ile Pro Leu Glu Thr Gly  
 450 455 460

Asn Cys Thr Ser Leu Val Arg Leu Arg Leu Val Asn Asn Arg Ile Thr  
 465 470 475 480

Gly Glu Ile Pro Lys Gly Ile Gly Phe Leu Gln Asn Leu Ser Phe Leu  
 485 490 495

Asp Leu Ser Glu Asn Asn Leu Ser Gly Pro Val Pro Leu Glu Ile Ser  
 500 505 510

Asn Cys Arg Gln Leu Gln Met Leu Asn Leu Ser Asn Asn Thr Leu Gln  
 515 520 525

Gly Tyr Leu Pro Leu Ser Leu Ser Ser Leu Thr Lys Leu Gln Val Leu  
 530 535 540

Asp Val Ser Ser Asn Asp Leu Thr Gly Lys Ile Pro Asp Ser Leu Gly  
 545 550 555 560

His Leu Ile Ser Leu Asn Arg Leu Ile Leu Ser Lys Asn Ser Phe Asn  
 565 570 575

Gly Glu Ile Pro Ser Ser Leu Gly His Cys Thr Asn Leu Gln Leu Leu  
 580 585 590

## crop01lpct.ST25.txt

Asp Leu Ser Ser Asn Asn Ile Ser Gly Thr Ile Pro Glu Glu Leu Phe  
 595 600 605  
 Asp Ile Gln Asp Leu Asp Ile Ala Leu Asn Leu Ser Trp Asn Ser Leu  
 610 615 620  
 Asp Gly Phe Ile Pro Glu Arg Ile Ser Ala Leu Asn Arg Leu Ser Val  
 625 630 635 640  
 Leu Asp Ile Ser His Asn Met Leu Ser Gly Asp Leu Ser Ala Leu Ser  
 645 650 655  
 Gly Leu Glu Asn Leu Val Ser Leu Asn Ile Ser His Asn Arg Phe Ser  
 660 665 670  
 Gly Tyr Leu Pro Asp Ser Lys Val Phe Arg Gln Leu Ile Gly Ala Glu  
 675 680 685  
 Met Glu Gly Asn Asn Gly Leu Cys Ser Lys Gly Phe Arg Ser Cys Phe  
 690 695 700  
 Val Ser Asn Ser Ser Gln Leu Thr Thr Gln Arg Gly Val His Ser His  
 705 710 715 720  
 Arg Leu Arg Ile Ala Ile Gly Leu Leu Ile Ser Val Thr Ala Val Leu  
 725 730 735  
 Ala Val Leu Gly Val Leu Ala Val Ile Arg Ala Lys Gln Met Ile Arg  
 740 745 750  
 Asp Asp Asn Asp Ser Glu Thr Gly Glu Asn Leu Trp Thr Trp Gln Phe  
 755 760 765  
 Thr Pro Phe Gln Lys Leu Asn Phe Thr Val Glu His Val Leu Lys Cys  
 770 775 780  
 Leu Val Glu Gly Asn Val Ile Gly Lys Gly Cys Ser Gly Ile Val Tyr  
 785 790 795 800  
 Lys Ala Glu Met Pro Asn Arg Glu Val Ile Ala Val Lys Lys Leu Trp  
 805 810 815  
 Pro Val Thr Val Pro Asn Leu Asn Glu Lys Thr Lys Ser Ser Gly Val  
 820 825 830  
 Arg Asp Ser Phe Ser Ala Glu Val Lys Thr Leu Gly Ser Ile Arg His  
 835 840 845

crop011pct.ST25.txt

Lys Asn Ile Val Arg Phe Leu Gly Cys Cys Trp Asn Lys Asn Thr Arg  
 850 855 860  
 Leu Leu Met Tyr Asp Tyr Met Ser Asn Gly Ser Leu Gly Ser Leu Leu  
 865 870 875 880  
 His Glu Arg Ser Gly Val Cys Ser Leu Gly Trp Glu Val Arg Tyr Lys  
 885 890 895  
 Ile Ile Leu Gly Ala Ala Gln Gly Leu Ala Tyr Leu His His Asp Cys  
 900 905 910  
 Val Pro Pro Ile Val His Arg Asp Ile Lys Ala Asn Asn Ile Leu Ile  
 915 920 925  
 Gly Pro Asp Phe Glu Pro Tyr Ile Gly Asp Phe Gly Leu Ala Lys Leu  
 930 935 940  
 Val Asp Asp Gly Asp Phe Ala Arg Ser Ser Asn Thr Ile Ala Gly Ser  
 945 950 955 960  
 Tyr Gly Tyr Ile Ala Pro Glu Tyr Gly Tyr Ser Met Lys Ile Thr Glu  
 965 970 975  
 Lys Ser Asp Val Tyr Ser Tyr Gly Val Val Val Leu Glu Val Leu Thr  
 980 985 990  
 Gly Lys Gln Pro Ile Asp Pro Thr Ile Pro Asp Gly Leu His Ile Val  
 995 1000 1005  
 Asp Trp Val Lys Lys Ile Arg Asp Ile Gln Val Ile Asp Gln Gly  
 1010 1015 1020  
 Leu Gln Ala Arg Pro Glu Ser Glu Val Glu Glu Met Met Gln Thr  
 1025 1030 1035  
 Leu Gly Val Ala Leu Leu Cys Ile Asn Pro Ile Pro Glu Asp Arg  
 1040 1045 1050  
 Pro Thr Met Lys Asp Val Ala Ala Met Leu Ser Glu Ile Cys Gln  
 1055 1060 1065  
 Glu Arg Glu Glu Ser Met Lys Val Asp Gly Cys Ser Gly Ser Cys  
 1070 1075 1080  
 Asn Asn Gly Arg Glu Arg Gly Lys Asp Asp Ser Thr Ser Ser Val  
 Page 13

crop011pct.ST25.txt

1085	1090	1095
------	------	------

Met Gln Gln Thr Ala Lys Tyr Leu Arg Ser Ser Ser Thr Ser Phe  
 1100 1105 1110

Ser Ala Ser Ser Leu Leu Tyr Ser Ser Ser Ser Ser Ala Thr Ser  
 1115 1120 1125

Asn Val Arg Pro Asn Leu Lys  
 1130 1135

<210> 5  
 <211> 25  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 5

Arg Tyr Lys Ile Ile Leu Gly Ala Ala Gln Gly Leu Ala Tyr Leu His  
 1 5 10 15

His Asp Cys Val Pro Pro Ile Val His  
 20 25

<210> 6  
 <211> 23  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 6  
 aayaayttma syggtkmdwt hcc 23

<210> 7  
 <211> 20  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 7  
 carythamcg gwgaaathcc 20

<210> 8  
 <211> 20  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 8  
 mayytsavyg gahctathcc 20

<210> 9  
 <211> 22  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>

crop01lpct.ST25.txt

<221> misc\_feature  
 <222> {17}..{17}  
 <223> n = a, c, g or t

<400> 9  
 thccwtwgs catgtantca ta 22

<210> 10  
 <211> 19  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 10  
 aytcdggdgc hahgtadcc 19

<210> 11  
 <211> 21  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 11  
 mayhccraar ctrtavacat c 21

<210> 12  
 <211> 19  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 12  
 cgatcagaca caagaacct 19

<210> 13  
 <211> 19  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 13  
 agcaatggtg ttggaagaa 19

<210> 14  
 <211> 19  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 14  
 tgcagatctt ygtgaagac 19

<210> 15  
 <211> 18  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 15  
 gactccttct ggatgttg 18

## crop011pct.ST25.txt

<210> 16  
 <211> 18  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 16  
 tgtaaaacga cggccagt

18

<210> 17  
 <211> 34  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 17  
 cacacaggat ccaatcggca ttgcaaaga cata

34

<210> 18  
 <211> 842  
 <212> DNA  
 <213> Arabidopsis thaliana

<400> 18  
 aagcttaaat gacaatttag taccttgggt tggatcatgat ttagagcggg acaaatatac 60  
 catacatcaa acgaggatat acagagaaaa ttcattggaag tatggaattt agaggacaat 120  
 ttctcttctg ggtacaaacg gaccggccca ttgctcatt taccagagg tatcgagttt 180  
 gtggactttt gatgcgcta gagactattg gcatcggatt gaaaaaatg ttacttcgt 240  
 tgtaacaat ttctgaatg caatatttct cttgtcatga atatttaaac ttgttattac 300  
 ttctttttag cttagggtg gacaattatg gagtttactt caaacgagga agaattctaa 360  
 acgctcgggt cagggtctga aaacaaacca actcacaatc ctgacttaat tgaggaaaac 420  
 aatgcaaac cactgcatg ctcccatatt tctatcataa tcttataaga aaaacacta 480  
 ctaagtgaat tgattctgta tatatataac caatgccttt tgttttgta tattttatgt 540  
 atatataact attgactttt gtcactatg gatagtgtct cgggtctctg gcaaacatat 600  
 ttcaaagaaa agttaatgac tgtaattaat taatctgaag ctgaaaacag aaccccgagg 660  
 taaaagaaaa agacagagca catgaagttt agtactttta tatatttaat atactattct 720  
 ttcttattgc ttatctctaa agcaaaaact tccctaaacc ctaagccaaa ggactcagat 780  
 cgatgcagaa ccaagaaggc ttgttttggg ttgagagcc aaatgcaag aaaaaaactc 840  
 tt 842

<210> 19  
 <211> 1130  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 19

Met Pro Ile Pro Arg Lys Lys Ala Leu Thr Val Ser His Phe Ser Ile

crop011pct.ST25.txt

1                      5                      10                      15

Thr Leu Ser Leu Phe Leu Ala Phe Phe Ile Ser Ser Thr Ser Ala Ser  
20                      25                      30

Thr Asn Glu Val Ser Ala Leu Ile Ser Trp Leu His Ser Ser Asn Ser  
35                      40                      45

Pro Pro Pro Ser Val Phe Ser Gly Trp Asn Pro Ser Asp Ser Asp Pro  
50                      55                      60

Cys Gln Trp Pro Tyr Ile Thr Cys Ser Ser Pro Asp Asn Lys Leu Val  
65                      70                      75                      80

Thr Glu Ile Asn Val Val Ser Val Gln Leu Ala Leu Pro Phe Pro Pro  
85                      90                      95

Asn Ile Ser Ser Phe Thr Ser Leu Gln Lys Leu Val Ile Ser Asn Thr  
100                      105                      110

Asn Leu Thr Gly Ala Ile Ser Ser Glu Ile Gly Asp Cys Ser Glu Leu  
115                      120                      125

Ile Val Ile Asp Leu Ser Ser Asn Ser Leu Val Gly Glu Ile Pro Ser  
130                      135                      140

Ser Leu Gly Lys Leu Lys Asn Leu Gln Glu Leu Cys Leu Asn Ser Asn  
145                      150                      155                      160

Gly Leu Thr Gly Lys Ile Pro Pro Glu Leu Gly Asp Cys Val Ser Leu  
165                      170                      175

Lys Asn Leu Glu Ile Phe Asp Asn Tyr Leu Ser Glu Asn Leu Pro Leu  
180                      185                      190

Glu Leu Gly Lys Ile Ser Thr Leu Glu Ser Ile Arg Ala Gly Gly Asn  
195                      200                      205

Ser Glu Leu Ser Gly Lys Ile Pro Glu Glu Ile Gly Asn Cys Arg Asn  
210                      215                      220

Leu Lys Val Leu Gly Leu Ala Ala Thr Lys Ile Ser Gly Ser Leu Pro  
225                      230                      235                      240

Val Ser Leu Gly Gln Leu Ser Lys Leu Gln Ser Leu Phe Val Tyr Ser  
245                      250                      255



crop011pct.ST25.txt

Thr Ser Leu Ser Gly Glu Ile Pro Lys Glu Leu Gly Asn Cys Ser Glu  
260 265 270

Leu Ile Asn Leu Phe Leu Tyr Asp Asn Asp Leu Ser Gly Thr Leu Pro  
280 285

Lys Val Leu Gly Lys Leu Gln Asn Leu Glu Lys Met Leu Leu Trp Gln  
295 300

Asn Asp Leu His Gly Pro Ile Pro Glu Glu Ile Gly Phe Met Lys Ser  
305 310 315 320

Leu Asn Ala Ile Asp Leu Ser Met Asn Tyr Phe Ser Gly Thr Ile Pro  
325 330 335

Lys Ser Ile Gly Asn Leu Ser Asn Leu Gln Glu Leu Met Leu Ser Ser  
340 345 350

Asn Asn Ile Thr Gly Ser Ile Pro Ser Ile Leu Ser Asp Cys Thr Lys  
365 370 375

Leu Val Gln Phe Gln Ile Asp Ala Asn Gln Ile Ser Gly Leu Ile Pro  
380 385 390 395

Pro Ile Ile Gly Leu Leu Lys Glu Leu Asn Ile Phe Gly Trp Gln Asn  
395 400 405

Lys Leu Glu Gly Asn Ile Pro Asp Glu Leu Ala Gly Cys Gln Asn Leu  
410 415 420

Gln Ala Leu Asp Leu Ser Gln Asn Tyr Leu Thr Gly Ser Leu Pro Ala  
425 430 435

Gly Leu Phe Gln Leu Arg Asn Leu Thr Lys Leu Leu Ile Ser Asn  
440 445 450

Ala Ile Ser Gly Val Ile Pro Leu Glu Thr Gly Asn Cys Thr Ser Leu  
455 460 465

Val Arg Leu Arg Leu Val Asn Asn Arg Ile Thr Gly Glu Ile Pro Lys  
470 475 480

Gly Ile Gly Phe Leu Gln Asn Leu Ser Phe Leu Asp Leu Ser Glu Asn  
485 490 495

Asn Leu Ser Gly Pro Val Pro Leu Glu Ile Ser Asn Cys Arg Gln Leu  
500 505 510

## crop011pct.ST25.txt

Gln Met Leu Asn Leu Ser Asn Asn Thr Leu Gln Gly Tyr Leu Pro Leu  
 515 520 525  
 Ser Leu Ser Ser Leu Thr Lys Leu Gln Val Leu Asp Val Ser Ser Asn  
 530 535 540  
 Asp Leu Thr Gly Lys Ile Pro Asp Ser Leu Gly His Leu Ile Ser Leu  
 545 550 555 560  
 Asn Arg Leu Ile Leu Ser Lys Asn Ser Phe Asn Gly Glu Ile Pro Ser  
 565 570 575  
 Ser Leu Gly His Cys Thr Asn Leu Gln Leu Leu Asp Leu Ser Ser Asn  
 580 585 590  
 Asn Ile Ser Gly Thr Ile Pro Glu Glu Leu Phe Asp Ile Gln Asp Leu  
 595 600 605  
 Asp Ile Ala Leu Asn Leu Ser Trp Asn Ser Leu Asp Gly Phe Ile Pro  
 610 615 620  
 Glu Arg Ile Ser Ala Leu Asn Arg Leu Ser Val Leu Asp Ile Ser His  
 625 630 635 640  
 Asn Met Leu Ser Gly Asp Leu Ser Ala Leu Ser Gly Leu Glu Asn Leu  
 645 650 655  
 Val Ser Leu Asn Ile Ser His Asn Arg Phe Ser Gly Tyr Leu Pro Asp  
 660 665 670  
 Ser Lys Val Phe Arg Gln Leu Ile Gly Ala Glu Met Glu Gly Asn Asn  
 675 680 685  
 Gly Leu Cys Ser Lys Gly Phe Arg Ser Cys Phe Val Ser Asn Ser Ser  
 690 695 700  
 Gln Leu Thr Thr Gln Arg Gly Val His Ser His Arg Leu Arg Ile Ala  
 705 710 715 720  
 Ile Gly Leu Leu Ile Ser Val Thr Ala Val Leu Ala Val Leu Gly Val  
 725 730 735  
 Leu Ala Val Ile Arg Ala Lys Gln Met Ile Arg Asp Asp Asn Asp Ser  
 740 745 750  
 Glu Thr Gly Glu Asn Leu Trp Thr Trp Gln Phe Thr Pro Phe Gln Lys  
 755 760 765

crop011pct.ST25.txt

Leu Asn Phe Thr Val Glu His Val Leu Lys Cys Leu Val Glu Gly Asn  
 770 775 780

Val Ile Gly Lys Gly Cys Ser Gly Ile Val Tyr Lys Ala Glu Met Pro  
 785 790 795 800

Asn Arg Glu Val Ile Ala Val Lys Lys Leu Trp Pro Val Thr Val Pro  
 805 810 815

Asn Leu Asn Glu Lys Thr Lys Ser Ser Gly Val Arg Asp Ser Phe Ser  
 820 825 830

Ala Glu Val Lys Thr Leu Gly Ser Ile Arg His Lys Asn Ile Val Arg  
 835 840 845

Phe Leu Gly Cys Cys Trp Asn Lys Asn Thr Arg Leu Leu Met Tyr Asp  
 850 855 860

Tyr Met Ser Asn Gly Ser Leu Gly Ser Leu Leu His Glu Arg Ser Gly  
 865 870 875 880

Val Cys Ser Leu Gly Trp Glu Val Arg Tyr Lys Ile Ile Leu Gly Ala  
 885 890 895

Ala Gln Gly Leu Ala Tyr Leu His His Asp Cys Val Pro Pro Ile Val  
 900 905 910

His Arg Asp Ile Lys Ala Asn Asn Ile Leu Ile Gly Pro Asp Phe Glu  
 915 920 925

Pro Tyr Ile Gly Asp Phe Gly Leu Ala Lys Leu Val Asp Asp Gly Asp  
 930 935 940

Phe Ala Arg Ser Ser Asn Thr Ile Ala Gly Ser Tyr Gly Tyr Ile Ala  
 945 950 955 960

Pro Glu Tyr Gly Tyr Ser Met Lys Ile Thr Glu Lys Ser Asp Val Tyr  
 965 970 975

Ser Tyr Gly Val Val Val Leu Glu Val Leu Thr Gly Lys Gln Pro Ile  
 980 985 990

Asp Pro Thr Ile Pro Asp Gly Leu His Ile Val Asp Trp Val Lys Lys  
 995 1000 1005

Ile Arg Asp Ile Gln Val Ile Asp Gln Gly Leu Gln Ala Arg Pro

crop011pct.ST25.txt

1010

1015

1020

Glu Ser Glu Val Glu Glu Met Met Gln Thr Leu Gly Val Ala Leu  
 1025 1030 1035

Leu Cys Ile Asn Pro Ile Pro Glu Asp Arg Pro Thr Met Lys Asp  
 1040 1045 1050

Val Ala Ala Met Leu Ser Glu Ile Cys Gln Glu Arg Glu Glu Ser  
 1055 1060 1065

Met Lys Val Asp Gly Cys Ser Gly Ser Cys Asn Asn Gly Arg Glu  
 1070 1075 1080

Arg Gly Lys Asp Asp Ser Thr Ser Ser Val Met Gln Gln Thr Ala  
 1085 1090 1095

Lys Tyr Leu Arg Ser Ser Ser Thr Ser Phe Ser Ala Ser Ser Leu  
 1100 1105 1110

Leu Thr Ser Ser Ser Ser Ser Ala Thr Ser Asn Val Arg Pro Asn  
 1115 1120 1125

Leu Lys  
 1130